OMEGA ACADEMY, NUMERICAL METHODS COURSE.

Erika Jissel Gutiérrez Beltrán

Daniel Fernandez Delgado

Frank Edward Daza González

Johanna Arias

Freddy Sebastian Garcia

Teacher:

Walter German Magaña

Matter:

Numerical Methods

Universidad de San Buenaventura Cali 2014



UNIT EIGHT

Polynomials roots.

The roots of a polynomial are the values that override the polynomial; a polynomial is a mathematical expression consisting of a finite set of variables and constants not determined. The functions or equations have a degree; the degree is the highest exponent of the equation.

Example I.

(Function Second Grade).

$$f(x) = x^2 - 5x + 6$$

Factorized

$$f(x) = (x-2)(x-3)$$

Then the solutions or roots of the polynomial are

$$x_0 = 2 y x_1 = 3$$

Replacing x0 and x1 will check the function

$$f(2) = 2^2 - 5(2) + 6$$

$$f(2) = 4 - 10 + 6$$

$$f(2)=0$$



$$f(3) = 3^{2} - 5(3) + 6$$
$$f(3) = 9 - 15 + 6$$
$$f(3) = 0$$

Example 2 (Function Fourth Grade):

$$f(x) = x^4 - 11x^3 + 41x^2 - 61x + 30$$

Synthetic division is made; the coefficient of the variables are taken



$$f(x) = (x^2 - 3x + 2)(x - 3)(x - 5)$$

$$f(x) = (x-1)(x-2)(x-3)(x-5)$$

The solutions or roots of the polynomial are:

$$x_0 = 1, x_1 = 2, x_2 = 3 \text{ y } x_3 = 5$$

It is found by replacing the x in the function

$$f(1) = (1)^4 - 11(1)^3 + 41(1)^2 - 61(1) + 30$$
$$f(1) = 1 - 11 + 42 - 61 + 30$$
$$f(1) = 72 - 72$$
$$f(1) = 0$$

$$f(2) = (2)^4 - 11(2)^3 + 41(2)^2 - 61(2) + 30$$
$$f(2) = 16 - 88 + 164 - 122 + 30$$
$$f(2) = 210 - 210$$
$$f(2) = 0$$

$$f(3) = (3)^4 - 11(3)^3 + 41(3)^2 - 61(3) + 30$$
$$f(3) = 81 - 297 + 369 - 183 + 30$$
$$f(3) = 480 - 480$$
$$f(3) = 0$$



$$f(5) = (5)^4 - 11(5)^3 + 41(5)^2 - 61(5) + 30$$
$$f(5) = 625 - 1375 + 1025 - 305 + 30$$
$$f(4) = 1680 - 1680$$
$$f(4) = 0$$

