Table S9. Model summaries for the relationship between body mass and cat effects on prey abundance. Model estimates ± 95% CIs and test statistics are reported along with sample sizes and residual unexplained heterogeneity (I^2), decomposed by hierarchical model levels (article and observation). The effect size used is given in the heading of each model (SMD=standardized mean difference or Hedges' g; Zr=correlation coefficient; lnOR=log odds ratio).

Term	Estimate	Test statistics
Abundance with/without cats | lnOR $N_{articles} = 9$; $N_{species} = 11$; $N_{observations} = 11$ $I^2_{total} = 65.2$; $I^2_{article} = 65.2$; $I^2_{obs} = 0$		
intercept	-0.23 ±[-5.18, 4.72]	$t_7 = -0.11, p = 0.91$
Mass (g, log10)	-0.18 ±[-2.26, 1.91]	$t_9 = -0.19, p = 0.85$
Abundance on islands with/without cats | lnOR $N_{articles} = 6$; $N_{species} = 8$; $N_{observations} = 8$ $I^2_{total} = 49$; $I^2_{article} = 49$; $I^2_{obs} = 0$		
intercept	1.13 ±[-5.23, 7.5]	$t_4 = 0.5, p = 0.65$
Mass (g, log10)	-0.42 ±[-2.73, 1.88]	$t_6 = -0.45, p = 0.67$
Abundance association | Zr $N_{articles} = 15$; $N_{species} = 14$; $N_{observations} = 23$ $I^2_{total} = 83.8$; $I^2_{article} = 59.4$; $I^2_{obs} = 24$		
intercept	0.4 ±[-1.53, 2.33]	$t_{13} = 0.45, p = 0.66$
Mass (g, log10)	-0.12 ±[-0.8, 0.55]	$t_{21} = -0.38, p = 0.71$
Abundance temporal association | Zr $N_{articles} = 11$; $N_{species} = 13$; $N_{observations} = 19$ $I^2_{total} = 86.4$; $I^2_{article} = 62.3$; $I^2_{obs} = 24$		
intercept	-0.1 ±[-3.9, 3.7]	$t_9 = -0.06, p = 0.95$
Mass (g, log10)	0.06 ±[-1.16, 1.27]	$t_{17} = 0.1, p = 0.92$