Table SX. Model summaries for the relationship between body mass and cat effects on prey abundance. Model estimates ± 95% CIs and test statistics are reported along with sample sizes and residual unexplained heterogeneity (I^2), decomposed by hierarchical model levels (article and observation). The effect size used is given in the heading of each model (SMD=standardized mean difference or Hedges' g; Zr=correlation coefficient; lnOR=log odds ratio).

Term	Estimate	Test statistics
Abundance with/without cats | SMD | $N_{articles} = 10$; $N_{species} = 13$; $N_{observations} = 15$ | $I^2_{total} = 59.2$; $I^2_{article} = 59.2$; $I^2_{obs} = 0$		
intercept	-0.12 ±[-2.45, 2.21]	$t_8 = -0.12, p = 0.91$
Mass (g, log10)	-0.15 ±[-1.11, 0.81]	$t_{13} = -0.34, p = 0.74$
Abundance on islands with/without cats | lnOR | $N_{articles} = 7$; $N_{species} = 11$; $N_{observations} = 11$ | $I^2_{total} = 42.1$; $I^2_{article} = 42.1$; $I^2_{obs} = 0$		
intercept	0.63 ±[-4.32, 5.58]	$t_5 = 0.33, p = 0.76$
Mass (g, log10)	-0.34 ±[-2.15, 1.48]	$t_9 = -0.42, p = 0.69$
Abundance temporal association | Zr | $N_{articles} = 11$; $N_{species} = 13$; $N_{observations} = 19$ | $I^2_{total} = 86.4$; $I^2_{article} = 62.3$; $I^2_{obs} = 24$		
intercept	-0.1 ±[-3.9, 3.7]	$t_9 = -0.06, p = 0.95$
Mass (g, log10)	0.06 ±[-1.16, 1.27]	$t_{17} = 0.1, p = 0.92$