

Table SX. Model summaries for intercept-only (main text) models as well as models evaluating influence of habitat, body mass, and continents versus islands. Model estimates $\pm$ 95% CIs and test statistics are reported along with sample sizes and residual unexplained heterogeneity (I^2), decomposed by hierarchical model levels (article and observation). The effect size used is given in the heading of each model (SMD=standardized mean difference or Hedges' g; Zr=correlation coefficient; lnOR=log odds ratio).

Term	Estimate	Moderator	Test statistics	Sample size	Residual heterogeneity
Abundance temporal association All species Continent vs island Zr					
island	-0.59 $\pm$ [-1.48, 0.29]	Continent vs island	$t_9 = -1.51, p = 0.16$	$N_{articles} = 3$ $N_{species} = 4$ $N_{observations} = 6$	$I^2_{total} = 82.8$ $I^2_{article} = 56.7$ $I^2_{obs} = 26$
mainland	0.3 $\pm$ [-0.22, 0.83]	Continent vs island	$t_9 = 1.3, p = 0.22$	$N_{articles} = 8$ $N_{species} = 9$ $N_{observations} = 13$	$I^2_{total} = 82.8$ $I^2_{article} = 56.7$ $I^2_{obs} = 26$
Abundance on islands with/without cats All species Ground foraging lnOR					
ground	0.07 $\pm$ [-2.04, 2.17]	Ground foraging	$t_9 = 0.07, p = 0.94$	$N_{articles} = 4$ $N_{species} = 4$ $N_{observations} = 4$	$I^2_{total} = 43.4$ $I^2_{article} = 43.4$ $I^2_{obs} = 0$
other	-0.26 $\pm$ [-2.08, 1.55]	Ground foraging	$t_9 = -0.33, p = 0.75$	$N_{articles} = 4$ $N_{species} = 7$ $N_{observations} = 7$	$I^2_{total} = 43.4$ $I^2_{article} = 43.4$ $I^2_{obs} = 0$
Abundance temporal association All species Ground foraging Zr					
ground	-0.81 $\pm$ [-1.71, 0.09]	Ground foraging	$t_{17} = -1.9, p = 0.07$	$N_{articles} = 3$ $N_{species} = 3$ $N_{observations} = 3$	$I^2_{total} = 81.6$ $I^2_{article} = 59.9$ $I^2_{obs} = 22$
other	0.23 $\pm$ [-0.21, 0.67]	Ground foraging	$t_{17} = 1.12, p = 0.28$	$N_{articles} = 9$ $N_{species} = 10$ $N_{observations} = 16$	$I^2_{total} = 81.6$ $I^2_{article} = 59.9$ $I^2_{obs} = 22$
Abundance on islands with/without cats All species Ground nesting/resting lnOR					
ground	0.62 $\pm$ [-1.22, 2.45]	Ground nesting/resting	$t_9 = 0.76, p = 0.47$	$N_{articles} = 3$ $N_{species} = 6$ $N_{observations} = 6$	$I^2_{total} = 23.7$ $I^2_{article} = 23.7$ $I^2_{obs} = 0$
other	-0.69 $\pm$ [-2.13, 0.74]	Ground nesting/resting	$t_9 = -1.09, p = 0.3$	$N_{articles} = 5$ $N_{species} = 5$ $N_{observations} = 5$	$I^2_{total} = 23.7$ $I^2_{article} = 23.7$ $I^2_{obs} = 0$
Abundance temporal association All species Ground nesting/resting Zr					

Table SX. Model summaries for intercept-only (main text) models as well as models evaluating influence of habitat, body mass, and continents versus islands. Model estimates $\pm$ 95% CIs and test statistics are reported along with sample sizes and residual unexplained heterogeneity (I^2), decomposed by hierarchical model levels (article and observation). The effect size used is given in the heading of each model (SMD=standardized mean difference or Hedges' g; Zr=correlation coefficient; lnOR=log odds ratio).

Term	Estimate	Moderator	Test statistics	Sample size	Residual heterogeneity
ground	-0.36 $\pm$ [-1.27, 0.55]	Ground nesting/resting	$t_{17} = -0.84, p = 0.41$	$N_{articles} = 3$ $N_{species} = 3$ $N_{observations} = 4$	$I^2_{total} = 86.1$ $I^2_{article} = 66.6$ $I^2_{obs} = 20$
other	0.16 $\pm$ [-0.35, 0.67]	Ground nesting/resting	$t_{17} = 0.66, p = 0.52$	$N_{articles} = 9$ $N_{species} = 10$ $N_{observations} = 15$	$I^2_{total} = 86.1$ $I^2_{article} = 66.6$ $I^2_{obs} = 20$
Abundance with/without cats All species Volant SMD					
non_volant	-0.32 $\pm$ [-1.2, 0.55]	Volant	$t_8 = -0.86, p = 0.42$	$N_{articles} = 6$ $N_{species} = 8$ $N_{observations} = 8$	$I^2_{total} = 57.8$ $I^2_{article} = 57.8$ $I^2_{obs} = 0$
volant	-0.58 $\pm$ [-1.63, 0.48]	Volant	$t_8 = -1.26, p = 0.24$	$N_{articles} = 4$ $N_{species} = 5$ $N_{observations} = 7$	$I^2_{total} = 57.8$ $I^2_{article} = 57.8$ $I^2_{obs} = 0$
Abundance on islands with/without cats All species Volant lnOR					
non_volant	0.16 $\pm$ [-1.88, 2.21]	Volant	$t_5 = 0.2, p = 0.85$	$N_{articles} = 4$ $N_{species} = 6$ $N_{observations} = 6$	$I^2_{total} = 39.7$ $I^2_{article} = 39.7$ $I^2_{obs} = 0$
volant	-0.56 $\pm$ [-2.99, 1.86]	Volant	$t_5 = -0.6, p = 0.58$	$N_{articles} = 3$ $N_{species} = 5$ $N_{observations} = 5$	$I^2_{total} = 39.7$ $I^2_{article} = 39.7$ $I^2_{obs} = 0$