

Table S10. Model summaries for habitat specific groupings, including foraging habitat (ground versus other), nesting habitat (ground versus other), locomotion (volant versus non-volant) and landform (continent vs island). Each habitat factor level was run independently in a subgroup intercept-only model. Model estimates $\pm$ 95% CIs and test statistics are reported along with sample sizes and residual unexplained heterogeneity (I^2), decomposed by hierarchical model levels (article and observation). The effect size used is given in the heading of each model (SMD=standardized mean difference or Hedges' g; Zr=correlation coefficient; lnOR=log odds ratio).

Term	Estimate	Moderator	Test statistics	Sample size	Residual heterogeneity
Abundance with/without cats Locomotion: volant lnOR					
overall	-0.72 $\pm$ [-8.12, 6.68]	Overall estimate	$t_2 = -0.42, p = 0.72$	$N_{articles} = 3$ $N_{species} = 3$ $N_{observations} = 3$	$I^2_{total} = 77.5$ $I^2_{article} = 38.7$ $I^2_{obs} = 39$
Abundance with/without cats Locomotion: Non-volant lnOR					
overall	-0.58 $\pm$ [-2.37, 1.21]	Overall estimate	$t_5 = -0.83, p = 0.44$	$N_{articles} = 6$ $N_{species} = 8$ $N_{observations} = 8$	$I^2_{total} = 58.2$ $I^2_{article} = 58.2$ $I^2_{obs} = 0$
Abundance with/without cats Foraging habitat: ground lnOR					
overall	0.27 $\pm$ [-3.57, 4.1]	Overall estimate	$t_3 = 0.22, p = 0.84$	$N_{articles} = 4$ $N_{species} = 4$ $N_{observations} = 4$	$I^2_{total} = 63.7$ $I^2_{article} = 31.9$ $I^2_{obs} = 32$
Abundance with/without cats Foraging habitat: other lnOR					
overall	-0.91 $\pm$ [-2.72, 0.9]	Overall estimate	$t_5 = -1.29, p = 0.25$	$N_{articles} = 6$ $N_{species} = 7$ $N_{observations} = 7$	$I^2_{total} = 59.2$ $I^2_{article} = 52.9$ $I^2_{obs} = 6$
Abundance with/without cats Ground nesting: ground lnOR					
overall	1.75 $\pm$ [-9.35, 12.85]	Overall estimate	$t_1 = 2, p = 0.29$	$N_{articles} = 2$ $N_{species} = 4$ $N_{observations} = 4$	$I^2_{total} = 0$ $I^2_{article} = 0$ $I^2_{obs} = 0$
Abundance with/without cats Ground nesting: other lnOR					
overall	-1.38 $\pm$ [-2.68, -0.08]	Overall estimate	$t_6 = -2.6, p = 0.04$	$N_{articles} = 7$ $N_{species} = 7$ $N_{observations} = 7$	$I^2_{total} = 43.2$ $I^2_{article} = 21.6$ $I^2_{obs} = 22$
Abundance with/without cats Landform: island lnOR					
overall	-0.27 $\pm$ [-1.64, 1.09]	Overall estimate	$t_7 = -0.47, p = 0.65$	$N_{articles} = 8$ $N_{species} = 10$ $N_{observations} = 10$	$I^2_{total} = 39.3$ $I^2_{article} = 39.3$ $I^2_{obs} = 0$

Table S10. Model summaries for habitat specific groupings, including foraging habitat (ground versus other), nesting habitat (ground versus other), locomotion (volant versus non-volant) and landform (continent vs island). Each habitat factor level was run independently in a subgroup intercept-only model. Model estimates $\pm$ 95% CIs and test statistics are reported along with sample sizes and residual unexplained heterogeneity (I^2), decomposed by hierarchical model levels (article and observation). The effect size used is given in the heading of each model (SMD=standardized mean difference or Hedges' g; Zr=correlation coefficient; lnOR=log odds ratio).

Term	Estimate	Moderator	Test statistics	Sample size	Residual heterogeneity
Abundance on islands with/without cats Locomotion: volant lnOR					
overall	0.39 $\pm$ [-30.5, 31.27]	Overall estimate	$t_1 = 0.16, p = 0.9$	$N_{articles} = 2$ $N_{species} = 2$ $N_{observations} = 2$	$I^2_{total} = 82.4$ $I^2_{article} = 41.2$ $I^2_{obs} = 41$
Abundance on islands with/without cats Locomotion: Non-volant lnOR					
overall	0.08 $\pm$ [-1.82, 1.99]	Overall estimate	$t_3 = 0.14, p = 0.9$	$N_{articles} = 4$ $N_{species} = 6$ $N_{observations} = 6$	$I^2_{total} = 13.3$ $I^2_{article} = 13.3$ $I^2_{obs} = 0$
Abundance on islands with/without cats Foraging habitat: ground lnOR					
overall	0.27 $\pm$ [-3.57, 4.1]	Overall estimate	$t_3 = 0.22, p = 0.84$	$N_{articles} = 4$ $N_{species} = 4$ $N_{observations} = 4$	$I^2_{total} = 63.7$ $I^2_{article} = 31.9$ $I^2_{obs} = 32$
Abundance on islands with/without cats Foraging habitat: other lnOR					
overall	0.16 $\pm$ [-2.72, 3.04]	Overall estimate	$t_2 = 0.23, p = 0.84$	$N_{articles} = 3$ $N_{species} = 4$ $N_{observations} = 4$	$I^2_{total} = 10.2$ $I^2_{article} = 0$ $I^2_{obs} = 10$
Abundance on islands with/without cats Ground nesting: ground lnOR					
overall	1.75 $\pm$ [-9.35, 12.85]	Overall estimate	$t_1 = 2, p = 0.29$	$N_{articles} = 2$ $N_{species} = 4$ $N_{observations} = 4$	$I^2_{total} = 0$ $I^2_{article} = 0$ $I^2_{obs} = 0$
Abundance on islands with/without cats Ground nesting: other lnOR					
overall	-0.69 $\pm$ [-2.37, 1]	Overall estimate	$t_3 = -1.3, p = 0.28$	$N_{articles} = 4$ $N_{species} = 4$ $N_{observations} = 4$	$I^2_{total} = 0$ $I^2_{article} = 0$ $I^2_{obs} = 0$
Abundance on islands with/without cats Landform: island lnOR					
overall	0.12 $\pm$ [-1.59, 1.84]	Overall estimate	$t_5 = 0.18, p = 0.86$	$N_{articles} = 6$ $N_{species} = 8$ $N_{observations} = 8$	$I^2_{total} = 39.6$ $I^2_{article} = 39.6$ $I^2_{obs} = 0$
Abundance before-after eradication Foraging habitat: other SMD					

Table S10. Model summaries for habitat specific groupings, including foraging habitat (ground versus other), nesting habitat (ground versus other), locomotion (volant versus non-volant) and landform (continent vs island). Each habitat factor level was run independently in a subgroup intercept-only model. Model estimates $\pm$ 95% CIs and test statistics are reported along with sample sizes and residual unexplained heterogeneity (I^2), decomposed by hierarchical model levels (article and observation). The effect size used is given in the heading of each model (SMD=standardized mean difference or Hedges' g; Zr=correlation coefficient; lnOR=log odds ratio).

Term	Estimate	Moderator	Test statistics	Sample size	Residual heterogeneity
overall	-0.75 $\pm$ [-7.93, 6.43]	Overall estimate	$t_1 = -1.33, p = 0.41$	$N_{articles} = 2$ $N_{species} = 2$ $N_{observations} = 2$	$I^2_{total} = 18.8$ $I^2_{article} = 9.4$ $I^2_{obs} = 9$
Abundance before-after eradication Ground nesting: other SMD					
overall	-0.75 $\pm$ [-7.93, 6.43]	Overall estimate	$t_1 = -1.33, p = 0.41$	$N_{articles} = 2$ $N_{species} = 2$ $N_{observations} = 2$	$I^2_{total} = 18.8$ $I^2_{article} = 9.4$ $I^2_{obs} = 9$
Abundance before-after eradication Landform: island SMD					
overall	-0.75 $\pm$ [-7.93, 6.43]	Overall estimate	$t_1 = -1.33, p = 0.41$	$N_{articles} = 2$ $N_{species} = 2$ $N_{observations} = 2$	$I^2_{total} = 18.8$ $I^2_{article} = 9.4$ $I^2_{obs} = 9$
Abundance association Locomotion: volant Zr					
overall	-0.57 $\pm$ [-8.52, 7.38]	Overall estimate	$t_1 = -0.91, p = 0.53$	$N_{articles} = 2$ $N_{species} = 2$ $N_{observations} = 2$	$I^2_{total} = 57.4$ $I^2_{article} = 28.7$ $I^2_{obs} = 29$
Abundance association Locomotion: Non-volant Zr					
overall	0.14 $\pm$ [-0.29, 0.56]	Overall estimate	$t_{12} = 0.7, p = 0.49$	$N_{articles} = 13$ $N_{species} = 12$ $N_{observations} = 21$	$I^2_{total} = 83.4$ $I^2_{article} = 58$ $I^2_{obs} = 25$
Abundance association Foraging habitat: ground Zr					
overall	-0.88 $\pm$ [-2.76, 1.01]	Overall estimate	$t_2 = -2, p = 0.18$	$N_{articles} = 3$ $N_{species} = 3$ $N_{observations} = 3$	$I^2_{total} = 51$ $I^2_{article} = 25.5$ $I^2_{obs} = 25$
Abundance association Foraging habitat: other Zr					
overall	0.18 $\pm$ [-0.21, 0.58]	Overall estimate	$t_{12} = 1, p = 0.34$	$N_{articles} = 13$ $N_{species} = 11$ $N_{observations} = 20$	$I^2_{total} = 81.1$ $I^2_{article} = 57.9$ $I^2_{obs} = 23$
Abundance association Ground nesting: ground Zr					

Table S10. Model summaries for habitat specific groupings, including foraging habitat (ground versus other), nesting habitat (ground versus other), locomotion (volant versus non-volant) and landform (continent vs island). Each habitat factor level was run independently in a subgroup intercept-only model. Model estimates $\pm$ 95% CIs and test statistics are reported along with sample sizes and residual unexplained heterogeneity (I^2), decomposed by hierarchical model levels (article and observation). The effect size used is given in the heading of each model (SMD=standardized mean difference or Hedges' g; Zr=correlation coefficient; lnOR=log odds ratio).

Term	Estimate	Moderator	Test statistics	Sample size	Residual heterogeneity
overall	-0.32 $\pm$ [-2.59, 1.96]	Overall estimate	$t_2 = -0.6, p = 0.61$	$N_{articles} = 3$ $N_{species} = 3$ $N_{observations} = 4$	$I^2_{total} = 74.3$ $I^2_{article} = 68.4$ $I^2_{obs} = 6$
Abundance association Ground nesting: other Zr					
overall	0.09 $\pm$ [-0.35, 0.53]	Overall estimate	$t_{12} = 0.46, p = 0.66$	$N_{articles} = 13$ $N_{species} = 11$ $N_{observations} = 19$	$I^2_{total} = 84.7$ $I^2_{article} = 63.6$ $I^2_{obs} = 21$
Abundance association Landform: mainland Zr					
overall	0.3 $\pm$ [-0.17, 0.77]	Overall estimate	$t_{10} = 1.43, p = 0.18$	$N_{articles} = 11$ $N_{species} = 10$ $N_{observations} = 16$	$I^2_{total} = 81.5$ $I^2_{article} = 54.8$ $I^2_{obs} = 27$
Abundance association Landform: island Zr					
overall	-0.45 $\pm$ [-1.28, 0.37]	Overall estimate	$t_3 = -1.75, p = 0.18$	$N_{articles} = 4$ $N_{species} = 4$ $N_{observations} = 7$	$I^2_{total} = 66$ $I^2_{article} = 31.2$ $I^2_{obs} = 35$
Abundance spatial association Locomotion: Non-volant Zr					
overall	0.1 $\pm$ [-1.25, 1.45]	Overall estimate	$t_3 = 0.24, p = 0.83$	$N_{articles} = 4$ $N_{species} = 2$ $N_{observations} = 4$	$I^2_{total} = 67.3$ $I^2_{article} = 33.6$ $I^2_{obs} = 34$
Abundance spatial association Foraging habitat: other Zr					
overall	0.1 $\pm$ [-1.25, 1.45]	Overall estimate	$t_3 = 0.24, p = 0.83$	$N_{articles} = 4$ $N_{species} = 2$ $N_{observations} = 4$	$I^2_{total} = 67.3$ $I^2_{article} = 33.6$ $I^2_{obs} = 34$
Abundance spatial association Ground nesting: other Zr					
overall	0.1 $\pm$ [-1.25, 1.45]	Overall estimate	$t_3 = 0.24, p = 0.83$	$N_{articles} = 4$ $N_{species} = 2$ $N_{observations} = 4$	$I^2_{total} = 67.3$ $I^2_{article} = 33.6$ $I^2_{obs} = 34$
Abundance spatial association Landform: mainland Zr					

Table S10. Model summaries for habitat specific groupings, including foraging habitat (ground versus other), nesting habitat (ground versus other), locomotion (volant versus non-volant) and landform (continent vs island). Each habitat factor level was run independently in a subgroup intercept-only model. Model estimates $\pm$ 95% CIs and test statistics are reported along with sample sizes and residual unexplained heterogeneity (I^2), decomposed by hierarchical model levels (article and observation). The effect size used is given in the heading of each model (SMD=standardized mean difference or Hedges' g; Zr=correlation coefficient; lnOR=log odds ratio).

Term	Estimate	Moderator	Test statistics	Sample size	Residual heterogeneity
overall	0.31 $\pm$ [-2.23, 2.84]	Overall estimate	$t_2 = 0.52, p = 0.65$	$N_{articles} = 3$ $N_{species} = 1$ $N_{observations} = 3$	$I^2_{total} = 70.9$ $I^2_{article} = 35.4$ $I^2_{obs} = 35$
Abundance temporal association Locomotion: volant Zr					
overall	-0.57 $\pm$ [-8.52, 7.38]	Overall estimate	$t_1 = -0.91, p = 0.53$	$N_{articles} = 2$ $N_{species} = 2$ $N_{observations} = 2$	$I^2_{total} = 57.4$ $I^2_{article} = 28.7$ $I^2_{obs} = 29$
Abundance temporal association Locomotion: Non-volant Zr					
overall	0.15 $\pm$ [-0.38, 0.67]	Overall estimate	$t_8 = 0.65, p = 0.53$	$N_{articles} = 9$ $N_{species} = 11$ $N_{observations} = 17$	$I^2_{total} = 86$ $I^2_{article} = 60.7$ $I^2_{obs} = 25$
Abundance temporal association Foraging habitat: ground Zr					
overall	-0.88 $\pm$ [-2.76, 1.01]	Overall estimate	$t_2 = -2, p = 0.18$	$N_{articles} = 3$ $N_{species} = 3$ $N_{observations} = 3$	$I^2_{total} = 51$ $I^2_{article} = 25.5$ $I^2_{obs} = 25$
Abundance temporal association Foraging habitat: other Zr					
overall	0.21 $\pm$ [-0.27, 0.69]	Overall estimate	$t_8 = 1.01, p = 0.34$	$N_{articles} = 9$ $N_{species} = 10$ $N_{observations} = 16$	$I^2_{total} = 83.5$ $I^2_{article} = 59.8$ $I^2_{obs} = 24$
Abundance temporal association Ground nesting: ground Zr					
overall	-0.32 $\pm$ [-2.59, 1.96]	Overall estimate	$t_2 = -0.6, p = 0.61$	$N_{articles} = 3$ $N_{species} = 3$ $N_{observations} = 4$	$I^2_{total} = 74.3$ $I^2_{article} = 68.4$ $I^2_{obs} = 6$
Abundance temporal association Ground nesting: other Zr					
overall	0.09 $\pm$ [-0.48, 0.65]	Overall estimate	$t_8 = 0.36, p = 0.73$	$N_{articles} = 9$ $N_{species} = 10$ $N_{observations} = 15$	$I^2_{total} = 87.7$ $I^2_{article} = 67.1$ $I^2_{obs} = 21$
Abundance temporal association Landform: mainland Zr					
overall	0.3 $\pm$ [-0.25, 0.86]	Overall estimate	$t_7 = 1.29, p = 0.24$	$N_{articles} = 8$ $N_{species} = 9$	$I^2_{total} = 83.7$ $I^2_{article} = 56.6$

Table S10. Model summaries for habitat specific groupings, including foraging habitat (ground versus other), nesting habitat (ground versus other), locomotion (volant versus non-volant) and landform (continent vs island). Each habitat factor level was run independently in a subgroup intercept-only model. Model estimates $\pm$ 95% CIs and test statistics are reported along with sample sizes and residual unexplained heterogeneity (I^2), decomposed by hierarchical model levels (article and observation). The effect size used is given in the heading of each model (SMD=standardized mean difference or Hedges' g; Zr=correlation coefficient; lnOR=log odds ratio).

Term	Estimate	Moderator	Test statistics	Sample size	Residual heterogeneity
				$N_{\text{observations}} = 13$	$I^2_{\text{obs}} = 27$
Abundance temporal association Landform: island Zr					
overall	-0.58 $\pm$ [-2.22, 1.05]	Overall estimate	$t_2 = -1.54, p = 0.26$	$N_{\text{articles}} = 3$ $N_{\text{species}} = 4$ $N_{\text{observations}} = 6$	$I^2_{\text{total}} = 77.5$ $I^2_{\text{article}} = 52.3$ $I^2_{\text{obs}} = 25$
Reproduction with/without cats Locomotion: volant lnOR					
overall	-0.39 $\pm$ [-3.52, 2.75]	Overall estimate	$t_2 = -0.53, p = 0.65$	$N_{\text{articles}} = 3$ $N_{\text{species}} = 3$ $N_{\text{observations}} = 5$	$I^2_{\text{total}} = 65.5$ $I^2_{\text{article}} = 28.2$ $I^2_{\text{obs}} = 37$
Reproduction with/without cats Foraging habitat: ground lnOR					
overall	-0.39 $\pm$ [-3.52, 2.75]	Overall estimate	$t_2 = -0.53, p = 0.65$	$N_{\text{articles}} = 3$ $N_{\text{species}} = 3$ $N_{\text{observations}} = 5$	$I^2_{\text{total}} = 65.5$ $I^2_{\text{article}} = 28.2$ $I^2_{\text{obs}} = 37$
Reproduction with/without cats Ground nesting: other lnOR					
overall	-0.39 $\pm$ [-3.52, 2.75]	Overall estimate	$t_2 = -0.53, p = 0.65$	$N_{\text{articles}} = 3$ $N_{\text{species}} = 3$ $N_{\text{observations}} = 5$	$I^2_{\text{total}} = 65.5$ $I^2_{\text{article}} = 28.2$ $I^2_{\text{obs}} = 37$
Reproduction with/without cats Landform: island lnOR					
overall	-0.39 $\pm$ [-3.52, 2.75]	Overall estimate	$t_2 = -0.53, p = 0.65$	$N_{\text{articles}} = 3$ $N_{\text{species}} = 3$ $N_{\text{observations}} = 5$	$I^2_{\text{total}} = 65.5$ $I^2_{\text{article}} = 28.2$ $I^2_{\text{obs}} = 37$
Reproduction inside/outside enclosure Locomotion: volant lnOR					
overall	-1.65 $\pm$ [-21.58, 18.28]	Overall estimate	$t_1 = -1.05, p = 0.48$	$N_{\text{articles}} = 1$ $N_{\text{species}} = 1$ $N_{\text{observations}} = 2$	$I^2_{\text{total}} = 70.2$ $I^2_{\text{article}} = NA$ $I^2_{\text{obs}} = NA$
Reproduction inside/outside enclosure Foraging habitat: ground lnOR					
overall	-1.65 $\pm$ [-21.58, 18.28]	Overall estimate	$t_1 = -1.05, p = 0.48$	$N_{\text{articles}} = 1$ $N_{\text{species}} = 1$ $N_{\text{observations}} = 2$	$I^2_{\text{total}} = 70.2$ $I^2_{\text{article}} = NA$ $I^2_{\text{obs}} = NA$

Table S10. Model summaries for habitat specific groupings, including foraging habitat (ground versus other), nesting habitat (ground versus other), locomotion (volant versus non-volant) and landform (continent vs island). Each habitat factor level was run independently in a subgroup intercept-only model. Model estimates $\pm$ 95% CIs and test statistics are reported along with sample sizes and residual unexplained heterogeneity (I^2), decomposed by hierarchical model levels (article and observation). The effect size used is given in the heading of each model (SMD=standardized mean difference or Hedges' g; Zr=correlation coefficient; lnOR=log odds ratio).

Term	Estimate	Moderator	Test statistics	Sample size	Residual heterogeneity
Reproduction inside/outside enclosure Ground nesting: other lnOR					
overall	-1.65 $\pm$ [-21.58, 18.28]	Overall estimate	$t_1 = -1.05, p = 0.48$	$N_{articles} = 1$ $N_{species} = 1$ $N_{observations} = 2$	$I^2_{total} = 70.2$ $I^2_{article} = NA$ $I^2_{obs} = NA$
Reproduction inside/outside enclosure Landform: island lnOR					
overall	-1.65 $\pm$ [-21.58, 18.28]	Overall estimate	$t_1 = -1.05, p = 0.48$	$N_{articles} = 1$ $N_{species} = 1$ $N_{observations} = 2$	$I^2_{total} = 70.2$ $I^2_{article} = NA$ $I^2_{obs} = NA$
Reproduction before-after eradication Locomotion: volant SMD					
overall	0.14 $\pm$ [-5.7, 5.99]	Overall estimate	$t_1 = 0.31, p = 0.81$	$N_{articles} = 2$ $N_{species} = 3$ $N_{observations} = 3$	$I^2_{total} = 54.4$ $I^2_{article} = 54.4$ $I^2_{obs} = 0$
Reproduction before-after eradication Foraging habitat: ground SMD					
overall	0.14 $\pm$ [-5.7, 5.99]	Overall estimate	$t_1 = 0.31, p = 0.81$	$N_{articles} = 2$ $N_{species} = 3$ $N_{observations} = 3$	$I^2_{total} = 54.4$ $I^2_{article} = 54.4$ $I^2_{obs} = 0$
Reproduction before-after eradication Ground nesting: other SMD					
overall	0.14 $\pm$ [-5.7, 5.99]	Overall estimate	$t_1 = 0.31, p = 0.81$	$N_{articles} = 2$ $N_{species} = 3$ $N_{observations} = 3$	$I^2_{total} = 54.4$ $I^2_{article} = 54.4$ $I^2_{obs} = 0$
Reproduction before-after eradication Landform: island SMD					
overall	0.14 $\pm$ [-5.7, 5.99]	Overall estimate	$t_1 = 0.31, p = 0.81$	$N_{articles} = 2$ $N_{species} = 3$ $N_{observations} = 3$	$I^2_{total} = 54.4$ $I^2_{article} = 54.4$ $I^2_{obs} = 0$
Reproduction association Locomotion: volant Zr					
overall	-0.51 $\pm$ [-11.08, 10.06]	Overall estimate	$t_1 = -0.62, p = 0.65$	$N_{articles} = 2$ $N_{species} = 2$ $N_{observations} = 5$	$I^2_{total} = 80.3$ $I^2_{article} = 80.3$ $I^2_{obs} = 0$
Reproduction association Foraging habitat: ground Zr					

Table S10. Model summaries for habitat specific groupings, including foraging habitat (ground versus other), nesting habitat (ground versus other), locomotion (volant versus non-volant) and landform (continent vs island). Each habitat factor level was run independently in a subgroup intercept-only model. Model estimates $\pm$ 95% CIs and test statistics are reported along with sample sizes and residual unexplained heterogeneity (I^2), decomposed by hierarchical model levels (article and observation). The effect size used is given in the heading of each model (SMD=standardized mean difference or Hedges' g; Zr=correlation coefficient; lnOR=log odds ratio).

Term	Estimate	Moderator	Test statistics	Sample size	Residual heterogeneity
overall	0.09 $\pm$ [-0.62, 0.8]	Overall estimate	$t_3 = 0.41, p = 0.71$	$N_{articles} = 1$ $N_{species} = 1$ $N_{observations} = 4$	$I^2_{total} = 0$ $I^2_{article} = NA$ $I^2_{obs} = NA$
Reproduction association Ground nesting: other Zr					
overall	-0.51 $\pm$ [-11.08, 10.06]	Overall estimate	$t_1 = -0.62, p = 0.65$	$N_{articles} = 2$ $N_{species} = 2$ $N_{observations} = 5$	$I^2_{total} = 80.3$ $I^2_{article} = 80.3$ $I^2_{obs} = 0$
Reproduction association Landform: island Zr					
overall	-0.51 $\pm$ [-11.08, 10.06]	Overall estimate	$t_1 = -0.62, p = 0.65$	$N_{articles} = 2$ $N_{species} = 2$ $N_{observations} = 5$	$I^2_{total} = 80.3$ $I^2_{article} = 80.3$ $I^2_{obs} = 0$
Reproduction temporal association Locomotion: volant Zr					
overall	0.09 $\pm$ [-0.62, 0.8]	Overall estimate	$t_3 = 0.41, p = 0.71$	$N_{articles} = 1$ $N_{species} = 1$ $N_{observations} = 4$	$I^2_{total} = 0$ $I^2_{article} = NA$ $I^2_{obs} = NA$
Reproduction temporal association Foraging habitat: ground Zr					
overall	0.09 $\pm$ [-0.62, 0.8]	Overall estimate	$t_3 = 0.41, p = 0.71$	$N_{articles} = 1$ $N_{species} = 1$ $N_{observations} = 4$	$I^2_{total} = 0$ $I^2_{article} = NA$ $I^2_{obs} = NA$
Reproduction temporal association Ground nesting: other Zr					
overall	0.09 $\pm$ [-0.62, 0.8]	Overall estimate	$t_3 = 0.41, p = 0.71$	$N_{articles} = 1$ $N_{species} = 1$ $N_{observations} = 4$	$I^2_{total} = 0$ $I^2_{article} = NA$ $I^2_{obs} = NA$
Reproduction temporal association Landform: island Zr					
overall	0.09 $\pm$ [-0.62, 0.8]	Overall estimate	$t_3 = 0.41, p = 0.71$	$N_{articles} = 1$ $N_{species} = 1$ $N_{observations} = 4$	$I^2_{total} = 0$ $I^2_{article} = NA$ $I^2_{obs} = NA$