

Table S10. Model summaries for habitat specific groupings, including foraging habitat (ground versus other), nesting habitat (ground versus other), locomotion (volant versus non-volant) and landform (continent vs island). Each habitat factor level was run independently in a subgroup intercept-only model. Model estimates $\pm$ 95% CIs and test statistics are reported along with sample sizes and residual unexplained heterogeneity (I^2), decomposed by hierarchical model levels (article and observation). The effect size used is given in the heading of each model (SMD=standardized mean difference or Hedges' g; Zr=correlation coefficient; lnOR=log odds ratio).

Term	Estimate	Moderator	Test statistics	Sample size	Residual heterogeneity
Abundance with/without cats Foraging habitat: ground SMD					
overall	0.11 $\pm$ [-1.89, 2.11]	Overall estimate	$t_3 = 0.18, p = 0.87$	$N_{articles} = 4$ $N_{species} = 4$ $N_{observations} = 4$	$I^2_{total} = 59.5$ $I^2_{article} = 29.7$ $I^2_{obs} = 30$
Abundance with/without cats Foraging habitat: other SMD					
overall	-0.58 $\pm$ [-1.33, 0.17]	Overall estimate	$t_6 = -1.9, p = 0.11$	$N_{articles} = 7$ $N_{species} = 9$ $N_{observations} = 11$	$I^2_{total} = 44.2$ $I^2_{article} = 44.2$ $I^2_{obs} = 0$
Abundance with/without cats Ground nesting: ground SMD					
overall	0.39 $\pm$ [-2.17, 2.96]	Overall estimate	$t_2 = 0.66, p = 0.58$	$N_{articles} = 3$ $N_{species} = 6$ $N_{observations} = 6$	$I^2_{total} = 42.1$ $I^2_{article} = 42.1$ $I^2_{obs} = 0$
Abundance with/without cats Ground nesting: other SMD					
overall	-0.77 $\pm$ [-1.39, -0.15]	Overall estimate	$t_7 = -2.96, p = 0.02$	$N_{articles} = 8$ $N_{species} = 7$ $N_{observations} = 9$	$I^2_{total} = 34.7$ $I^2_{article} = 34.7$ $I^2_{obs} = 0$
Abundance with/without cats Locomotion: Non-volant SMD					
overall	-0.33 $\pm$ [-1.28, 0.63]	Overall estimate	$t_5 = -0.88, p = 0.42$	$N_{articles} = 6$ $N_{species} = 8$ $N_{observations} = 8$	$I^2_{total} = 55.8$ $I^2_{article} = 55.8$ $I^2_{obs} = 0$
Abundance with/without cats Locomotion: volant SMD					
overall	-0.55 $\pm$ [-2.34, 1.24]	Overall estimate	$t_3 = -0.98, p = 0.4$	$N_{articles} = 4$ $N_{species} = 5$ $N_{observations} = 7$	$I^2_{total} = 56$ $I^2_{article} = 56$ $I^2_{obs} = 0$
Abundance with/without cats Landform: island SMD					
overall	-0.28 $\pm$ [-0.9, 0.35]	Overall estimate	$t_8 = -1.02, p = 0.34$	$N_{articles} = 9$ $N_{species} = 12$ $N_{observations} = 14$	$I^2_{total} = 31.1$ $I^2_{article} = 31.1$ $I^2_{obs} = 0$

Table S10. Model summaries for habitat specific groupings, including foraging habitat (ground versus other), nesting habitat (ground versus other), locomotion (volant versus non-volant) and landform (continent vs island). Each habitat factor level was run independently in a subgroup intercept-only model. Model estimates $\pm$ 95% CIs and test statistics are reported along with sample sizes and residual unexplained heterogeneity (I^2), decomposed by hierarchical model levels (article and observation). The effect size used is given in the heading of each model (SMD=standardized mean difference or Hedges' g; Zr=correlation coefficient; lnOR=log odds ratio).

Term	Estimate	Moderator	Test statistics	Sample size	Residual heterogeneity
Abundance before-after eradication Foraging habitat: other SMD					
overall	-0.91 $\pm$ [-8.73, 6.91]	Overall estimate	$t_1 = -1.48, p = 0.38$	$N_{articles} = 2$ $N_{species} = 2$ $N_{observations} = 3$	$I^2_{total} = 38$ $I^2_{article} = 38$ $I^2_{obs} = 0$
Abundance before-after eradication Ground nesting: other SMD					
overall	-0.91 $\pm$ [-8.73, 6.91]	Overall estimate	$t_1 = -1.48, p = 0.38$	$N_{articles} = 2$ $N_{species} = 2$ $N_{observations} = 3$	$I^2_{total} = 38$ $I^2_{article} = 38$ $I^2_{obs} = 0$
Abundance before-after eradication Landform: island SMD					
overall	-0.91 $\pm$ [-8.73, 6.91]	Overall estimate	$t_1 = -1.48, p = 0.38$	$N_{articles} = 2$ $N_{species} = 2$ $N_{observations} = 3$	$I^2_{total} = 38$ $I^2_{article} = 38$ $I^2_{obs} = 0$
Abundance on islands with/without cats Foraging habitat: ground lnOR					
overall	0.27 $\pm$ [-3.57, 4.1]	Overall estimate	$t_3 = 0.22, p = 0.84$	$N_{articles} = 4$ $N_{species} = 4$ $N_{observations} = 4$	$I^2_{total} = 63.7$ $I^2_{article} = 31.9$ $I^2_{obs} = 32$
Abundance on islands with/without cats Ground nesting: ground lnOR					
overall	0.77 $\pm$ [-4.36, 5.9]	Overall estimate	$t_2 = 0.65, p = 0.58$	$N_{articles} = 3$ $N_{species} = 6$ $N_{observations} = 6$	$I^2_{total} = 49.7$ $I^2_{article} = 49.7$ $I^2_{obs} = 0$
Abundance on islands with/without cats Ground nesting: other lnOR					
overall	-0.76 $\pm$ [-2.15, 0.63]	Overall estimate	$t_4 = -1.52, p = 0.2$	$N_{articles} = 5$ $N_{species} = 5$ $N_{observations} = 5$	$I^2_{total} = 0$ $I^2_{article} = 0$ $I^2_{obs} = 0$
Abundance on islands with/without cats Locomotion: Non-volant lnOR					
overall	0.08 $\pm$ [-1.82, 1.99]	Overall estimate	$t_3 = 0.14, p = 0.9$	$N_{articles} = 4$ $N_{species} = 6$ $N_{observations} = 6$	$I^2_{total} = 13.3$ $I^2_{article} = 13.3$ $I^2_{obs} = 0$
Abundance on islands with/without cats Locomotion: volant lnOR					

Table S10. Model summaries for habitat specific groupings, including foraging habitat (ground versus other), nesting habitat (ground versus other), locomotion (volant versus non-volant) and landform (continent vs island). Each habitat factor level was run independently in a subgroup intercept-only model. Model estimates $\pm$ 95% CIs and test statistics are reported along with sample sizes and residual unexplained heterogeneity (I^2), decomposed by hierarchical model levels (article and observation). The effect size used is given in the heading of each model (SMD=standardized mean difference or Hedges' g; Zr=correlation coefficient; lnOR=log odds ratio).

Term	Estimate	Moderator	Test statistics	Sample size	Residual heterogeneity
overall	-0.35 $\pm$ [-6.25, 5.55]	Overall estimate	$t_2 = -0.25, p = 0.82$	$N_{articles} = 3$ $N_{species} = 5$ $N_{observations} = 5$	$I^2_{total} = 65.8$ $I^2_{article} = 65.8$ $I^2_{obs} = 0$
Abundance on islands with/without cats Landform: island lnOR					
overall	-0.16 $\pm$ [-1.56, 1.24]	Overall estimate	$t_6 = -0.29, p = 0.78$	$N_{articles} = 7$ $N_{species} = 11$ $N_{observations} = 11$	$I^2_{total} = 34$ $I^2_{article} = 34$ $I^2_{obs} = 0$
Abundance spatial association Foraging habitat: other Zr					
overall	0.1 $\pm$ [-1.25, 1.45]	Overall estimate	$t_3 = 0.24, p = 0.83$	$N_{articles} = 4$ $N_{species} = 2$ $N_{observations} = 4$	$I^2_{total} = 67.3$ $I^2_{article} = 33.6$ $I^2_{obs} = 34$
Abundance spatial association Ground nesting: other Zr					
overall	0.1 $\pm$ [-1.25, 1.45]	Overall estimate	$t_3 = 0.24, p = 0.83$	$N_{articles} = 4$ $N_{species} = 2$ $N_{observations} = 4$	$I^2_{total} = 67.3$ $I^2_{article} = 33.6$ $I^2_{obs} = 34$
Abundance spatial association Locomotion: Non-volant Zr					
overall	0.1 $\pm$ [-1.25, 1.45]	Overall estimate	$t_3 = 0.24, p = 0.83$	$N_{articles} = 4$ $N_{species} = 2$ $N_{observations} = 4$	$I^2_{total} = 67.3$ $I^2_{article} = 33.6$ $I^2_{obs} = 34$
Abundance spatial association Landform: mainland Zr					
overall	0.31 $\pm$ [-2.23, 2.84]	Overall estimate	$t_2 = 0.52, p = 0.65$	$N_{articles} = 3$ $N_{species} = 1$ $N_{observations} = 3$	$I^2_{total} = 70.9$ $I^2_{article} = 35.4$ $I^2_{obs} = 35$
Abundance temporal association Foraging habitat: ground Zr					
overall	-0.88 $\pm$ [-2.76, 1.01]	Overall estimate	$t_2 = -2, p = 0.18$	$N_{articles} = 3$ $N_{species} = 3$ $N_{observations} = 3$	$I^2_{total} = 51$ $I^2_{article} = 25.5$ $I^2_{obs} = 25$
Abundance temporal association Foraging habitat: other Zr					

Table S10. Model summaries for habitat specific groupings, including foraging habitat (ground versus other), nesting habitat (ground versus other), locomotion (volant versus non-volant) and landform (continent vs island). Each habitat factor level was run independently in a subgroup intercept-only model. Model estimates $\pm$ 95% CIs and test statistics are reported along with sample sizes and residual unexplained heterogeneity (I^2), decomposed by hierarchical model levels (article and observation). The effect size used is given in the heading of each model (SMD=standardized mean difference or Hedges' g; Zr=correlation coefficient; lnOR=log odds ratio).

Term	Estimate	Moderator	Test statistics	Sample size	Residual heterogeneity
overall	0.21 $\pm$ [-0.27, 0.69]	Overall estimate	$t_8 = 1.01, p = 0.34$	$N_{articles} = 9$ $N_{species} = 10$ $N_{observations} = 16$	$I^2_{total} = 83.5$ $I^2_{article} = 59.8$ $I^2_{obs} = 24$
Abundance temporal association Ground nesting: ground Zr					
overall	-0.32 $\pm$ [-2.59, 1.96]	Overall estimate	$t_2 = -0.6, p = 0.61$	$N_{articles} = 3$ $N_{species} = 3$ $N_{observations} = 4$	$I^2_{total} = 74.3$ $I^2_{article} = 68.4$ $I^2_{obs} = 6$
Abundance temporal association Ground nesting: other Zr					
overall	0.09 $\pm$ [-0.48, 0.65]	Overall estimate	$t_8 = 0.36, p = 0.73$	$N_{articles} = 9$ $N_{species} = 10$ $N_{observations} = 15$	$I^2_{total} = 87.7$ $I^2_{article} = 67.1$ $I^2_{obs} = 21$
Abundance temporal association Locomotion: Non-volant Zr					
overall	0.15 $\pm$ [-0.38, 0.67]	Overall estimate	$t_8 = 0.65, p = 0.53$	$N_{articles} = 9$ $N_{species} = 11$ $N_{observations} = 17$	$I^2_{total} = 86$ $I^2_{article} = 60.7$ $I^2_{obs} = 25$
Abundance temporal association Landform: island Zr					
overall	-0.58 $\pm$ [-2.22, 1.05]	Overall estimate	$t_2 = -1.54, p = 0.26$	$N_{articles} = 3$ $N_{species} = 4$ $N_{observations} = 6$	$I^2_{total} = 77.5$ $I^2_{article} = 52.3$ $I^2_{obs} = 25$
Abundance temporal association Landform: mainland Zr					
overall	0.3 $\pm$ [-0.25, 0.86]	Overall estimate	$t_7 = 1.29, p = 0.24$	$N_{articles} = 8$ $N_{species} = 9$ $N_{observations} = 13$	$I^2_{total} = 83.7$ $I^2_{article} = 56.6$ $I^2_{obs} = 27$
Reproduction with/without cats Foraging habitat: ground lnOR					
overall	-0.39 $\pm$ [-3.52, 2.75]	Overall estimate	$t_2 = -0.53, p = 0.65$	$N_{articles} = 3$ $N_{species} = 3$ $N_{observations} = 5$	$I^2_{total} = 65.5$ $I^2_{article} = 28.2$ $I^2_{obs} = 37$
Reproduction with/without cats Ground nesting: other lnOR					

Table S10. Model summaries for habitat specific groupings, including foraging habitat (ground versus other), nesting habitat (ground versus other), locomotion (volant versus non-volant) and landform (continent vs island). Each habitat factor level was run independently in a subgroup intercept-only model. Model estimates $\pm$ 95% CIs and test statistics are reported along with sample sizes and residual unexplained heterogeneity (I^2), decomposed by hierarchical model levels (article and observation). The effect size used is given in the heading of each model (SMD=standardized mean difference or Hedges' g; Zr=correlation coefficient; lnOR=log odds ratio).

Term	Estimate	Moderator	Test statistics	Sample size	Residual heterogeneity
overall	-0.39 $\pm$ [-3.52, 2.75]	Overall estimate	$t_2 = -0.53, p = 0.65$	$N_{articles} = 3$ $N_{species} = 3$ $N_{observations} = 5$	$I^2_{total} = 65.5$ $I^2_{article} = 28.2$ $I^2_{obs} = 37$
Reproduction with/without cats Locomotion: volant lnOR					
overall	-0.39 $\pm$ [-3.52, 2.75]	Overall estimate	$t_2 = -0.53, p = 0.65$	$N_{articles} = 3$ $N_{species} = 3$ $N_{observations} = 5$	$I^2_{total} = 65.5$ $I^2_{article} = 28.2$ $I^2_{obs} = 37$
Reproduction with/without cats Landform: island lnOR					
overall	-0.39 $\pm$ [-3.52, 2.75]	Overall estimate	$t_2 = -0.53, p = 0.65$	$N_{articles} = 3$ $N_{species} = 3$ $N_{observations} = 5$	$I^2_{total} = 65.5$ $I^2_{article} = 28.2$ $I^2_{obs} = 37$
Reproduction before-after eradication Foraging habitat: ground SMD					
overall	0.14 $\pm$ [-5.7, 5.99]	Overall estimate	$t_1 = 0.31, p = 0.81$	$N_{articles} = 2$ $N_{species} = 3$ $N_{observations} = 3$	$I^2_{total} = 54.4$ $I^2_{article} = 54.4$ $I^2_{obs} = 0$
Reproduction before-after eradication Ground nesting: other SMD					
overall	0.14 $\pm$ [-5.7, 5.99]	Overall estimate	$t_1 = 0.31, p = 0.81$	$N_{articles} = 2$ $N_{species} = 3$ $N_{observations} = 3$	$I^2_{total} = 54.4$ $I^2_{article} = 54.4$ $I^2_{obs} = 0$
Reproduction before-after eradication Locomotion: volant SMD					
overall	0.14 $\pm$ [-5.7, 5.99]	Overall estimate	$t_1 = 0.31, p = 0.81$	$N_{articles} = 2$ $N_{species} = 3$ $N_{observations} = 3$	$I^2_{total} = 54.4$ $I^2_{article} = 54.4$ $I^2_{obs} = 0$
Reproduction before-after eradication Landform: island SMD					
overall	0.14 $\pm$ [-5.7, 5.99]	Overall estimate	$t_1 = 0.31, p = 0.81$	$N_{articles} = 2$ $N_{species} = 3$ $N_{observations} = 3$	$I^2_{total} = 54.4$ $I^2_{article} = 54.4$ $I^2_{obs} = 0$
Reproduction temporal association Foraging habitat: ground Zr					

Table S10. Model summaries for habitat specific groupings, including foraging habitat (ground versus other), nesting habitat (ground versus other), locomotion (volant versus non-volant) and landform (continent vs island). Each habitat factor level was run independently in a subgroup intercept-only model. Model estimates $\pm$ 95% CIs and test statistics are reported along with sample sizes and residual unexplained heterogeneity (I^2), decomposed by hierarchical model levels (article and observation). The effect size used is given in the heading of each model (SMD=standardized mean difference or Hedges' g; Zr=correlation coefficient; lnOR=log odds ratio).

Term	Estimate	Moderator	Test statistics	Sample size	Residual heterogeneity
overall	0.09 $\pm$ [-0.62, 0.8]	Overall estimate	$t_3 = 0.41, p = 0.71$	$N_{articles} = 1$ $N_{species} = 1$ $N_{observations} = 4$	$I^2_{total} = 0$ $I^2_{article} = NA$ $I^2_{obs} = NA$
Reproduction temporal association Ground nesting: other Zr					
overall	0.09 $\pm$ [-0.62, 0.8]	Overall estimate	$t_3 = 0.41, p = 0.71$	$N_{articles} = 1$ $N_{species} = 1$ $N_{observations} = 4$	$I^2_{total} = 0$ $I^2_{article} = NA$ $I^2_{obs} = NA$
Reproduction temporal association Locomotion: volant Zr					
overall	0.09 $\pm$ [-0.62, 0.8]	Overall estimate	$t_3 = 0.41, p = 0.71$	$N_{articles} = 1$ $N_{species} = 1$ $N_{observations} = 4$	$I^2_{total} = 0$ $I^2_{article} = NA$ $I^2_{obs} = NA$
Reproduction temporal association Landform: island Zr					
overall	0.09 $\pm$ [-0.62, 0.8]	Overall estimate	$t_3 = 0.41, p = 0.71$	$N_{articles} = 1$ $N_{species} = 1$ $N_{observations} = 4$	$I^2_{total} = 0$ $I^2_{article} = NA$ $I^2_{obs} = NA$