

Table S6. Model summaries for intercept-only (main text) models as well as models evaluating influence of habitat, body mass, and continents versus islands. Model estimates $\pm$ 95% CIs and test statistics are reported along with sample sizes and residual unexplained heterogeneity (I^2), decomposed by hierarchical model levels (article and observation). The effect size used is given in the heading of each model (SMD=standardized mean difference or Hedges' g; Zr=correlation coefficient; lnOR=log odds ratio).

Term	Estimate	Moderator	Test statistics	Sample size	Residual heterogeneity
Abundance with/without cats All species Overall estimate SMD					
overall	-0.44 $\pm$ [-1.07, 0.18]	Overall estimate	$t_9 = -1.6, p = 0.14$	$N_{articles} = 10$ $N_{species} = 13$ $N_{observations} = 15$	$I^2_{total} = 54$ $I^2_{article} = 54$ $I^2_{obs} = 0$
Abundance before-after eradication All species Overall estimate SMD					
overall	-0.91 $\pm$ [-6.27, 4.45]	Overall estimate	$t_1 = -2.16, p = 0.28$	$N_{articles} = 2$ $N_{species} = 2$ $N_{observations} = 3$	$I^2_{total} = 26.5$ $I^2_{article} = 26.5$ $I^2_{obs} = 0$
Abundance on islands with/without cats All species Overall estimate lnOR					
overall	-0.16 $\pm$ [-1.56, 1.24]	Overall estimate	$t_6 = -0.29, p = 0.78$	$N_{articles} = 7$ $N_{species} = 11$ $N_{observations} = 11$	$I^2_{total} = 34$ $I^2_{article} = 34$ $I^2_{obs} = 0$
Abundance spatial association All species Overall estimate Zr					
overall	0.1 $\pm$ [-1.25, 1.45]	Overall estimate	$t_3 = 0.24, p = 0.83$	$N_{articles} = 4$ $N_{species} = 2$ $N_{observations} = 4$	$I^2_{total} = 67.3$ $I^2_{article} = 33.6$ $I^2_{obs} = 34$
Abundance temporal association All species Overall estimate Zr					
overall	0.06 $\pm$ [-0.42, 0.54]	Overall estimate	$t_{10} = 0.28, p = 0.78$	$N_{articles} = 11$ $N_{species} = 13$ $N_{observations} = 19$	$I^2_{total} = 85.3$ $I^2_{article} = 60.5$ $I^2_{obs} = 25$
Reproduction with/without cats All species Overall estimate lnOR					
overall	-0.08 $\pm$ [-2.62, 2.47]	Overall estimate	$t_2 = -0.13, p = 0.91$	$N_{articles} = 3$ $N_{species} = 3$ $N_{observations} = 5$	$I^2_{total} = 68.8$ $I^2_{article} = 68.8$ $I^2_{obs} = 0$
Reproduction before-after eradication All species Overall estimate SMD					
overall	0.16 $\pm$ [-4.5, 4.82]	Overall estimate	$t_1 = 0.43, p = 0.74$	$N_{articles} = 2$ $N_{species} = 3$ $N_{observations} = 3$	$I^2_{total} = 75$ $I^2_{article} = 75$ $I^2_{obs} = 0$

Table S6. Model summaries for intercept-only (main text) models as well as models evaluating influence of habitat, body mass, and continents versus islands. Model estimates $\pm$ 95% CIs and test statistics are reported along with sample sizes and residual unexplained heterogeneity (I^2), decomposed by hierarchical model levels (article and observation). The effect size used is given in the heading of each model (SMD=standardized mean difference or Hedges' g; Zr=correlation coefficient; lnOR=log odds ratio).

Term	Estimate	Moderator	Test statistics	Sample size	Residual heterogeneity
Reproduction temporal association All species Overall estimate Zr					
overall	0.09 $\pm$ [-0.62, 0.8]	Overall estimate	$t_3 = 0.41, p = 0.71$	$N_{articles} = 1$ $N_{species} = 1$ $N_{observations} = 4$	$I^2_{total} = 0$ $I^2_{article} = NA$ $I^2_{obs} = NA$
Abundance with/without cats Mammals Overall estimate SMD					
overall	-0.59 $\pm$ [-2.19, 1]	Overall estimate	$t_3 = -1.18, p = 0.32$	$N_{articles} = 4$ $N_{species} = 6$ $N_{observations} = 6$	$I^2_{total} = 60.4$ $I^2_{article} = 60.4$ $I^2_{obs} = 0$
Abundance on islands with/without cats Mammals Overall estimate lnOR					
overall	-0.25 $\pm$ [-20.55, 20.04]	Overall estimate	$t_1 = -0.16, p = 0.9$	$N_{articles} = 2$ $N_{species} = 4$ $N_{observations} = 4$	$I^2_{total} = 60.1$ $I^2_{article} = 60.1$ $I^2_{obs} = 0$
Abundance spatial association Mammals Overall estimate Zr					
overall	0.1 $\pm$ [-1.25, 1.45]	Overall estimate	$t_3 = 0.24, p = 0.83$	$N_{articles} = 4$ $N_{species} = 2$ $N_{observations} = 4$	$I^2_{total} = 67.3$ $I^2_{article} = 33.6$ $I^2_{obs} = 34$
Abundance temporal association Mammals Overall estimate Zr					
overall	0.15 $\pm$ [-0.38, 0.67]	Overall estimate	$t_8 = 0.65, p = 0.53$	$N_{articles} = 9$ $N_{species} = 11$ $N_{observations} = 17$	$I^2_{total} = 86$ $I^2_{article} = 60.7$ $I^2_{obs} = 25$
Abundance with/without cats Birds Overall estimate SMD					
overall	-0.45 $\pm$ [-2, 1.09]	Overall estimate	$t_3 = -0.94, p = 0.42$	$N_{articles} = 4$ $N_{species} = 5$ $N_{observations} = 7$	$I^2_{total} = 54.7$ $I^2_{article} = 54.7$ $I^2_{obs} = 0$
Abundance on islands with/without cats Birds Overall estimate lnOR					
overall	-0.13 $\pm$ [-5.5, 5.24]	Overall estimate	$t_2 = -0.11, p = 0.93$	$N_{articles} = 3$ $N_{species} = 5$ $N_{observations} = 5$	$I^2_{total} = 58.5$ $I^2_{article} = 58.5$ $I^2_{obs} = 0$
Reproduction with/without cats Birds Overall estimate lnOR					

Table S6. Model summaries for intercept-only (main text) models as well as models evaluating influence of habitat, body mass, and continents versus islands. Model estimates $\pm$ 95% CIs and test statistics are reported along with sample sizes and residual unexplained heterogeneity (I^2), decomposed by hierarchical model levels (article and observation). The effect size used is given in the heading of each model (SMD=standardized mean difference or Hedges' g; Zr=correlation coefficient; lnOR=log odds ratio).

Term	Estimate	Moderator	Test statistics	Sample size	Residual heterogeneity
overall	-0.08 $\pm$ [-2.62, 2.47]	Overall estimate	$t_2 = -0.13, p = 0.91$	$N_{articles} = 3$ $N_{species} = 3$ $N_{observations} = 5$	$I^2_{total} = 68.8$ $I^2_{article} = 68.8$ $I^2_{obs} = 0$
Reproduction before-after eradication Birds Overall estimate SMD					
overall	0.16 $\pm$ [-4.5, 4.82]	Overall estimate	$t_1 = 0.43, p = 0.74$	$N_{articles} = 2$ $N_{species} = 3$ $N_{observations} = 3$	$I^2_{total} = 75$ $I^2_{article} = 75$ $I^2_{obs} = 0$
Reproduction temporal association Birds Overall estimate Zr					
overall	0.09 $\pm$ [-0.62, 0.8]	Overall estimate	$t_3 = 0.41, p = 0.71$	$N_{articles} = 1$ $N_{species} = 1$ $N_{observations} = 4$	$I^2_{total} = 0$ $I^2_{article} = NA$ $I^2_{obs} = NA$
Abundance with/without cats All species Mass SMD					
intercept	-0.2 $\pm$ [-2.37, 1.96]	Mass	$t_8 = -0.22, p = 0.83$	$N_{articles} = 10$ $N_{species} = 13$ $N_{observations} = 15$	$I^2_{total} = 57.7$ $I^2_{article} = 57.7$ $I^2_{obs} = 0$
Mass (g, log10)	-0.1 $\pm$ [-0.99, 0.78]	Mass	$t_{13} = -0.25, p = 0.81$	$N_{articles} = 10$ $N_{species} = 13$ $N_{observations} = 15$	$I^2_{total} = 57.7$ $I^2_{article} = 57.7$ $I^2_{obs} = 0$
Abundance on islands with/without cats All species Mass lnOR					
intercept	0.63 $\pm$ [-4.32, 5.58]	Mass	$t_5 = 0.33, p = 0.76$	$N_{articles} = 7$ $N_{species} = 11$ $N_{observations} = 11$	$I^2_{total} = 42.1$ $I^2_{article} = 42.1$ $I^2_{obs} = 0$
Mass (g, log10)	-0.34 $\pm$ [-2.15, 1.48]	Mass	$t_9 = -0.42, p = 0.69$	$N_{articles} = 7$ $N_{species} = 11$ $N_{observations} = 11$	$I^2_{total} = 42.1$ $I^2_{article} = 42.1$ $I^2_{obs} = 0$
Abundance temporal association All species Mass Zr					
intercept	-0.1 $\pm$ [-3.9, 3.7]	Mass	$t_9 = -0.06, p = 0.95$	$N_{articles} = 11$ $N_{species} = 13$ $N_{observations} = 19$	$I^2_{total} = 86.4$ $I^2_{article} = 62.3$ $I^2_{obs} = 24$

Table S6. Model summaries for intercept-only (main text) models as well as models evaluating influence of habitat, body mass, and continents versus islands. Model estimates $\pm$ 95% CIs and test statistics are reported along with sample sizes and residual unexplained heterogeneity (I^2), decomposed by hierarchical model levels (article and observation). The effect size used is given in the heading of each model (SMD=standardized mean difference or Hedges' g; Zr=correlation coefficient; lnOR=log odds ratio).

Term	Estimate	Moderator	Test statistics	Sample size	Residual heterogeneity
Mass (g, log10)	0.06 $\pm$ [-1.16, 1.27]	Mass	$t_{17} = 0.1, p = 0.92$	$N_{articles} = 11$ $N_{species} = 13$ $N_{observations} = 19$	$I^2_{total} = 86.4$ $I^2_{article} = 62.3$ $I^2_{obs} = 24$
Abundance temporal association All species Continent vs island Zr					
island (Intercept)	-0.59 $\pm$ [-1.48, 0.29]	Continent vs island	$t_9 = -1.51, p = 0.16$	$N_{articles} = 8$ $N_{species} = 9$ $N_{observations} = 13$	$I^2_{total} = 82.8$ $I^2_{article} = 56.7$ $I^2_{obs} = 26$
mainland	0.9 $\pm$ [-0.13, 1.93]	Continent vs island	$t_9 = 1.97, p = 0.08$	$N_{articles} = 8$ $N_{species} = 9$ $N_{observations} = 13$	$I^2_{total} = 82.8$ $I^2_{article} = 56.7$ $I^2_{obs} = 26$
Abundance with/without cats All species Ground foraging SMD					
ground (Intercept)	-0.03 $\pm$ [-1.2, 1.15]	Ground foraging	$t_8 = -0.05, p = 0.96$	$N_{articles} = 4$ $N_{species} = 4$ $N_{observations} = 4$	$I^2_{total} = 53.3$ $I^2_{article} = 53.3$ $I^2_{obs} = 0$
other	-0.57 $\pm$ [-1.82, 0.69]	Ground foraging	$t_{13} = -0.98, p = 0.35$	$N_{articles} = 7$ $N_{species} = 9$ $N_{observations} = 11$	$I^2_{total} = 53.3$ $I^2_{article} = 53.3$ $I^2_{obs} = 0$
Abundance on islands with/without cats All species Ground foraging lnOR					
ground (Intercept)	0.07 $\pm$ [-2.32, 2.46]	Ground foraging	$t_5 = 0.07, p = 0.94$	$N_{articles} = 4$ $N_{species} = 4$ $N_{observations} = 4$	$I^2_{total} = 43.4$ $I^2_{article} = 43.4$ $I^2_{obs} = 0$
other	-0.33 $\pm$ [-2.98, 2.32]	Ground foraging	$t_9 = -0.28, p = 0.78$	$N_{articles} = 4$ $N_{species} = 7$ $N_{observations} = 7$	$I^2_{total} = 43.4$ $I^2_{article} = 43.4$ $I^2_{obs} = 0$
Abundance temporal association All species Ground foraging Zr					
ground (Intercept)	-0.81 $\pm$ [-1.78, 0.15]	Ground foraging	$t_9 = -1.9, p = 0.09$	$N_{articles} = 3$ $N_{species} = 3$ $N_{observations} = 3$	$I^2_{total} = 81.6$ $I^2_{article} = 59.9$ $I^2_{obs} = 22$
other	1.04 $\pm$ [0.1, 1.99]	Ground foraging	$t_{17} = 2.33, p = 0.03$	$N_{articles} = 9$ $N_{species} = 10$ $N_{observations} = 16$	$I^2_{total} = 81.6$ $I^2_{article} = 59.9$ $I^2_{obs} = 22$

Table S6. Model summaries for intercept-only (main text) models as well as models evaluating influence of habitat, body mass, and continents versus islands. Model estimates $\pm$ 95% CIs and test statistics are reported along with sample sizes and residual unexplained heterogeneity (I^2), decomposed by hierarchical model levels (article and observation). The effect size used is given in the heading of each model (SMD=standardized mean difference or Hedges' g; Zr=correlation coefficient; lnOR=log odds ratio).

Term	Estimate	Moderator	Test statistics	Sample size	Residual heterogeneity
Abundance with/without cats All species Ground nesting/resting SMD					
ground (Intercept)	0.3 $\pm$ [-0.75, 1.34]	Ground nesting/resting	$t_8 = 0.65, p = 0.53$	$N_{articles} = 3$ $N_{species} = 6$ $N_{observations} = 6$	$I^2_{total} = 34.8$ $I^2_{article} = 34.8$ $I^2_{obs} = 0$
other	-1.05 $\pm$ [-2.14, 0.05]	Ground nesting/resting	$t_{13} = -2.07, p = 0.06$	$N_{articles} = 8$ $N_{species} = 7$ $N_{observations} = 9$	$I^2_{total} = 34.8$ $I^2_{article} = 34.8$ $I^2_{obs} = 0$
Abundance temporal association All species Ground nesting/resting Zr					
ground (Intercept)	-0.36 $\pm$ [-1.33, 0.61]	Ground nesting/resting	$t_9 = -0.84, p = 0.42$	$N_{articles} = 3$ $N_{species} = 3$ $N_{observations} = 4$	$I^2_{total} = 86.1$ $I^2_{article} = 66.6$ $I^2_{obs} = 20$
other	0.52 $\pm$ [-0.44, 1.48]	Ground nesting/resting	$t_{17} = 1.14, p = 0.27$	$N_{articles} = 9$ $N_{species} = 10$ $N_{observations} = 15$	$I^2_{total} = 86.1$ $I^2_{article} = 66.6$ $I^2_{obs} = 20$
Abundance on islands with/without cats All species Volant lnOR					
non-volant (Intercept)	0.16 $\pm$ [-1.88, 2.21]	Volant	$t_5 = 0.2, p = 0.85$	$N_{articles} = 4$ $N_{species} = 6$ $N_{observations} = 6$	$I^2_{total} = 39.7$ $I^2_{article} = 39.7$ $I^2_{obs} = 0$
volant	-0.73 $\pm$ [-3.9, 2.45]	Volant	$t_5 = -0.59, p = 0.58$	$N_{articles} = 3$ $N_{species} = 5$ $N_{observations} = 5$	$I^2_{total} = 39.7$ $I^2_{article} = 39.7$ $I^2_{obs} = 0$