

Table S6. Model summaries for intercept-only main-text models across all species and by taxonomic class. Model estimates $\pm$ 95% CIs and test statistics are reported along with sample sizes and residual unexplained heterogeneity (I^2), decomposed by hierarchical model levels (article and observation). The effect size used is given in the heading of each model (SMD=standardized mean difference or Hedges' g; Zr=correlation coefficient; lnOR=log odds ratio).

Term	Estimate	Test statistics	Sample size	Residual heterogeneity
Abundance with/without cats All classes lnOR				
overall	-0.64 $\pm$ [-2.08, 0.8]	$t_8 = -1.03, p = 0.33$	$N_{articles} = 9$ $N_{species} = 11$ $N_{observations} = 11$	$I^2_{total} = 60.6$ $I^2_{article} = 60.6$ $I^2_{obs} = 0$
Abundance on islands with/without cats All classes lnOR				
overall	0.12 $\pm$ [-1.59, 1.84]	$t_5 = 0.18, p = 0.86$	$N_{articles} = 6$ $N_{species} = 8$ $N_{observations} = 8$	$I^2_{total} = 39.6$ $I^2_{article} = 39.6$ $I^2_{obs} = 0$
Abundance association All classes Zr				
overall	0.07 $\pm$ [-0.33, 0.47]	$t_{14} = 0.38, p = 0.71$	$N_{articles} = 15$ $N_{species} = 14$ $N_{observations} = 23$	$I^2_{total} = 82.7$ $I^2_{article} = 57.8$ $I^2_{obs} = 25$
Abundance spatial association All classes Zr				
overall	0.1 $\pm$ [-1.25, 1.45]	$t_3 = 0.24, p = 0.83$	$N_{articles} = 4$ $N_{species} = 2$ $N_{observations} = 4$	$I^2_{total} = 67.3$ $I^2_{article} = 33.6$ $I^2_{obs} = 34$
Abundance temporal association All classes Zr				
overall	0.06 $\pm$ [-0.42, 0.54]	$t_{10} = 0.28, p = 0.78$	$N_{articles} = 11$ $N_{species} = 13$ $N_{observations} = 19$	$I^2_{total} = 85.3$ $I^2_{article} = 60.5$ $I^2_{obs} = 25$
Abundance with/without cats Mammals lnOR				
overall	-1.06 $\pm$ [-4.05, 1.93]	$t_3 = -1.13, p = 0.34$	$N_{articles} = 4$ $N_{species} = 6$ $N_{observations} = 6$	$I^2_{total} = 62.6$ $I^2_{article} = 62.6$ $I^2_{obs} = 0$
Abundance on islands with/without cats Mammals lnOR				
overall	-0.25 $\pm$ [-20.55, 20.04]	$t_1 = -0.16, p = 0.9$	$N_{articles} = 2$ $N_{species} = 4$ $N_{observations} = 4$	$I^2_{total} = 60.1$ $I^2_{article} = 60.1$ $I^2_{obs} = 0$
Abundance association Mammals Zr				

Table S6. Model summaries for intercept-only main-text models across all species and by taxonomic class. Model estimates $\pm$ 95% CIs and test statistics are reported along with sample sizes and residual unexplained heterogeneity (I^2), decomposed by hierarchical model levels (article and observation). The effect size used is given in the heading of each model (SMD=standardized mean difference or Hedges' g; Zr=correlation coefficient; lnOR=log odds ratio).

Term	Estimate	Test statistics	Sample size	Residual heterogeneity
overall	0.14 $\pm$ [-0.29, 0.56]	$t_{12} = 0.7, p = 0.49$	$N_{articles} = 13$ $N_{species} = 12$ $N_{observations} = 21$	$I^2_{total} = 83.4$ $I^2_{article} = 58$ $I^2_{obs} = 25$
Abundance spatial association Mammals Zr				
overall	0.1 $\pm$ [-1.25, 1.45]	$t_3 = 0.24, p = 0.83$	$N_{articles} = 4$ $N_{species} = 2$ $N_{observations} = 4$	$I^2_{total} = 67.3$ $I^2_{article} = 33.6$ $I^2_{obs} = 34$
Abundance temporal association Mammals Zr				
overall	0.15 $\pm$ [-0.38, 0.67]	$t_8 = 0.65, p = 0.53$	$N_{articles} = 9$ $N_{species} = 11$ $N_{observations} = 17$	$I^2_{total} = 86$ $I^2_{article} = 60.7$ $I^2_{obs} = 25$
Abundance with/without cats Birds lnOR				
overall	-0.46 $\pm$ [-7.58, 6.66]	$t_2 = -0.28, p = 0.81$	$N_{articles} = 3$ $N_{species} = 3$ $N_{observations} = 3$	$I^2_{total} = 74.7$ $I^2_{article} = 37.3$ $I^2_{obs} = 37$
Reproduction with/without cats All classes lnOR				
overall	-0.39 $\pm$ [-3.52, 2.75]	$t_2 = -0.53, p = 0.65$	$N_{articles} = 3$ $N_{species} = 3$ $N_{observations} = 5$	$I^2_{total} = 65.5$ $I^2_{article} = 28.2$ $I^2_{obs} = 37$
Reproduction before-after eradication All classes SMD				
overall	0.14 $\pm$ [-5.7, 5.99]	$t_1 = 0.31, p = 0.81$	$N_{articles} = 2$ $N_{species} = 3$ $N_{observations} = 3$	$I^2_{total} = 54.4$ $I^2_{article} = 54.4$ $I^2_{obs} = 0$
Reproduction association All classes Zr				
overall	-0.51 $\pm$ [-11.08, 10.06]	$t_1 = -0.62, p = 0.65$	$N_{articles} = 2$ $N_{species} = 2$ $N_{observations} = 5$	$I^2_{total} = 80.3$ $I^2_{article} = 80.3$ $I^2_{obs} = 0$
Reproduction temporal association All classes Zr				
overall	0.09 $\pm$ [-0.62, 0.8]	$t_3 = 0.41, p = 0.71$	$N_{articles} = 1$ $N_{species} = 1$ $N_{observations} = 4$	$I^2_{total} = 0$ $I^2_{article} = NA$ $I^2_{obs} = NA$

Table S6. Model summaries for intercept-only main-text models across all species and by taxonomic class. Model estimates $\pm$ 95% CIs and test statistics are reported along with sample sizes and residual unexplained heterogeneity (I^2), decomposed by hierarchical model levels (article and observation). The effect size used is given in the heading of each model (SMD=standardized mean difference or Hedges' g; Zr=correlation coefficient; lnOR=log odds ratio).

Term	Estimate	Test statistics	Sample size	Residual heterogeneity
Reproduction before-after eradication Birds SMD				
overall	0.14 $\pm$ [-5.7, 5.99]	$t_1 = 0.31, p = 0.81$	$N_{articles} = 2$ $N_{species} = 3$ $N_{observations} = 3$	$I^2_{total} = 54.4$ $I^2_{article} = 54.4$ $I^2_{obs} = 0$
Reproduction with/without cats Birds lnOR				
overall	-0.39 $\pm$ [-3.52, 2.75]	$t_2 = -0.53, p = 0.65$	$N_{articles} = 3$ $N_{species} = 3$ $N_{observations} = 5$	$I^2_{total} = 65.5$ $I^2_{article} = 28.2$ $I^2_{obs} = 37$
Reproduction association Birds Zr				
overall	-0.51 $\pm$ [-11.08, 10.06]	$t_1 = -0.62, p = 0.65$	$N_{articles} = 2$ $N_{species} = 2$ $N_{observations} = 5$	$I^2_{total} = 80.3$ $I^2_{article} = 80.3$ $I^2_{obs} = 0$
Reproduction temporal association Birds Zr				
overall	0.09 $\pm$ [-0.62, 0.8]	$t_3 = 0.41, p = 0.71$	$N_{articles} = 1$ $N_{species} = 1$ $N_{observations} = 4$	$I^2_{total} = 0$ $I^2_{article} = NA$ $I^2_{obs} = NA$