

```

#
↪ =====
# LECTURE 27: van der Waals Residual Properties Calculator
# Author: Edward Maginn, CBE 20260
# Description:
#   This notebook solves the cubic van der Waals equation of
↪   state for a variety of
#   industrial gases, calculates the molar volume roots, and
↪   computes the residual
#   thermodynamic properties ( $V^R$ ,  $U^R$ ,  $H^R$ ,  $S^R$ ) for the real
↪   physical roots.
#
↪ =====

# Quietly install CoolProp if running on Google Colab
try:
    import CoolProp
except ImportError:
    import subprocess
    import sys
    subprocess.check_call([sys.executable, "-m", "pip",
↪   "install", "-q", "CoolProp"])
    print("CoolProp successfully installed.")

import numpy as np
import matplotlib.pyplot as plt
import ipywidgets as widgets
from IPython.display import display, clear_output
import CoolProp.CoolProp as CP

# Dictionary mapping friendly names to CoolProp string
↪   identifiers
fluid_dict = {
    'Methane': 'Methane',
    'Nitrogen': 'Nitrogen',
    'Oxygen': 'Oxygen',
    'Ethane': 'Ethane',
    'Ammonia': 'Ammonia',
    'Carbon Dioxide': 'CarbonDioxide',
    'Carbon Monoxide': 'CarbonMonoxide',
    'Argon': 'Argon',
    'Water': 'Water',
    'R-32': 'R32'
}

def compute_residuals_and_plot(fluid_name, T_val, T_unit, P_val,
↪   P_unit):
    fluid = fluid_dict[fluid_name]

```

```

# --- 1. Unit Conversions to Standard Math Units (Kelvin &
↪ bar) ---
if T_unit == '°C':
    T = T_val + 273.15
elif T_unit == '°F':
    T = (T_val - 32) * 5/9 + 273.15
else:
    T = T_val

if P_unit == 'atm':
    P = P_val * 1.01325
elif P_unit == 'Pa':
    P = P_val / 1e5
elif P_unit == 'MPa':
    P = P_val * 10
elif P_unit == 'psi':
    P = P_val / 14.5038
else:
    P = P_val

if T <= 0 or P <= 0:
    print("Temperature and Pressure must be greater than
↪ absolute zero.")
    return

# --- 2. Fetch Critical Properties ---
Tc = CP.PropsSI('TCRIT', fluid)
Pc_Pa = CP.PropsSI('PCRIT', fluid)
Pc = Pc_Pa / 1e5 # Convert Pa to bar

# --- 3. Calculate VDW Parameters ---
R_Lbar = 0.0831446 # L*bar/(mol*K)
R_J = 8.31446 # J/(mol*K)

a = 27 * (R_Lbar * Tc)**2 / (64 * Pc)
b = R_Lbar * Tc / (8 * Pc)

# --- 4. Solve the Cubic Equation for Roots ---
# V^3 - (b + RT/P)V^2 + (a/P)V - (ab/P) = 0
C2 = -(b + R_Lbar * T / P)
C1 = a / P
C0 = -a * b / P

roots = np.roots([1, C2, C1, C0])

# Filter out complex roots and non-physical roots (V < b)
real_roots = roots[np.isclose(roots.imag, 0)].real

```

```

real_roots = np.sort(real_roots[real_roots > b])

# --- 5. Print Output Summary & Residuals ---
print("="*65)
print(f"FLUID: {fluid_name}")
print(f"Critical State: Tc = {Tc:.2f} K | Pc = {Pc:.2f}
↪ bar")
print(f"VDW Parameters: a = {a:.4f} L2·bar/mol2 | b =
↪ {b:.5f} L/mol")
print("-" * 65)
print(f"TARGET STATE: T = {T:.2f} K | P = {P:.2f} bar")

V_ideal = R_Lbar * T / P
print(f"Ideal Gas Vol: V_ig = {V_ideal:.4f} L/mol")
print("="*65)

for i, root in enumerate(real_roots):
    # Determine phase label
    if len(real_roots) == 1:
        phase = "Supercritical Fluid" if T > Tc else
↪ ("Liquid" if root < 5*b else "Vapor")
    elif len(real_roots) == 3:
        if i == 0: phase = "Liquid Root"
        elif i == 1: phase = "Unstable Root (Non-Physical)"
        else: phase = "Vapor Root"

    print(f"\n[{phase}]\n")
    print(f" Molar Volume (V) = {root:.4f} L/mol")

    # Calculate Residuals for physical roots
    if phase != "Unstable Root (Non-Physical)":
        # 1. Residual Volume (L/mol)
        V_R = root - V_ideal

        # 2. Residual Internal Energy (J/mol)
        # U^R = -a/V (in L·bar/mol) -> multiply by 100 to
        ↪ get J/mol
        U_R = (-a / root) * 100

        # 3. Residual Enthalpy (J/mol)
        # H^R = U^R + P(V - V_ideal) -> PV terms must be
        ↪ converted to J/mol
        H_R = U_R + 100 * P * V_R

        # 4. Residual Entropy (J/(mol K))
        # S^R = R * ln[ P(V-b) / RT ]
        arg = P * (root - b) / (R_Lbar * T)
        S_R = R_J * np.log(arg)

```

```

        print(f"    Residual Volume      (V^R) = {V_R:.4f}
              ↪ L/mol")
        print(f"    Residual Internal (U^R) = {U_R:.1f}
              ↪ J/mol")
        print(f"    Residual Enthalpy (H^R) = {H_R:.1f}
              ↪ J/mol")
        print(f"    Residual Entropy  (S^R) = {S_R:.3f}
              ↪ J/(mol·K)")

    print("\n" + "="*65)

    # --- 6. Plotting the P-V Diagram ---
    V_max = max(V_ideal * 1.5, 10 * b)
    if len(real_roots) == 3:
        V_max = max(V_max, real_roots[2] * 1.3)

    V_arr = np.linspace(b * 1.02, V_max, 2000)

    P_iso = (R_Lbar * T) / (V_arr - b) - a / (V_arr**2)
    P_crit = (R_Lbar * Tc) / (V_arr - b) - a / (V_arr**2)

    plt.figure(figsize=(9, 6))

    # Plot Isotherms
    plt.plot(V_arr, P_crit, 'r--', lw=1.5, label=f'Critical
    ↪ Isotherm (Tc = {Tc:.1f} K)')
    plt.plot(3*b, Pc, 'rs', markersize=6, label='Critical
    ↪ Point')
    plt.plot(V_arr, P_iso, 'b-', lw=2, label=f'Target Isotherm
    ↪ (T = {T:.1f} K)')
    plt.axhline(P, color='k', linestyle=':', lw=1.5,
    ↪ label=f'Target Pressure (P = {P:.1f} bar)')

    # Plot roots
    for i, root in enumerate(real_roots):
        if len(real_roots) == 3 and i == 1:
            plt.plot(root, P, 'wo', markeredgecolor='k',
            ↪ markersize=8, label='Unstable Root' if i==1 else "")
        else:
            plt.plot(root, P, 'go', markersize=8,
            ↪ label='Physical Root(s)' if i==0 else "")

    plt.ylim(0, max(P * 1.5, Pc * 1.5))
    plt.xlim(0, V_max)
    plt.xlabel('Molar Volume, V (L/mol)', fontsize=12)
    plt.ylabel('Pressure, P (bar)', fontsize=12)

```

```

plt.title(f'van der Waals Properties for {fluid_name}',
↪ fontsize=14, fontweight='bold')
plt.legend(loc='upper right')
plt.grid(True, linestyle='--', alpha=0.6)

plt.show()

# --- 7. Interactive UI Setup ---
style = {'description_width': 'initial'}

fluid_dropdown =
↪ widgets.DropDown(options=list(fluid_dict.keys()),
↪ value='Methane', description='Fluid:')

T_input = widgets.FloatText(value=300,
↪ description='Temperature:', style=style,
↪ layout=widgets.Layout(width='200px'))
T_unit = widgets.DropDown(options=['K', '°C', '°F'], value='K',
↪ layout=widgets.Layout(width='80px'))

P_input = widgets.FloatText(value=100, description='Pressure:',
↪ style=style, layout=widgets.Layout(width='200px'))
P_unit = widgets.DropDown(options=['bar', 'atm', 'Pa', 'MPa',
↪ 'psi'], value='bar', layout=widgets.Layout(width='80px'))

T_box = widgets.HBox([T_input, T_unit])
P_box = widgets.HBox([P_input, P_unit])

ui = widgets.VBox([fluid_dropdown, T_box, P_box])
out = widgets.interactive_output(compute_residuals_and_plot, {
    'fluid_name': fluid_dropdown,
    'T_val': T_input,
    'T_unit': T_unit,
    'P_val': P_input,
    'P_unit': P_unit
})

display(ui, out)

```

VBox(children=(DropDown(description='Fluid:', options=('Methane', 'Nitrogen

Output())