

Timeline Check In

Macey Hartmann and Erick Morales Oyola

2026-05-22

Setup

Packages

We loaded packages for data wrangling, file organization, missing-data exploration, and visualization. The main workflow uses `tidyverse` for cleaning and summarizing vegetation data, `here` for file paths, and plotting packages for exploratory figures.

Data

We used two vegetation datasets. The main dataset, `veg.csv`, contains vegetation survey observations, including species identity, percent cover, cover category, habitat type, transect, site, and year. The metadata file, `vp_veg_metadata.csv`, provides supporting information about vegetation transects, including the number of quadrats sampled within each transect-year.

Ideas for analysis and background

Our project focuses on salt marsh vegetation at North Campus Open Space (NCOS), with the goal of understanding which plant species characterize this restored wetland habitat and how those patterns have changed across survey years. Rather than treating vegetation cover as only a set of numbers, we are using percent cover to describe how the salt marsh is developing as a plant community over time. In this sense, our project asks how restoration becomes visible through the species that take root, persist, expand, or decline across the marsh.

Salt marsh vegetation is especially useful for understanding restoration trajectory because these plant communities are shaped by strong environmental filters, including salinity, tidal flooding, saturated soils, and seasonal changes in water availability. In southern California salt marshes, native halophytes such as pickleweed (*Salicornia pacifica*) and saltgrass (*Distichlis spicata*)

often help define the structure and identity of the habitat (Ferren 1985; North Campus Open Space 2025). Because these species are adapted to stressful coastal wetland conditions, changes in their mean percent cover can help indicate whether the restored salt marsh is developing toward a recognizable native marsh community.

This question is particularly relevant at NCOS because the site was restored from the former Ocean Meadows Golf Course into a mosaic of wetland and upland habitats, including salt marsh, brackish marsh, seasonal wetlands, and transitional habitats (Cheadle Center for Biodiversity and Ecological Restoration and Environmental Science Associates 2016; Rickard 2024). Broader NCOS monitoring reports evaluate restoration progress across habitat types, while our project narrows in on salt marsh vegetation specifically. By focusing on dominant species and native/non-native cover through time, we can connect the survey data to larger questions about restoration progress, habitat identity, and community development.

Previous work on California wetland restoration has shown that vegetation establishment can be slow, variable, and strongly shaped by local site conditions such as elevation, soil salinity, soil moisture, planting, and adaptive management (Beheshti et al. 2023). This matters for our analysis because shifts in dominant plant cover are not just visual patterns in a graph. They may reflect ecological processes such as native plant establishment, non-native plant expansion, hydrologic change, disturbance, or interannual variability in rainfall and growing conditions.

Ecological background

Salt marshes are dynamic coastal wetlands where plant communities exist at the edge of land and water. Their vegetation patterns are shaped by both physical stress and biological interactions, making them useful indicators of environmental conditions and restoration progress. In southern California, seasonal rainfall and dry summers can influence freshwater input, soil moisture, salinity, and plant growth, which may affect how vegetation cover changes from year to year (Ferren 1985).

At NCOS, we expect dominant salt marsh species to include native salt-tolerant plants such as *Salicornia pacifica* and *Distichlis spicata*. The NCOS coastal salt marsh habitat page describes pickleweed as usually the dominant species in coastal salt marshes and also identifies saltgrass, alkali heath, fleshy jaumea, and California saltbush as characteristic salt marsh plants (North Campus Open Space 2025). These expectations give us a useful ecological reference point for interpreting our data: if the restored salt marsh is developing toward a recognizable coastal salt marsh community, we would expect these native halophytes to contribute strongly to vegetation cover.

Native and non-native cover are also important to examine because restoration success depends not only on whether plants are present, but on which plants are shaping the habitat. A marsh with high total vegetation cover may still be ecologically different depending on whether that cover is dominated by native salt marsh species or by non-native plants. NCOS monitoring reports describe vegetation cover, native biodiversity, and relative native cover as important

indicators of restoration progress (Rickard 2024). Our project builds from that monitoring context by focusing more closely on the species-level patterns that contribute to salt marsh identity through time.

Planned analyses

Our first analysis will identify which plant species most strongly characterize salt marsh and closely related brackish marsh vegetation at NCOS. To do this, we will summarize mean percent cover for each plant species across survey years and transects, then rank species from highest to lowest mean cover. We will use the 90th percentile of mean percent cover as a cutoff for identifying dominant species, which gives us a clearer mathematical basis than selecting species only by visual inspection.

Our second analysis will examine how dominant species cover changes across survey years. After identifying species at or above the 90th percentile of mean percent cover across the full filtered dataset, we will summarize their mean percent cover by year and visualize them with a stacked bar plot. In this figure, the y-axis represents the summed mean percent cover of the dominant species in each survey year, so bar height shows how much cover those high-contributing species represent through time.

Our third analysis will focus specifically on non-native species in the filtered salt marsh/brackish marsh dataset. Rather than only comparing total native and non-native cover, we will identify the five non-native species with the highest mean percent cover across all survey years, then summarize their mean percent cover by year. We will visualize these species with a stacked bar plot so we can see whether non-native cover is being driven by the same species through time or whether different non-native plants become more important in different years. This analysis is useful because a simple native/non-native proportion plot can show the relative amount of non-native cover, but it does not show which species are responsible for that pattern. By focusing on the top non-native species, we can connect non-native cover to actual plant identities and better interpret whether non-native vegetation reflects a consistent community pattern, a temporary increase in a few species, or a shift in the plants contributing to marsh composition.

Our fourth analysis will provide broader context by examining total vegetation cover across survey years. Unlike the dominant-species figure, this analysis includes all native and non-native plant species in the filtered salt marsh/brackish marsh dataset. We will summarize total mean vegetation cover by year and visualize it as a line plot with standard error, which will help us interpret whether changes in species composition are happening alongside broader increases or decreases in overall vegetation cover.

Together, these analyses connect our figures to the main project questions. The species-rank plot identifies the plants that most strongly characterize the habitat, the dominant-species plot shows how the top five species change through time, the proportional native/non-native plot shows how plant origin contributes to habitat identity, and the dominant non-native species

plot shows which non-native plants are driving non-native cover across survey years. The total-cover plot then provides broader context for overall vegetation development across the monitoring period.

Data cleaning and wrangling

We filtered the vegetation dataset to focus on salt marsh and closely related brackish marsh vegetation, then limited the surveys to August and September because late summer and early fall are when vegetation cover is expected to be highest and most comparable across years. We used `complete()` to make implicit absences explicit as zero percent cover, then joined the vegetation data with the metadata file so we could account for the number of quadrats sampled in each transect-year. Finally, we filtered to native and non-native plant cover categories so our analyses focused on vegetation rather than bare ground, thatch, or other non-plant cover types.

```
saltmarsh <- veg |>
  # filter to salt marsh habitat only
  filter(str_detect(habitat, "Salt Marsh") | str_detect(habitat, "Brackish
    ↪ Marsh")) |>
  # parse date column and filter to August and September
  mutate(date = as.Date(date),
         month = month(date)) |>
  filter(month %in% c(8, 9))
```

```
saltmarsh_complete <- saltmarsh |>
  complete(year, transect_name, psoc, transect_distance,
         fill = list(percent_cover = 0)) |>
  group_by(year, transect_name, psoc, cover_category) |>
  summarize(sum_pc = sum(percent_cover, na.rm = TRUE)) |>
  ungroup() |>
  unite("year_pool", year, transect_name, remove = FALSE) |>
  left_join(metadata, by = "year_pool") |>
  select(-year.y, -transect_name.y) |>
  rename(year = year.x,
         transect_name = transect_name.x) |>
  mutate(mean_pc = sum_pc / num_quad)
```

```
saltmarsh_plants <- saltmarsh_complete |>
  filter(cover_category %in% c("NATIVE COVER", "NON-NATIVE COVER")) |>
  mutate(species_label = str_replace_all(psoc, "_", " "))
```

Planned visualizations and descriptive analyses

For this stage of the project, our work focuses mainly on descriptive analyses and visualizations that help us understand salt marsh/brackish marsh vegetation patterns at NCOS. These summaries allow us to identify dominant species, compare native and non-native cover, and explore how vegetation cover changes across survey years. For the final project, we plan to add a simple statistical analysis using linear models to test whether vegetation cover changes through time.

Our planned statistical analysis will use survey year as the explanatory variable and mean percent cover as the response variable. We will first test whether total mean vegetation cover changes across years, then consider separate statistical tests for native cover, non-native cover, or individual dominant species if the data structure supports those comparisons. These tests will help us move beyond visual interpretation by asking whether the patterns we see in the figures suggest directional changes in vegetation cover over the monitoring period.

```
#Creating a species-level summary table for the rank plot
species_rank <- saltmarsh_plants |>
  #Grouping by species and cover category so each species keeps its
  ↪ native/non-native status
  group_by(psoc, species_label, cover_category) |>
  #Calculating mean percent cover, standard error, and sample size for each
  ↪ species
  summarize(mean_pc = mean(mean_pc, na.rm = TRUE),
             se_pc = sd(mean_pc, na.rm = TRUE) / sqrt(n()),
             n = n()) |>
  #Removing grouping so later steps do not accidentally stay grouped
  ungroup() |>
  #Ordering species from highest to lowest mean percent cover
  arrange(desc(mean_pc))
```

```
# Calculating the 90th percentile of species mean percent cover
dominance_cutoff <- quantile(species_rank$mean_pc,
                             probs = 0.90,
                             na.rm = TRUE)

# Identifying dominant species as those at or above the 90th percentile
dominant_species <- species_rank |>
  filter(mean_pc >= dominance_cutoff) |>
  pull(psoc)

# Creating a table that ranks dominant species from highest to lowest mean
↪ cover
```

```

dominant_species_order <- species_rank |>
  filter(psoc %in% dominant_species) |>
  arrange(desc(mean_pc)) |>
  select(psoc, species_label, overall_mean_pc = mean_pc)

# Summarizing dominant species cover by year
dominant_by_year <- saltmarsh_plants |>
  # Keeping only species at or above the 90th percentile cutoff
  filter(psoc %in% dominant_species) |>
  # Grouping by year, transect, species, and cover category
  group_by(year, transect_name, psoc, species_label, cover_category) |>
  # Calculating species cover within each transect-year
  summarize(sum_pc = sum(mean_pc, na.rm = TRUE)) |>
  ungroup() |>
  # Averaging across transects within each year and species
  group_by(year, psoc, species_label, cover_category) |>
  summarize(mean_pc = mean(sum_pc, na.rm = TRUE)) |>
  ungroup() |>
  # Joining the overall species rank back into the plotting dataset
  left_join(dominant_species_order, by = c("psoc", "species_label")) |>
  # Ordering species so the highest-cover species appears at the bottom of
  ↪ the stack
  mutate(species_label = factor(species_label,
                                levels =
                                ↪ rev(dominant_species_order$species_label)))

#Keeping the top 20 species for the rank plot
species_rank_top20 <- species_rank |>
  filter(n > 1) |>
  slice_max(mean_pc, n = 20)

#Creating color palette for species within the 90th percentile
species_rank_palette <- c(
  #Native species
  "Salicornia pacifica"      = "#00441B",
  "Distichlis spicata"      = "#238B45",
  "Atriplex lentiformis"    = "#74C476",
  "Distichlis littoralis"   = "#006D2C",
  "Suaeda taxifolia"        = "#41AB5D",
  "Eleocharis macrostachya" = "#C4E8D9",
  "Elymus triticoides"      = "#A1D99B",
  "Isocoma menziesii"       = "#2CA25F",
  "Frankenia salina"        = "#99D8C9",

```

```

#Non-native species
"Raphanus sativus"      = "#D85A30",
"Festuca perennis"      = "#B8322A"
)

```

```

native_nonnative_points <- saltmarsh_plants |>
  # Grouping by year, transect, and cover category to keep transect-level
  ↪ observations
  group_by(year, transect_name, cover_category) |>
  # Summing native or non-native cover within each transect-year
  summarize(total_pc = sum(mean_pc, na.rm = TRUE)) |>
  ungroup()

native_nonnative_raw <- saltmarsh_plants |>
  # Starting with transect-level native and non-native cover values
  group_by(year, transect_name, cover_category) |>
  summarize(total_pc = sum(mean_pc, na.rm = TRUE)) |>
  ungroup() |>
  # Grouping by year and cover category to calculate yearly means
  group_by(year, cover_category) |>
  # Calculating mean cover and standard error for each cover category in each
  ↪ year
  summarize(mean_cover = mean(total_pc, na.rm = TRUE),
            se_cover = sd(total_pc, na.rm = TRUE) / sqrt(n())) |>
  ungroup()

```

```

#Identifying the five non-native species with the highest average cover
top_nonnative_species <- saltmarsh_plants |>
  filter(cover_category == "NON-NATIVE COVER") |>
  group_by(psoc) |>
  summarize(mean_pc = mean(mean_pc, na.rm = TRUE)) |>
  ungroup() |>
  slice_max(mean_pc, n = 5) |>
  pull(psoc)

#Summarizing top non-native species by year
top_nonnative_by_year <- saltmarsh_plants |>
  filter(cover_category == "NON-NATIVE COVER",
        psoc %in% top_nonnative_species) |>
  mutate(species_label = str_replace_all(psoc, "_", " ")) |>
  group_by(year, transect_name, psoc, species_label) |>

```

```

summarize(sum_pc = sum(mean_pc, na.rm = TRUE)) |>
ungroup() |>
group_by(year, psoc, species_label) |>
summarize(mean_pc = mean(sum_pc, na.rm = TRUE)) |>
ungroup()

```

```

# Creating transect-level total vegetation cover values for underlying points
total_cover_points <- saltmarsh_plants |>
  group_by(year, transect_name) |>
  summarize(total_pc = sum(mean_pc, na.rm = TRUE)) |>
  ungroup()

# Summarizing total vegetation cover by year for the line and ribbon
total_cover <- total_cover_points |>
  group_by(year) |>
  summarize(mean_total = mean(total_pc, na.rm = TRUE),
            se_total = sd(total_pc, na.rm = TRUE) / sqrt(n())) |>
  ungroup()

```

Visualizations directly relevant to answering questions

The following four figures directly address our project questions by connecting plant identity, cover, and origin through time. Figure 1 focuses on the five dominant plant species overall, Figure 2 compares the proportional contribution of native and non-native cover, Figure 3 examines which non-native species contribute most to non-native cover, and Figure 4 ranks the top species by mean cover to justify our dominant-species cutoff.

Figure 1: Dominant species percent cover across survey years

```

#Creating Figure 1: dominant species cover across survey years
fig1 <- ggplot(data = dominant_by_year,
               aes(x = year,
                   y = mean_pc,
                   fill = species_label,
                   group = species_label)) +
  #Creating stacked bars to show how dominant species contribute to cover
  ↪ each year
  geom_col(position = "stack") +
  #Applying custom species colors and keeping the legend ordered from highest
  ↪ to lowest cover

```

```

scale_fill_manual(values = species_rank_palette,
                  breaks = dominant_species_order$species_label) +
#Showing each survey year as its own x-axis break
scale_x_continuous(breaks = seq(2018, 2025, 1)) +
#Adding axis labels and legend title
labs(x = "Survey year",
     y = "Mean percent cover (%)",
     fill = "Species") +
#Using a clean theme for readability
theme_minimal() +
#Formatting the legend and italicizing species names
theme(legend.position = "bottom",
      legend.text = element_text(face = "italic"),
      legend.title = element_text(face = "bold"))

#Displaying Figure 1
fig1

```




Figure 1: Mean percent cover of dominant salt marsh/brackish marsh plant species at NCOS across survey years, averaged across transects within each year. Dominant species were identified as species at or above the 90th percentile of mean percent cover across all survey years. Bar height reflects total mean cover of the dominant species and varies across years. Green shades indicate native species; coral indicates non-native species.

The original figure was unclear about what the y-axis represented, so we revised the wrangling approach to make the summary more interpretable. Rather than averaging directly across all observations, we first calculated transect-level totals for each dominant species within each year, then averaged those values across transects. This means the y-axis represents mean percent cover averaged across transects within each survey year, and bar height varies across years based on the summed cover of the dominant species. Dominant species were identified using the 90th percentile of mean percent cover across the full filtered dataset, rather than by visual inspection. The “Other” category was removed so the figure focuses only on species that met the dominant-species cutoff, and species are stacked within each year from highest to lowest mean cover.

Figure 2: Proportional native and non-native cover across survey years

```
#Creating Figure 2: native and non-native vegetation cover across survey
  ↪ years
fig2 <- ggplot(data = native_nonnative_raw,
               aes(x = year,
                   y = mean_cover,
                   color = cover_category,
                   fill = cover_category)) +
#Adding faint points to show the underlying transect-level observations
#Each point represents total native or non-native cover within one
  ↪ transect-year
geom_point(data = native_nonnative_points,
           aes(x = year,
               y = total_pc,
               color = cover_category),
           inherit.aes = FALSE,
           alpha = 0.35,
           size = 1.8,
           position = position_jitter(width = 0.08, height = 0)) +
#Adding shaded ribbons to show ± 1 standard error around each yearly mean
geom_ribbon(aes(ymin = mean_cover - se_cover,
                ymax = mean_cover + se_cover),
           alpha = 0.2,
           color = NA) +
#Adding lines to connect yearly mean percent cover values through time
geom_line(linewidth = 1) +
#Adding larger points to show the yearly mean percent cover values
geom_point(size = 3) +
#Using green for native cover and coral/red for non-native cover
scale_color_manual(values = c("NATIVE COVER" = "#1D9E75",
                              "NON-NATIVE COVER" = "#D85A30")) +
#Matching the ribbon fill colors to the line and point colors
scale_fill_manual(values = c("NATIVE COVER" = "#1D9E75",
                             "NON-NATIVE COVER" = "#D85A30")) +
#Showing each survey year as its own x-axis break
scale_x_continuous(breaks = seq(2018, 2025, 1)) +
#Adding axis labels and legend titles
labs(x = "Survey year",
     y = "Mean percent cover (± 1 SE)",
     color = "Cover category",
     fill = "Cover category") +
#Using a clean theme for readability
```

```

theme_minimal() +
#Moving the legend below the plot and bolding the legend title
theme(legend.position = "bottom",
      legend.title = element_text(face = "bold"))

#Displaying Figure 2
fig2

```




Figure 2: Mean percent cover (± 1 SE) of native and non-native vegetation in salt marsh habitat at NCOS across survey years. Points represent means and shaded bands represent ± 1 standard error. Native cover is shown in green; non-native cover in coral.

The original proportional stacked bar was nearly identical to panel (b) of figure 9 in the NCOS monitoring report, making it redundant. We replaced it with a line plot showing raw mean percent cover of native and non-native vegetation over time, with ± 1 standard error ribbons

around each line. This is distinct from the monitoring report because it uses a continuous line geometry that emphasizes temporal trend, places both cover categories on the same axis for direct visual comparison, and includes a measure of uncertainty that the monitoring report's bar charts do not show.

Figure 3: Dominant non-native species percent cover across survey years

```
# Creating a red/coral color palette for the top non-native species
# Each color corresponds to one of the five highest-cover non-native species
nonnative_palette <- c("#7F1D1D", "#B8322A", "#D85A30", "#E9825A", "#F3B49C")

# Creating Figure 3: top non-native species cover across survey years
fig_3 <- ggplot(data = top_nonnative_by_year,
               aes(x = year,
                  y = mean_pc,
                  fill = species_label)) +
  # Creating stacked bars to show how the top non-native species contribute
  # to cover each year
  geom_col(position = "stack") +
  # Applying the custom red/coral palette to distinguish non-native species
  scale_fill_manual(values = nonnative_palette) +
  # Showing each survey year as its own x-axis break
  scale_x_continuous(breaks = seq(2018, 2025, 1)) +
  # Adding axis labels and legend title
  labs(x = "Survey year",
       y = "Mean percent cover (%)",
       fill = "Non-native species") +
  # Using a clean theme for readability
  theme_minimal() +
  # Moving the legend below the plot and italicizing species names
  theme(legend.position = "bottom",
        legend.text = element_text(face = "italic"),
        legend.title = element_text(face = "bold"))

# Displaying Figure 3
fig_3
```




Figure 3: Mean percent cover of the five most common non-native plant species in the filtered salt marsh/brackish marsh dataset across survey years, averaged across transects within each year. Bar height reflects the summed mean cover of these non-native species in each year.

Figure 4: Species rank plot

```
# Creating Figure 4: species rank plot
# This figure ranks the top 20 plant species by mean percent cover

# Setting the x-axis location of the dashed cutoff line
# This cutoff comes from the 90th percentile of species mean percent cover
cutoff_x <- dominance_cutoff

fig4 <- ggplot(data = species_rank_top20,
               aes(x = mean_pc,
                   y = reorder(species_label, mean_pc),
                   color = cover_category)) +
  # Adding points and horizontal error bars for each species
  # Points show mean percent cover and bars show ± 1 standard error
  geom_pointrange(aes(xmin = mean_pc - se_pc,
                      xmax = mean_pc + se_pc)) +
```

```

# Adding a dashed vertical line to show the 90th percentile dominance
↪ cutoff
geom_vline(xintercept = cutoff_x,
           linetype = "dashed",
           color = "gray50") +
# Using green for native species and coral/red for non-native species
scale_color_manual(values = c("NATIVE COVER" = "#1D9E75",
                              "NON-NATIVE COVER" = "#D85A30")) +
# Adding axis labels and legend title
labs(x = "Mean percent cover (± 1 SE)",
     y = "Species",
     color = "Cover category") +
# Using a clean theme for readability
theme_minimal() +
# Italicizing species names and placing the legend below the plot
theme(axis.text.y = element_text(face = "italic", size = 9),
      legend.position = "bottom",
      legend.title = element_text(face = "bold"))

# Displaying Figure 4
fig4

```




Figure 4: Mean percent cover (± 1 SE) of the top 20 plant species recorded in the filtered salt marsh/brackish marsh dataset, averaged across transects and survey years and ranked from highest to lowest mean cover. Points represent means and horizontal bars represent ± 1 standard error. Green indicates native species; coral indicates non-native species. Dashed line marks the 90th percentile cutoff used to identify dominant species for figure 1.

The dashed line represents the 75th percentile of mean percent cover across all species in the filtered dataset. Species to the right of this line were classified as dominant because their average cover was higher than at least 75% of recorded plant species. This gives the dominant-species selection a clearer mathematical basis than choosing a visual cutoff by eye.

Other exploratory visualizations

Figure 5: Total vegetation cover across survey years

```

# Creating Figure 5: total vegetation cover across survey years
fig5 <- ggplot(data = total_cover,
               aes(x = year,
                   y = mean_total)) +
# Adding faint points to show underlying transect-level observations
# Each point represents total native + non-native vegetation cover within
  ↪ one transect-year
geom_point(data = total_cover_points,
           aes(x = year,
               y = total_pc),
           inherit.aes = FALSE,
           alpha = 0.35,
           size = 1.8,
           position = position_jitter(width = 0.08, height = 0),
           color = "#6B5B95") +
# Adding a shaded ribbon to show ± 1 standard error around yearly mean
  ↪ total cover
geom_ribbon(aes(ymin = mean_total - se_total,
               ymax = mean_total + se_total),
           fill = "#6B5B95",
           alpha = 0.2) +
# Adding a line to connect yearly mean total vegetation cover through time
geom_line(color = "#6B5B95",
          linewidth = 1) +
# Adding larger points to show yearly mean total vegetation cover values
geom_point(color = "#6B5B95",
           size = 3) +
# Adding axis labels
labs(x = "Survey year",
     y = "Total mean percent cover (%)") +
# Using a clean theme for readability
theme_minimal()

# Displaying Figure 5
fig5

```




Figure 5: Total mean percent vegetation cover (native + non-native combined) in salt marsh habitat at NCOS across survey years. Shaded band represents ± 1 standard error around the mean.

Plan for elective (if completing)

For our advanced elective, we are creating a National Geographic-inspired educational magazine about the salt marsh plants of the North Campus Open Space (NCOS). The magazine will translate our vegetation analysis into a more accessible and visual format for younger students and community members, introducing readers to the dominant plant species that characterize salt marsh habitat at NCOS while explaining why salt marshes matter for restoration, wildlife habitat, and coastal resilience. The magazine will be designed in Canva, printed in color at the dorm print facilities, and delivered to Ms. McCarter's fourth grade classroom with one copy per table group (approximately 3–4 copies total).

Elective concept

The magazine is inspired by educational nature publications such as *National Geographic Kids* and is designed for a fourth grade audience. It will include species spotlight pages profiling the dominant salt marsh plants with illustrated profiles, fun facts, and native/non-native status; a “Meet the Marsh” illustrated map of NCOS showing habitat zones; a data feature communicating our project findings in an accessible visual format; a “Native vs. Non-Native vs. Invasive” explanation page; a salt marsh vocabulary word search; a “What Plant Are You?” personality flowchart; and a how-to-get-involved page describing how students can visit and learn more about NCOS.

Draft sections (include some description/image of drafts)

We have finalized the magazine structure, section outline, and visual style. Based on our preliminary data exploration, we have identified the dominant salt marsh species to feature: *Salicornia pacifica*, *Distichlis spicata*, *Atriplex lentiformis*, and *Raphanus sativus*. Each species will receive a spotlight page with a drawing or photograph, its native or non-native status, and a brief description of its role in the marsh community. For images, we will use Creative Commons licensed photographs from iNaturalist, Wikimedia Commons, and the USDA PLANTS Database, supplement with our own photographs taken at NCOS, and create original botanical-style illustrations for species that are difficult to photograph clearly.



Figure 6: Cover page draft



Figure 7: Table of contents draft

ISSUE 01 /

TABLE OF CONTENTS



04.

ABOUT

05.

IMPORTANT
DEFINITIONS

06.

MEET THE MARSH:
A MAP OF NCOS

07

NATIVE VS NON-NATIVE VS
INVASIVE

08.-09.

ECO WORD SEARCH

10.

THE DATA

11.

ENVIRONMENTAL
CROSSWORD

12-13

WHAT PLANT ARE YOU?

14.

SPECIES SPOTLIGHTS

15.

HOW TO VISIT

Figure 8: Table of contents draft 2

ABOUT



Educate, Inspire, Excite


Eco-Explorers Magazine is a children's magazine created for the students of Santa Barbara and Goleta. This edition dives into the salt marsh habitat of the North Campus Open Space (NCOS) — a restored wetland right in our own backyard — and the remarkable plants that call it home. From dominant native species like *Salicornia pacifica* and *Distichlis spicata* to non-native invaders, readers will discover who lives in the marsh, why it matters, and how they can help protect it. Through species spotlights, games, maps, and activities, this magazine aims to educate, inspire, and excite the next generation of environmental stewards.

/ ECO-EXPLORERS

Figure 9: About page draft



Figure 10: Word search draft



CONSERVATION CROSSWORD

Down:

1. Abbreviation for the restored wetland near UCSB (abbr.)
2. Scientists count plants by doing this in the field
4. To bring a habitat back to its natural state
5. A non-native species that spreads and harms the ecosystem
7. Caused by the rise and fall of the ocean
8. A habitat where the ground is soaked with water

Across:

2. A group of plants or animals of the same kind
3. Scientists measure plant ___ to see how much ground is filled
6. The underground parts of a plant that absorb water
9. Where a river meets the ocean, creating a mix of fresh and salt water
10. A plant that naturally belongs in an ecosystem
11. Information scientists collect during a study
12. Someone who protects and cares for the environment

Figure 11: Crossword draft

Plan for the next two weeks

Over the next two weeks, we will use our preliminary analysis to select the species featured in the magazine. We will also continue revising our figures so the main results are clear enough to translate into a more visual and accessible format for younger students and community readers.

By Week 9, we will have completed all remaining sections including the data feature, word search, flowchart, and how-to-get-involved page, and will have begun design and layout in Canva. By Week 10 we will finalize and polish the full magazine for presentation, revising the writing for clarity and ensuring the layout is cohesive and visually engaging. The completed magazine will be delivered to Ms. McCarter's classroom during finals week.

Next steps

Our next steps are to refine the statistical analyses and continue revising the figures so our interpretations are clear and well-supported. For the final project, we will add more complete citations, connect our results more directly to NCOS restoration goals, and continue developing the advanced elective magazine alongside the data analysis.

Before submission, we will make sure the GitHub repository includes the rendered PDF, the QMD, the data files, the bibliography file, and a README that clearly links to the rendered document.

References

- Beheshti, Kathryn M., Stephen C. Schroeter, Andres A. Deza, Daniel C. Reed, Rachel S. Smith, and Henry M. Page. 2023. "Large-Scale Field Studies Inform Adaptive Management of California Wetland Restoration." *Restoration Ecology* 31 (5): e13936. <https://doi.org/10.1111/rec.13936>.
- Cheadle Center for Biodiversity and Ecological Restoration, and Environmental Science Associates. 2016. *North Campus Open Space Restoration Project Restoration Plan*. University of California, Santa Barbara. <https://escholarship.org/uc/item/9505h60h>.
- Ferren, Wayne R. Jr. 1985. *Carpinteria Salt Marsh: Environment, History, and Botanical Resources of a Southern California Estuary*. Publication Number 4. The Herbarium, Department of Biological Sciences, University of California, Santa Barbara.
- North Campus Open Space. 2025. *Coastal Salt Marsh*. <https://www.ncos.ccber.ucsb.edu/native-plant-habitats/coastal-salt-marsh>.

Rickard, Alison. 2024. *North Campus Open Space Restoration Project Monitoring Report: Year 7, December 2024*. University of California, Santa Barbara Cheadle Center for Biodiversity; Ecological Restoration. <https://escholarship.org/uc/item/7bq618m8>.