

Salt Marsh Vegetation at North Campus Open Space

Dominant species, native/non-native cover, and restoration trajectories

Macey Hartmann and Erick Morales Oyola

2026-06-10

Table of contents

Setup	1
Introduction	1
Methods	2
Study system and datasets	2
Data cleaning and summary approach	3
Results	3
Dominant species identity	3
Dominant species cover through time	5
Native and non-native vegetation cover	5
Total vegetation cover	6
Discussion	8
References	8

Setup

Introduction

North Campus Open Space (NCOS) is a restored coastal wetland system in Santa Barbara, California, created from the former Ocean Meadows Golf Course and designed to support a

mosaic of wetland and upland habitats, including salt marsh, brackish marsh, seasonal wetlands, and transitional habitats (Cheadle Center for Biodiversity and Ecological Restoration and Environmental Science Associates 2016; Rickard 2024). Because restored wetlands develop over time, vegetation monitoring provides a useful way to evaluate whether habitat structure is moving toward intended ecological conditions. In salt marshes especially, plant cover and species identity can reveal more than whether vegetation is present; they can show which species are shaping habitat identity, whether native halophytes are establishing, and how non-native species contribute to community composition.

Salt marsh vegetation is shaped by strong environmental filters, including salinity, tidal flooding, saturated soils, elevation, and seasonal water availability. In southern California marshes, native salt-tolerant species such as pickleweed (*Salicornia pacifica*) and saltgrass (*Distichlis spicata*) often help define the structure and identity of coastal salt marsh habitat (Ferren 1985; North Campus Open Space 2025). At NCOS, monitoring reports evaluate restoration progress across multiple habitat types, including vegetation cover, native biodiversity, and relative native cover (Rickard 2024). Our project builds from that broader monitoring context by focusing specifically on salt marsh and brackish marsh vegetation.

Previous work on California wetland restoration shows that vegetation establishment can be slow, variable, and shaped by local conditions such as elevation, salinity, soil moisture, planting, disturbance, and adaptive management (Beheshti et al. 2023). For this reason, changes in vegetation cover should be interpreted as part of a restoration trajectory rather than as a single fixed measure of success. A restored marsh with high total vegetation cover may still differ ecologically depending on whether that cover is dominated by native salt marsh species or by non-native plants.

This study asks three main questions: Which dominant plant species characterize salt marsh and brackish marsh vegetation at NCOS? How has mean percent cover of dominant salt marsh species changed across survey years? How has native and non-native vegetation cover changed across survey years?

Methods

Study system and datasets

We used vegetation monitoring data from NCOS to examine plant community composition in salt marsh and brackish marsh habitat. The main dataset, `veg.csv`, contains vegetation survey observations, including plant species identity, percent cover, cover category, habitat type, transect, site, date, and year. The metadata file, `vp_veg_metadata.csv`, provides supporting information about vegetation transects, including the number of quadrats sampled within each transect-year.

The dataset covers vegetation surveys from 2021 through 2025 in the filtered salt marsh and brackish marsh habitat categories. To make survey years more comparable, we focused on August and September surveys, when late-summer and early-fall vegetation cover is expected to be relatively high. We filtered the dataset to include only native and non-native plant cover categories, excluding bare ground, thatch, and other non-plant cover types.

Data cleaning and summary approach

We summarized vegetation cover as mean percent cover by year, transect, species, and cover category. Because sampling effort differed among transects and years, we used the metadata file to account for the number of quadrats sampled in each transect-year. We also used `complete()` to make implicit absences explicit as zero percent cover, so that species not recorded in a transect-year were treated as absent rather than missing.

Dominant species were identified using the 90th percentile of mean percent cover across all plant species in the filtered dataset. Species at or above this threshold were classified as dominant because their average cover exceeded that of most species recorded in the salt marsh/brackish marsh data. We then summarized dominant species cover by survey year and visualized these patterns with a stacked bar plot.

To compare native and non-native vegetation, we summarized transect-level native and non-native cover by year and then calculated yearly means and standard errors. We also summarized total vegetation cover by year to provide broader context for changes in vegetation abundance through time. These analyses are descriptive and are intended to identify patterns in community composition and cover through time.

Results

Dominant species identity

Species differed strongly in mean percent cover across the filtered salt marsh and brackish marsh dataset. *Salicornia pacifica* had the highest mean cover overall, followed by *Distichlis spicata* and *Atriplex lentiformis*. Several non-native species, including *Raphanus sativus* and *Festuca perennis*, also appeared among the higher-cover species, showing that non-native plants contributed meaningfully to marsh vegetation composition even when native species dominated overall.




Figure 1: Mean percent cover (± 1 SE) of the top 20 plant species recorded in filtered salt marsh/brackish marsh habitat at NCOS. Species are ranked by mean cover across transects and survey years. Green indicates native species and coral indicates non-native species. The dashed line marks the 90th percentile cutoff used to identify dominant species.

Dominant species cover through time

Dominant species cover varied across survey years. The summed mean cover of dominant species was highest around 2023 and lower in 2025, suggesting that dominant vegetation cover may fluctuate across years rather than increasing steadily. Native salt marsh species, especially *Salicornia pacifica* and *Distichlis spicata*, were major contributors across the survey period. However, non-native dominant species also contributed to cover in some years, especially *Raphanus sativus* and *Festuca perennis*.




Figure 2: Mean percent cover of dominant salt marsh/brackish marsh plant species at NCOS across survey years. Dominant species were identified as species at or above the 90th percentile of mean percent cover across all survey years. Bar height reflects total mean cover of the dominant species and varies across years. Green shades indicate native species and coral/red shades indicate non-native species.

Native and non-native vegetation cover

Native vegetation cover remained higher than non-native vegetation cover across all survey years. Native cover increased from 2021 to 2022 and remained relatively high through 2024 before declining in 2025. Non-native cover was lower overall but increased from 2022 to

2024 before also declining in 2025. These patterns suggest that native vegetation was the stronger component of marsh cover, while non-native vegetation still contributed noticeably to community composition.




Figure 3: Mean percent cover (± 1 SE) of native and non-native vegetation in salt marsh/brackish marsh habitat at NCOS across survey years. Faint points show transect-level observations; lines connect yearly means; shaded bands represent ± 1 standard error. Native cover is shown in green and non-native cover in coral.

Total vegetation cover

Total vegetation cover, combining native and non-native plants, increased from 2021 to 2024 and then declined in 2025. This broader pattern provides context for the species-level and native/non-native patterns above: changes in relative composition occurred alongside changes in total cover, suggesting that interannual variability may influence both overall vegetation abundance and the species contributing to that cover.




Figure 4: Total mean percent vegetation cover (native + non-native combined) in salt marsh/brackish marsh habitat at NCOS across survey years. Faint points show transect-level observations; the line connects yearly means; the shaded band represents ± 1 standard error around the mean.

Discussion

References

- Beheshti, Kathryn M., Stephen C. Schroeter, Andres A. Deza, Daniel C. Reed, Rachel S. Smith, and Henry M. Page. 2023. "Large-Scale Field Studies Inform Adaptive Management of California Wetland Restoration." *Restoration Ecology* 31 (5): e13936. <https://doi.org/10.1111/rec.13936>.
- Cheadle Center for Biodiversity and Ecological Restoration, and Environmental Science Associates. 2016. *North Campus Open Space Restoration Project Restoration Plan*. University of California, Santa Barbara. <https://escholarship.org/uc/item/9505h60h>.
- Ferren, Wayne R. Jr. 1985. *Carpinteria Salt Marsh: Environment, History, and Botanical Resources of a Southern California Estuary*. Publication Number 4. The Herbarium, Department of Biological Sciences, University of California, Santa Barbara.
- North Campus Open Space. 2025. *Coastal Salt Marsh*. <https://www.ncos.ccber.ucsb.edu/native-plant-habitats/coastal-salt-marsh>.
- Rickard, Alison. 2024. *North Campus Open Space Restoration Project Monitoring Report: Year 7, December 2024*. University of California, Santa Barbara Cheadle Center for Biodiversity; Ecological Restoration. <https://escholarship.org/uc/item/7bq618m8>.