

Lecture 6

Eric Tovar

Qualifying Prep Course – Numerical

07-10-2020



1 January 2010 Problem 3

2 August 2009 Exam

- Problem 3
- Problem 2



Problem 3

Consider the following initial boundary value problem: find $u(x, t)$ such that

$$u_t - u_{xx} + u = 0, \quad 0 < x < 1, \quad t > 0 \quad (1.1)$$

$$u_x(0, t) = u_x(1, t) = 0, \quad t > 0 \quad (1.2)$$

$$u(x, 0) = g(x), \quad 0 < x < 1. \quad (1.3)$$

- (a) Derive the semi-discrete approximation of this problem using linear finite elements over a uniform partition of $(0, 1)$. Write it as a system of linear ordinary differential equations for the coefficient vector.
- (b) Further, derive discretizations in time using backward Euler and Crank-Nicolson methods, respectively.
- (c) Show that both fully discrete schemes are unconditionally stable with respect to the initial data in the spatial $L^2(0, 1)$ -norm.



Solution to (a)

Derive the semi-discrete approximation of this problem using linear finite elements over a uniform partition of $(0, 1)$. Write it as a system of linear ordinary differential equations for the coefficient vector.



Solution to (a)

Q: What does it mean to be semi-discrete?



Solution to (a)

Q: What does it mean to be semi-discrete?

Let us proceed “informally”.



Solution to (a)

Q: What does it mean to be semi-discrete?

Let us proceed “informally”. We first multiply the PDE in our IBVP by a sufficiently smooth test function $v(x)$ in some real vector space V , and integrate over the domain $(0, 1)$:

$$0 = \int_0^1 \left(u_t(x, t)v(x) - u_{xx}(x, t)v(x) + u(x, t)v(x) \right) dx,$$



Solution to (a)

Q: What does it mean to be semi-discrete?

Let us proceed “informally”. We first multiply the PDE in our IBVP by a sufficiently smooth test function $v(x)$ in some real vector space V , and integrate over the domain $(0, 1)$:

$$0 = \int_0^1 \left(u_t(x, t)v(x) - u_{xx}(x, t)v(x) + u(x, t)v(x) \right) dx,$$

$$\stackrel{\text{IBP}}{=} (u_t, v) + \int_0^1 \left(u_x(x, t)v_x(x) + u(x, t)v(x) \right) dx - \cancel{[u_x(x, t)v(x)]}_0^1$$



Solution to (a)

Q: What does it mean to be semi-discrete?

Let us proceed “informally”. We first multiply the PDE in our IBVP by a sufficiently smooth test function $v(x)$ in some real vector space V , and integrate over the domain $(0, 1)$:

$$\begin{aligned} 0 &= \int_0^1 \left(u_t(x, t)v(x) - u_{xx}(x, t)v(x) + u(x, t)v(x) \right) dx, \\ &\stackrel{\text{IBP}}{=} (u_t, v) + \int_0^1 \left(u_x(x, t)v_x(x) + u(x, t)v(x) \right) dx - \cancel{[u_x(x, t)v(x)]_0^1} \\ &= (u_t, v) + a(u, v), \end{aligned}$$

where we define the inner product (\cdot, \cdot) and bilinear form $a(\cdot, \cdot)$ to be:

$$\begin{aligned} (u_t, v) &= \int_0^1 u_t v \, dx \\ a(u, v) &= \int_0^1 (u_x v_x + uv) \, dx. \end{aligned}$$



For the integrals above, it is sufficient to take our function space to be $H^1(0, 1)$.



For the integrals above, it is sufficient to take our function space to be $H^1(0, 1)$. Now let V_h be the space of piecewise linear functions defined over the uniform partition of $(0, 1)$.



For the integrals above, it is sufficient to take our function space to be $H^1(0, 1)$. Now let V_h be the space of piecewise linear functions defined over the uniform partition of $(0, 1)$. More precisely,

$$V_h := \{v \in H^1(0, 1) : v|_{[x_i, x_{i+1}]} \in \mathbb{P}_1, i = 0, \dots, N-1\}.$$



For the integrals above, it is sufficient to take our function space to be $H^1(0, 1)$. Now let V_h be the space of piecewise linear functions defined over the uniform partition of $(0, 1)$. More precisely,

$$V_h := \{v \in H^1(0, 1) : v|_{[x_i, x_{i+1}]} \in \mathbb{P}_1, i = 0, \dots, N - 1\}.$$

Thus, our weak formulation in the finite element discretization becomes: Find $u_h(t) \in V_h$ such that $(u_{h,t}, v_h) + a(u_h, v_h) = 0$ for all $v_h \in V_h$. Note, $u_h(t) \in V_h$ for each $t > 0$.



Now, let us consider the standard tent functions as our basis for V_h : $\text{span}\{\phi_i\}_{i=0}^N$. We set $v = \phi_j(x)$ and express u_h in terms of the basis functions

$$u_h(x, t) = \sum_{i=0}^N u_i(t) \phi_i(x),$$

where $\{u_i\}_{i=0}^N$ are the unknown coefficients.



Now, let us consider the standard tent functions as our basis for V_h : $\text{span}\{\phi_i\}_{i=0}^N$. We set $v = \phi_j(x)$ and express u_h in terms of the basis functions

$$u_h(x, t) = \sum_{i=0}^N u_i(t) \phi_i(x),$$

where $\{u_i\}_{i=0}^N$ are the unknown coefficients. Substituting these into our weak formulation yields:



Now, let us consider the standard tent functions as our basis for V_h : $\text{span}\{\phi_i\}_{i=0}^N$. We set $v = \phi_j(x)$ and express u_h in terms of the basis functions

$$u_h(x, t) = \sum_{i=0}^N u_i(t) \phi_i(x),$$

where $\{u_i\}_{i=0}^N$ are the unknown coefficients. Substituting these into our weak formulation yields:

$$\int_0^1 \sum_{i=0}^N u_i'(t) \phi_i(x) \phi_j(x) dx + \int_0^1 \left(\sum_{i=0}^N u_i(t) \phi_i'(x) \phi_j'(x) + \sum_{i=0}^N u_i(t) \phi_i(x) \phi_j(x) \right) dx,$$

for $j = 0, \dots, N$.



Using the vector/matrix notation, $\mathbf{U}(t) = [u_i(t)]_{i=1}^N$, $\mathbf{M} = [(\phi_i, \phi_j)]_{i,j=1}^N$,
and $\mathbf{A} = [a(\phi_i, \phi_j)]_{i,j=1}^N$,



Using the vector/matrix notation, $\mathbf{U}(t) = [u_i(t)]_{i=1}^N$, $\mathbf{M} = [(\phi_i, \phi_j)]_{i,j=1}^N$, and $\mathbf{A} = [a(\phi_i, \phi_j)]_{i,j=1}^N$, we can rewrite our semi-discrete variational equation as,

$$\begin{cases} \mathbf{M}\mathbf{U}'(t) + \mathbf{A}\mathbf{U}(t) = \mathbf{0}, \\ \mathbf{U}(0) = \mathbf{G}, \end{cases}$$

where $\mathbf{G} = [g(x_i)]_{i=1}^N$.



Using the vector/matrix notation, $\mathbf{U}(t) = [u_i(t)]_{i=1}^N$, $\mathbf{M} = [(\phi_i, \phi_j)]_{i,j=1}^N$, and $\mathbf{A} = [a(\phi_i, \phi_j)]_{i,j=1}^N$, we can rewrite our semi-discrete variational equation as,

$$\begin{cases} \mathbf{M}\mathbf{U}'(t) + \mathbf{A}\mathbf{U}(t) = \mathbf{0}, \\ \mathbf{U}(0) = \mathbf{G}, \end{cases}$$

where $\mathbf{G} = [g(x_i)]_{i=1}^N$. This is our system of linear ordinary differential equations.



Solution to (b)

Further, derive discretizations in time using backward Euler and Crank-Nicolson methods, respectively.

We now want to provide the discretization in time using both the backward Euler and Crank-Nicolson methods.



Solution to (b)

We now want to provide the discretization in time using both the backward Euler and Crank-Nicolson methods. Let us define $t^n = n\Delta t$ for $n = 0, 1, \dots, M$ with $\Delta t = \frac{T}{M}$ where T is the final time.



Solution to (b)

We now want to provide the discretization in time using both the backward Euler and Crank-Nicolson methods. Let us define $t^n = n\Delta t$ for $n = 0, 1, \dots, M$ with $\Delta t = \frac{T}{M}$ where T is the final time. Then, let us consider the general discretization in time (θ -method):

$$M \frac{\mathbf{U}^{n+1} - \mathbf{U}^n}{\Delta t} + \mathbf{A}(\theta \mathbf{U}^{n+1} + (1 - \theta) \mathbf{U}^n) = \mathbf{0},$$

where $\theta \in [0, 1]$.



Solution to (b)

We now want to provide the discretization in time using both the backward Euler and Crank-Nicolson methods. Let us define $t^n = n\Delta t$ for $n = 0, 1, \dots, M$ with $\Delta t = \frac{T}{M}$ where T is the final time. Then, let us consider the general discretization in time (θ -method):

$$\mathbf{M} \frac{\mathbf{U}^{n+1} - \mathbf{U}^n}{\Delta t} + \mathbf{A}(\theta \mathbf{U}^{n+1} + (1 - \theta) \mathbf{U}^n) = \mathbf{0},$$

where $\theta \in [0, 1]$. The backward Euler scheme is obtained by using $\theta = 1$



Solution to (b)

We now want to provide the discretization in time using both the backward Euler and Crank-Nicolson methods. Let us define $t^n = n\Delta t$ for $n = 0, 1, \dots, M$ with $\Delta t = \frac{T}{M}$ where T is the final time. Then, let us consider the general discretization in time (θ -method):

$$M \frac{\mathbf{U}^{n+1} - \mathbf{U}^n}{\Delta t} + \mathbf{A}(\theta \mathbf{U}^{n+1} + (1 - \theta) \mathbf{U}^n) = \mathbf{0},$$

where $\theta \in [0, 1]$. The backward Euler scheme is obtained by using $\theta = 1$ and the Crank-Nicolson scheme by using $\theta = \frac{1}{2}$.



Solution to (c)

Show that both fully discrete schemes are unconditionally stable with respect to the initial data in the spatial $L^2(0, 1)$ -norm. Let us consider the following variational problem:

$$\left(\frac{u^{n+1} - u^n}{\Delta t}, v \right) + a(\theta u^{n+1} + (1 - \theta)u^n, v) = 0, \quad (1.4)$$

where $u^n := u(x, n\Delta t)$.



Solution to (c)

Let us consider the following variational problem:

$$\left(\frac{u^{n+1} - u^n}{\Delta t}, v \right) + a(\theta u^{n+1} + (1 - \theta)u^n, v) = 0, \quad (1.4)$$

where $u^n := u(x, n\Delta t)$. Let us first consider the backward Euler method, $\theta = 1$:



Solution to (c)

Let us consider the following variational problem:

$$\left(\frac{u^{n+1} - u^n}{\Delta t}, v \right) + a(\theta u^{n+1} + (1 - \theta)u^n, v) = 0, \quad (1.4)$$

where $u^n := u(x, n\Delta t)$. Let us first consider the backward Euler method, $\theta = 1$:

$$\left(\frac{u^{n+1} - u^n}{\Delta t}, v \right) + a(u^{n+1}, v) = 0,$$



Let us make the *ansatz* $v = u^{n+1}$.



Let us make the *ansatz* $v = u^{n+1}$. Substituting into the above weak formulation, we have that

$$\left(\frac{u^{n+1} - u^n}{\Delta t}, u^{n+1} \right) + a(u^{n+1}, u^{n+1}) = 0.$$



Let us make the *ansatz* $v = u^{n+1}$. Substituting into the above weak formulation, we have that

$$\left(\frac{u^{n+1} - u^n}{\Delta t}, u^{n+1} \right) + a(u^{n+1}, u^{n+1}) = 0.$$

Note that $a(u^{n+1}, u^{n+1}) \geq 0$,



Let us make the *ansatz* $v = u^{n+1}$. Substituting into the above weak formulation, we have that

$$\left(\frac{u^{n+1} - u^n}{\Delta t}, u^{n+1} \right) + a(u^{n+1}, u^{n+1}) = 0.$$

Note that $a(u^{n+1}, u^{n+1}) \geq 0$, Dropping this term gives us that $LHS \leq 0$.



Let us make the *ansatz* $v = u^{n+1}$. Substituting into the above weak formulation, we have that

$$\left(\frac{u^{n+1} - u^n}{\Delta t}, u^{n+1} \right) + a(u^{n+1}, u^{n+1}) = 0.$$

Note that $a(u^{n+1}, u^{n+1}) \geq 0$. Dropping this term gives us that $LHS \leq 0$. Expanding and rearranging yields the following inequality:

$$\|u^{n+1}\|_{L^2(0,1)}^2 = (u^{n+1}, u^{n+1}) \leq (u^n, u^{n+1}) \leq \|u^n\|_{L^2(0,1)} \|u^{n+1}\|_{L^2(0,1)}.$$

Then, dividing by $\|u^{n+1}\|_{L^2(0,1)}$, we have that:

$$\|u^{n+1}\|_{L^2(0,1)} \leq \|u^n\|_{L^2(0,1)} \leq \cdots \leq \|u^0\|_{L^2(0,1)} = \|g\|_{L^2(0,1)}.$$

Thus the backward Euler is unconditionally stable.



Let us now consider the Crank-Nicolson method, $\theta = \frac{1}{2}$:



Let us now consider the Crank-Nicolson method, $\theta = \frac{1}{2}$:

$$\left(\frac{u^{n+1} - u^n}{\Delta t}, v\right) + a\left(\frac{u^{n+1} + u^n}{2}, v\right) = 0.$$



Let us now consider the Crank-Nicolson method, $\theta = \frac{1}{2}$:

$$\left(\frac{u^{n+1} - u^n}{\Delta t}, v\right) + a\left(\frac{u^{n+1} + u^n}{2}, v\right) = 0.$$

We make the *ansatz* $v = \frac{u^{n+1} + u^n}{2}$ and proceed similarly as in the backward Euler case:

$$\left(\frac{u^{n+1} - u^n}{\Delta t}, \frac{u^{n+1} + u^n}{2}\right) \leq 0.$$



Let us now consider the Crank-Nicolson method, $\theta = \frac{1}{2}$:

$$\left(\frac{u^{n+1} - u^n}{\Delta t}, v\right) + a\left(\frac{u^{n+1} + u^n}{2}, v\right) = 0.$$

We make the *ansatz* $v = \frac{u^{n+1} + u^n}{2}$ and proceed similarly as in the backward Euler case:

$$\left(\frac{u^{n+1} - u^n}{\Delta t}, \frac{u^{n+1} + u^n}{2}\right) \leq 0.$$

Expanding the inner product yields:



Let us now consider the Crank-Nicolson method, $\theta = \frac{1}{2}$:

$$\left(\frac{u^{n+1} - u^n}{\Delta t}, v\right) + a\left(\frac{u^{n+1} + u^n}{2}, v\right) = 0.$$

We make the *ansatz* $v = \frac{u^{n+1} + u^n}{2}$ and proceed similarly as in the backward Euler case:

$$\left(\frac{u^{n+1} - u^n}{\Delta t}, \frac{u^{n+1} + u^n}{2}\right) \leq 0.$$

Expanding the inner product yields:

$$\|u^{n+1}\|_{L^2(0,1)}^2 \leq \|u^n\|_{L^2(0,1)}^2.$$

Thus, we have that

$$\|u^{n+1}\|_{L^2(0,1)} \leq \|g\|_{L^2(0,1)}.$$

and the Crank-Nicolson scheme is unconditionally stable.



1 January 2010 Problem 3

2 August 2009 Exam

- Problem 3
- Problem 2



Problem 3

Let $\Omega_e = \{x \in \mathbb{R}^2 : \|x\| > 1\}$. Show that the Poincare inequality does not hold in $H_0^1(\Omega)$, i.e., there does not exist a constant $c > 0$ satisfying

$$c\|u\|_{L^2(\Omega_e)}^2 \leq \int_{\Omega_e} \|\nabla u\|^2 dx, \quad \text{for all } u \in H_0^1(\Omega_e).$$

The space $H_0^1(\Omega_e)$ is the completion of $C_0^\infty(\Omega_e)$ in the norm:

$$\|u\|_{H^1(\Omega)} = \left(\|v\|_{L^2(\Omega_e)}^2 + \|\nabla v\|_{(L^2(\Omega_e))^2}^2 \right)^{1/2}$$

(Hint: Consider dilating a fixed function.)



Loosely speaking, the Poincare inequality defined above allows us to obtain bounds on our function u using bounds on its gradient.



Loosely speaking, the Poincare inequality defined above allows us to obtain bounds on our function u using bounds on its gradient. For the Poincare inequality to hold, both the norms $\|u\|_{L^2(\Omega_e)}^2$ and $\|\nabla v\|_{(L^2(\Omega_e))^2}^2$ need to make sense.



Loosely speaking, the Poincare inequality defined above allows us to obtain bounds on our function u using bounds on its gradient. For the Poincare inequality to hold, both the norms $\|u\|_{L^2(\Omega_e)}^2$ and $\|\nabla v\|_{(L^2(\Omega_e))^2}^2$ need to make sense. We will show that this is not the case.



Recall that our domain is the the unbounded annulus

$\Omega_e = \{x \in \mathbb{R}^2 : \|x\| > 1\}$ and so it will be convenient for us to work in polar coordinates.



Recall that our domain is the the unbounded annulus

$\Omega_e = \{x \in \mathbb{R}^2 : \|x\| > 1\}$ and so it will be convenient for us to work in polar coordinates. Let us define $r := \|x\|$ and consider the following bump function in polar coordinates:

$$\phi(r, \theta) := \phi(r) := \begin{cases} \exp(\frac{-1}{r(1-r)}) & \text{for } 0 < r < 1 \\ 0 & \text{for } r \geq 1, \end{cases}$$

which is only a function of r .



Recall that our domain is the the unbounded annulus $\Omega_e = \{x \in \mathbb{R}^2 : \|x\| > 1\}$ and so it will be convenient for us to work in polar coordinates. Let us define $r := \|x\|$ and consider the following bump function in polar coordinates:

$$\phi(r, \theta) := \phi(r) := \begin{cases} \exp(\frac{-1}{r(1-r)}) & \text{for } 0 < r < 1 \\ 0 & \text{for } r \geq 1, \end{cases}$$

which is only a function of r . Let us now consider the sequence of bump functions defined by,

$$\phi_n(r) := \phi(\frac{r-1}{n}).$$

Note that ϕ_n has the support $\text{supp}(\phi_n) \subset [1, n+1]$.



Now, let us consider the L^2 norm of ϕ_n :



Now, let us consider the L^2 norm of ϕ_n :

$$\|\phi_n\|_{L^2(\Omega_e)}^2 = \int_0^{2\pi} \int_1^\infty \phi_n^2(r) r \, dr \, d\theta$$



Now, let us consider the L^2 norm of ϕ_n :

$$\begin{aligned}\|\phi_n\|_{L^2(\Omega_e)}^2 &= \int_0^{2\pi} \int_1^\infty \phi_n^2(r) r \, dr \, d\theta \\ &= 2\pi \int_1^{n+1} \phi^2\left(\frac{r-1}{n}\right) r \, dr\end{aligned}$$



Now, let us consider the L^2 norm of ϕ_n :

$$\begin{aligned}\|\phi_n\|_{L^2(\Omega_e)}^2 &= \int_0^{2\pi} \int_1^\infty \phi_n^2(r) r \, dr \, d\theta \\ &= 2\pi \int_1^{n+1} \phi^2\left(\frac{r-1}{n}\right) r \, dr \\ &= 2\pi \int_0^1 \phi^2(s) n(ns+1) \, ds\end{aligned}$$



Now, let us consider the L^2 norm of ϕ_n :

$$\begin{aligned}\|\phi_n\|_{L^2(\Omega_e)}^2 &= \int_0^{2\pi} \int_1^\infty \phi_n^2(r) r \, dr \, d\theta \\ &= 2\pi \int_1^{n+1} \phi^2\left(\frac{r-1}{n}\right) r \, dr \\ &= 2\pi \int_0^1 \phi^2(s) n(ns+1) \, ds \\ &= 2\pi n \int_0^1 \phi^2(s) \, ds + 2\pi n^2 \int_0^1 s\phi^2(s) \, ds\end{aligned}$$



Now, let us consider the L^2 norm of ϕ_n :

$$\begin{aligned} \|\phi_n\|_{L^2(\Omega_e)}^2 &= \int_0^{2\pi} \int_1^\infty \phi_n^2(r) r \, dr \, d\theta \\ &= 2\pi \int_1^{n+1} \phi^2\left(\frac{r-1}{n}\right) r \, dr \\ &= 2\pi \int_0^1 \phi^2(s) n(ns+1) \, ds \\ &= 2\pi n \int_0^1 \phi^2(s) \, ds + 2\pi n^2 \int_0^1 s\phi^2(s) \, ds \\ &\geq 2\pi n^2 \int_0^1 s\phi^2(s) \, ds \end{aligned}$$



Now, let us consider the L^2 norm of ϕ_n :

$$\begin{aligned}\|\phi_n\|_{L^2(\Omega_e)}^2 &= \int_0^{2\pi} \int_1^\infty \phi_n^2(r) r \, dr \, d\theta \\ &= 2\pi \int_1^{n+1} \phi^2\left(\frac{r-1}{n}\right) r \, dr \\ &= 2\pi \int_0^1 \phi^2(s) n(ns+1) \, ds \\ &= 2\pi n \int_0^1 \phi^2(s) \, ds + 2\pi n^2 \int_0^1 s\phi^2(s) \, ds \\ &\geq 2\pi n^2 \int_0^1 s\phi^2(s) \, ds \\ &= C_0 n^2.\end{aligned}$$

Thus $\lim_{n \rightarrow \infty} \|\phi_n\|_{L^2(\Omega_e)} = \infty$.



Now, let us consider the H^1 -semi norm of ϕ_n :



Now, let us consider the H^1 -semi norm of ϕ_n :

$$|\phi_n|_{H^1(\Omega_e)}^2 = \int_0^{2\pi} \int_1^\infty |(\phi_n(r))'|^2 r \, dr \, d\theta$$



Now, let us consider the H^1 -semi norm of ϕ_n :

$$\begin{aligned} |\phi_n|_{H^1(\Omega_e)}^2 &= \int_0^{2\pi} \int_1^\infty |(\phi_n(r))'|^2 r \, dr \, d\theta \\ &= 2\pi \int_1^{n+1} \left| \frac{\partial}{\partial r} \phi\left(\frac{r-1}{n}\right) \right|^2 r \, dr \end{aligned}$$



Now, let us consider the H^1 -semi norm of ϕ_n :

$$\begin{aligned} |\phi_n|_{H^1(\Omega_e)}^2 &= \int_0^{2\pi} \int_1^\infty |(\phi_n(r))'|^2 r \, dr \, d\theta \\ &= 2\pi \int_1^{n+1} \left| \frac{\partial}{\partial r} \phi\left(\frac{r-1}{n}\right) \right|^2 r \, dr \\ &= 2\pi \int_0^1 \left| \frac{1}{n} \frac{\partial}{\partial s} \phi(s) \right|^2 n(ns+1) \, ds \\ &= 2\pi \int_0^1 \left| \frac{\partial}{\partial s} \phi(s) \right|^2 \frac{n(ns+1)}{n^2} \, ds \end{aligned}$$



Now, let us consider the H^1 -semi norm of ϕ_n :

$$\begin{aligned} |\phi_n|_{H^1(\Omega_e)}^2 &= \int_0^{2\pi} \int_1^\infty |(\phi_n(r))'|^2 r \, dr \, d\theta \\ &= 2\pi \int_1^{n+1} \left| \frac{\partial}{\partial r} \phi\left(\frac{r-1}{n}\right) \right|^2 r \, dr \\ &= 2\pi \int_0^1 \left| \frac{1}{n} \frac{\partial}{\partial s} \phi(s) \right|^2 n(ns+1) \, ds \\ &= 2\pi \int_0^1 \left| \frac{\partial}{\partial s} \phi(s) \right|^2 \frac{n(ns+1)}{n^2} \, ds \\ &\leq 2\pi \frac{n(n+1)}{n^2} \int_0^1 |\phi'(s)|^2 \, ds \end{aligned}$$



Now, let us consider the H^1 -semi norm of ϕ_n :

$$\begin{aligned} |\phi_n|_{H^1(\Omega_e)}^2 &= \int_0^{2\pi} \int_1^\infty |(\phi_n(r))'|^2 r \, dr \, d\theta \\ &= 2\pi \int_1^{n+1} \left| \frac{\partial}{\partial r} \phi\left(\frac{r-1}{n}\right) \right|^2 r \, dr \\ &= 2\pi \int_0^1 \left| \frac{1}{n} \frac{\partial}{\partial s} \phi(s) \right|^2 n(ns+1) \, ds \\ &= 2\pi \int_0^1 \left| \frac{\partial}{\partial s} \phi(s) \right|^2 \frac{n(ns+1)}{n^2} \, ds \\ &\leq 2\pi \frac{n(n+1)}{n^2} \int_0^1 |\phi'(s)|^2 \, ds \\ &\leq C_1 \end{aligned}$$



Now, let us consider the H^1 -semi norm of ϕ_n :

$$\begin{aligned} |\phi_n|_{H^1(\Omega_e)}^2 &= \int_0^{2\pi} \int_1^\infty |(\phi_n(r))'|^2 r \, dr \, d\theta \\ &= 2\pi \int_1^{n+1} \left| \frac{\partial}{\partial r} \phi\left(\frac{r-1}{n}\right) \right|^2 r \, dr \\ &= 2\pi \int_0^1 \left| \frac{1}{n} \frac{\partial}{\partial s} \phi(s) \right|^2 n(ns+1) \, ds \\ &= 2\pi \int_0^1 \left| \frac{\partial}{\partial s} \phi(s) \right|^2 \frac{n(ns+1)}{n^2} \, ds \\ &\leq 2\pi \frac{n(n+1)}{n^2} \int_0^1 |\phi'(s)|^2 \, ds \\ &\leq C_1 \end{aligned}$$

Thus since the H^1 -semi norm is bounded for all n and the L^2 norm blows up to infinity, this implies that we cannot have a Poincare inequality on this domain.



Problem 2

Consider the Neumann Problem:

$$-\Delta u = f \quad \text{in } \Omega \quad (2.1)$$

$$\frac{\partial u}{\partial n} = g \quad \text{on } \partial\Omega. \quad (2.2)$$

Here Ω is a bounded domain in \mathbb{R}^2 and f and g are suitably smooth.

- (a) Derive a weak form of the above problem using a test function in $H^1(\Omega)$.
- (b) Discuss when the weak form of Part a. has a solution and if it is unique.
- (c) Describe a variational formulation of (2.1) in terms of an appropriate Hilbert space V . Be sure to explicitly define V .
- (d) Prove coercivity of the form of Part a. on the V of Part c. when $\Omega = (0, 1)^2$.



Solution to (a)

We want to derive a weak formulation of the BVP.



Solution to (a)

We want to derive a weak formulation of the BVP. We begin by multiplying the PDE by a test function $v \in H^1(\Omega)$ and integrating over Ω :

$$\begin{aligned} - \int_{\Omega} \Delta u v \, dx &= \int_{\Omega} \nabla u \cdot \nabla v \, dx - \int_{\partial\Omega} \frac{\partial u}{\partial n} v \, ds \\ &= \int_{\Omega} \nabla u \cdot \nabla v \, dx - \int_{\partial\Omega} g v \, ds. \end{aligned}$$



Solution to (a)

We want to derive a weak formulation of the BVP. We begin by multiplying the PDE by a test function $v \in H^1(\Omega)$ and integrating over Ω :

$$\begin{aligned} - \int_{\Omega} \Delta u v \, dx &= \int_{\Omega} \nabla u \cdot \nabla v \, dx - \int_{\partial\Omega} \frac{\partial u}{\partial n} v \, ds \\ &= \int_{\Omega} \nabla u \cdot \nabla v \, dx - \int_{\partial\Omega} g v \, ds. \end{aligned}$$

So our weak formulation becomes: Find $u \in H^1(\Omega)$ such that $a(u, v) = L(v)$ for all $v \in H^1(\Omega)$ where

$$a(u, v) := \int_{\Omega} \nabla u \cdot \nabla v \, dx = \int_{\Omega} f v \, dx + \int_{\partial\Omega} g v \, ds =: L(v).$$



Solution to (b)

We first note (**HW: show this**) that a and F can both be shown to be continuous (for F you need to invoke the Trace Lemma 3.3 in Grossmann book).



Solution to (b)

We first note (**HW: show this**) that a and F can both be shown to be continuous (for F you need to invoke the Trace Lemma 3.3 in Grossmann book).

Then, we know from the Lax-Milgram lemma that a solution exists and is unique if a is coercive.



Solution to (b)

We first note (**HW: show this**) that a and F can both be shown to be continuous (for F you need to invoke the Trace Lemma 3.3 in Grossmann book).

Then, we know from the Lax-Milgram lemma that a solution exists and is unique if a is coercive. However, in our case coercivity will not hold since we will not have a Poincare inequality that will help us bound $a(u, u)$ below. (**Why?**)



Solution to (b)

We first note (**HW: show this**) that a and F can both be shown to be continuous (for F you need to invoke the Trace Lemma 3.3 in Grossmann book).

Then, we know from the Lax-Milgram lemma that a solution exists and is unique if a is coercive. However, in our case coercivity will not hold since we will not have a Poincare inequality that will help us bound $a(u, u)$ below. (**Why?**)

Consider $v = \text{constant} \in H^1(\Omega)$. Specifically, let $v = 1$ and substitute into the weak formulation



Solution to (b)

We first note (**HW: show this**) that a and F can both be shown to be continuous (for F you need to invoke the Trace Lemma 3.3 in Grossmann book).

Then, we know from the Lax-Milgram lemma that a solution exists and is unique if a is coercive. However, in our case coercivity will not hold since we will not have a Poincare inequality that will help us bound $a(u, u)$ below. (**Why?**)

Consider $v = \text{constant} \in H^1(\Omega)$. Specifically, let $v = 1$ and substitute into the weak formulation

$$\int_{\Omega} f \, dx + \int_{\partial\Omega} g \, ds = 0.$$



Solution to (b)

We first note (**HW: show this**) that a and F can both be shown to be continuous (for F you need to invoke the Trace Lemma 3.3 in Grossmann book).

Then, we know from the Lax-Milgram lemma that a solution exists and is unique if a is coercive. However, in our case coercivity will not hold since we will not have a Poincare inequality that will help us bound $a(u, u)$ below. (**Why?**)

Consider $v = \text{constant} \in H^1(\Omega)$. Specifically, let $v = 1$ and substitute into the weak formulation

$$\int_{\Omega} f \, dx + \int_{\partial\Omega} g \, ds = 0.$$

This is called the **compatibility condition** or **solvability condition** and is necessary for existence of the solution. (**See Remark 3.27 on Neumann boundary conditions in Grossmann book**)



Now, note that if u is a solution then $u + c$ must also be a solution for $c \in \mathbb{R}$.



Now, note that if u is a solution then $u + c$ must also be a solution for $c \in \mathbb{R}$.

In order to guarantee a *unique* solution, we can impose the following condition on u :

$$\int_{\Omega} u \, dx = 0.$$



Now, note that if u is a solution then $u + c$ must also be a solution for $c \in \mathbb{R}$.

In order to guarantee a *unique* solution, we can impose the following condition on u :

$$\int_{\Omega} u \, dx = 0.$$

This assumption will allow us to derive a Poincaré inequality, which will then allow us to apply the Lax-Milgram lemma.



Solution to (c)

Describe a variational formulation of (2.1) in terms of an appropriate Hilbert space V . Be sure to explicitly define V . The variational formulation will be as follows: Find $u \in V := \{v \in H^1(\Omega) : \int_{\Omega} v \, ds = 0\}$ such that $a(u, v) = L(v)$ for all $v \in V$.



Solution to (c)

The variational formulation will be as follows: Find $u \in V := \{v \in H^1(\Omega) : \int_{\Omega} v \, ds = 0\}$ such that $a(u, v) = L(v)$ for all $v \in V$.



Solution to (d)

We want to show coercivity of the bilinear form $a(\cdot, \cdot)$. Recall that $a(\cdot, \cdot)$ is coercive when there exists a $\gamma > 0$ such that $\gamma \|u\|_V^2 \leq a(u, u)$ for all $u \in V$.



Solution to (d)

We want to show coercivity of the bilinear form $a(\cdot, \cdot)$. Recall that $a(\cdot, \cdot)$ is coercive when there exists a $\gamma > 0$ such that $\gamma \|u\|_V^2 \leq a(u, u)$ for all $u \in V$.

Note that our domain $\Omega = [0, 1]^2$ is a Lipschitz domain.



Solution to (d)

We want to show coercivity of the bilinear form $a(\cdot, \cdot)$. Recall that $a(\cdot, \cdot)$ is coercive when there exists a $\gamma > 0$ such that $\gamma \|u\|_V^2 \leq a(u, u)$ for all $u \in V$.

Note that our domain $\Omega = [0, 1]^2$ is a Lipschitz domain. Let us recall the Poincaré–Steklov Lemma:

Lemma 3.24 (Poincaré–Steklov). *Let D be a Lipschitz domain in \mathbb{R}^d . Let $\ell_D := \text{diam}(D)$. Let $p \in [1, \infty]$. There is $C_{\text{PS},p}$ (the subscript p is omitted when $p = 2$) s.t.*

$$C_{\text{PS},p} \|v - \underline{v}_D\|_{L^p(D)} \leq \ell_D |v|_{W^{1,p}(D)}, \quad \forall v \in W^{1,p}(D), \quad (3.8)$$

where $\underline{v}_D := \frac{1}{|D|} \int_D v \, dx$. The following holds true when D is convex:

$$C_{\text{PS},1} = 2, \quad C_{\text{PS}} := C_{\text{PS},2} = \pi, \quad C_{\text{PS},p} \geq \frac{1}{2} \left(\frac{2}{p} \right)^{\frac{1}{p}}, \quad p > 1. \quad (3.9)$$



Since we have that $\int_{\Omega} v \, ds = 0$, coercivity follows in the usual way with an application of the Poincare-Steklov Lemma.

