

# Lecture 2

Eric Tovar

Qualifying Prep Course – Numerical

07-02-2020



- 1 January 2009 Exam
  - Problem 2



# Problem 2

## Problem 2.

Let  $\Omega = (0, 1)^2$  and  $u$  be the solution of the second order elliptic problem:

$$-\Delta u : -u_{x_1 x_1} - u_{x_2 x_2} = f(x), \quad \text{for } x \in \Omega \quad (1.1)$$

$$\frac{\partial u}{\partial n} + u = g(x), \quad \text{for } x \in \partial\Omega \quad (1.2)$$

where  $n$  is the outward normal unit vector to the boundary  $\partial\Omega$  and  $f(x)$  and  $g(x)$  are given functions.



# Problem 2

## Problem 2.

Let  $\Omega = (0, 1)^2$  and  $u$  be the solution of the second order elliptic problem:

$$-\Delta u : -u_{x_1x_1} - u_{x_2x_2} = f(x), \quad \text{for } x \in \Omega \quad (1.1)$$

$$\frac{\partial u}{\partial n} + u = g(x), \quad \text{for } x \in \partial\Omega \quad (1.2)$$

where  $n$  is the outward normal unit vector to the boundary  $\partial\Omega$  and  $f(x)$  and  $g(x)$  are given functions.

**Question 1:** What kind of boundary condition do we have?



# Problem 2

## Problem 2.

Let  $\Omega = (0, 1)^2$  and  $u$  be the solution of the second order elliptic problem:

$$-\Delta u : -u_{x_1x_1} - u_{x_2x_2} = f(x), \quad \text{for } x \in \Omega \quad (1.1)$$

$$\frac{\partial u}{\partial n} + u = g(x), \quad \text{for } x \in \partial\Omega \quad (1.2)$$

where  $n$  is the outward normal unit vector to the boundary  $\partial\Omega$  and  $f(x)$  and  $g(x)$  are given functions.

**Question 1:** What kind of boundary condition do we have?

**Remark 1:**  $\frac{\partial u}{\partial n} = \nabla u \cdot n$



- (a) Derive the weak formulation of this problem in the form  $a(u, v) = F(v)$ , where  $a(u, v)$  and  $F(v)$  are the appropriate bilinear and linear forms defined on the Sobolev space  $H^1(\Omega)$ .



- (a) Derive the weak formulation of this problem in the form  $a(u, v) = F(v)$ , where  $a(u, v)$  and  $F(v)$  are the appropriate bilinear and linear forms defined on the Sobolev space  $H^1(\Omega)$ .
- (b) Let  $S_h$  be a finite element space of continuous piece-wise polynomial functions defined over a regular partitioning of  $\Omega$  into triangles and let  $a_h(u, v)$  and  $F_h(v)$  be the bilinear forms where all integrals are computed approximately. Derive Strang's lemma for the error of the FEM: find  $u_h \in S_h$  such that  $a_h(u_h, v) = F_h(v)$ ,  $\forall v \in S_h$ .



- (a) Derive the weak formulation of this problem in the form  $a(u, v) = F(v)$ , where  $a(u, v)$  and  $F(v)$  are the appropriate bilinear and linear forms defined on the Sobolev space  $H^1(\Omega)$ .
- (b) Let  $S_h$  be a finite element space of continuous piece-wise polynomial functions defined over a regular partitioning of  $\Omega$  into triangles and let  $a_h(u, v)$  and  $F_h(v)$  be the bilinear forms where all integrals are computed approximately. Derive Strang's lemma for the error of the FEM: find  $u_h \in S_h$  such that  $a_h(u_h, v) = F_h(v)$ ,  $\forall v \in S_h$ .
- (c) Let  $S_h$  be the finite element space of piece-wise linear functions. Let all integrals in  $a(u, v)$  and  $F(v)$  be computed using quadratures. Namely, for  $\tau$  and  $e$  being triangle and edge defined by the vertexes  $P_1, P_2, P_3$  and  $P_1, P_2$  respectively,

$$\int_{\tau} w(x) dx \approx \frac{|\tau|}{3} (w(P_1) + w(P_2) + w(P_3)), \quad \int_e w(x) ds \approx \frac{|e|}{2} (w(\alpha) + w(\beta)) \quad (1.3)$$

where  $|\tau|$  is the area of  $\tau$  and  $|e|$  is the length of  $e$ , and  $\alpha$  and  $\beta$  are the Gaussian quadrature nodes. Explain why  $a(w, v) = a_h(w, v)$  for all  $w, v \in S_h$ .





## Solution to (a)

Let  $v \in V$  be sufficiently smooth. We proceed “formally”: We multiply (1.1) by  $v$  and integrate over the domain:



## Solution to (a)

Let  $v \in V$  be sufficiently smooth. We proceed “formally”: We multiply (1.1) by  $v$  and integrate over the domain:

**LHS:**

$$\begin{aligned} - \int_{\Omega} (\Delta u) v dx &\stackrel{\text{IBP}}{=} - \int_{\partial\Omega} v (\mathbf{n} \cdot \nabla u) ds + \int_{\Omega} \nabla u \cdot \nabla v dx \\ &\stackrel{(1.2)}{=} - \int_{\partial\Omega} v (g - u) ds + \int_{\Omega} \nabla u \cdot \nabla v dx \end{aligned}$$



## Solution to (a)

Let  $v \in V$  be sufficiently smooth. We proceed “formally”: We multiply (1.1) by  $v$  and integrate over the domain:

**LHS:**

$$\begin{aligned} - \int_{\Omega} (\Delta u) v dx &\stackrel{\text{IBP}}{=} - \int_{\partial\Omega} v (\mathbf{n} \cdot \nabla u) ds + \int_{\Omega} \nabla u \cdot \nabla v dx \\ &\stackrel{(1.2)}{=} - \int_{\partial\Omega} v (g - u) ds + \int_{\Omega} \nabla u \cdot \nabla v dx \\ &= \int_{\Omega} \nabla u \cdot \nabla v dx + \int_{\partial\Omega} u v ds - \int_{\partial\Omega} g v ds \end{aligned}$$



## Solution to (a)

Let  $v \in V$  be sufficiently smooth. We proceed “formally”: We multiply (1.1) by  $v$  and integrate over the domain:

**LHS:**

$$\begin{aligned} - \int_{\Omega} (\Delta u) v dx &\stackrel{\text{IBP}}{=} - \int_{\partial\Omega} v (\mathbf{n} \cdot \nabla u) ds + \int_{\Omega} \nabla u \cdot \nabla v dx \\ &\stackrel{(1.2)}{=} - \int_{\partial\Omega} v (g - u) ds + \int_{\Omega} \nabla u \cdot \nabla v dx \\ &= \int_{\Omega} \nabla u \cdot \nabla v dx + \int_{\partial\Omega} u v ds - \int_{\partial\Omega} g v ds \end{aligned}$$

**RHS:**

$$\int_{\Omega} f(x) v dx = \int_{\Omega} f(x) v dx$$



## Solution to (a)

Let  $v \in V$  be sufficiently smooth. We proceed “formally”: We multiply (1.1) by  $v$  and integrate over the domain:

**LHS:**

$$\begin{aligned} - \int_{\Omega} (\Delta u) v dx &\stackrel{\text{IBP}}{=} - \int_{\partial\Omega} v (\mathbf{n} \cdot \nabla u) ds + \int_{\Omega} \nabla u \cdot \nabla v dx \\ &\stackrel{(1.2)}{=} - \int_{\partial\Omega} v (g - u) ds + \int_{\Omega} \nabla u \cdot \nabla v dx \\ &= \int_{\Omega} \nabla u \cdot \nabla v dx + \int_{\partial\Omega} u v ds - \int_{\partial\Omega} g v ds \end{aligned}$$

**RHS:**

$$\int_{\Omega} f(x) v dx = \int_{\Omega} f(x) v dx$$

**Q: How do we group these terms?**



The weak formulation is given as follows:

Find  $u \in V := H^1(\Omega)$  such that for any  $v \in V$

$$a(u, v) = F(v)$$

where

$$a(u, v) := \int_{\Omega} \nabla u \cdot \nabla v dx + \int_{\partial\Omega} uv ds \quad (1.4)$$

$$F(v) := \int_{\Omega} f(x)v dx + \int_{\partial\Omega} gv ds \quad (1.5)$$



## (First) Solution to (b)

Let  $S_h$  be a finite element space of continuous piece-wise polynomial functions defined over a regular partitioning of  $\Omega$  into triangles and let  $a_h(u, v)$  and  $F_h(v)(!!!)$  be the bilinear forms where all **integrals are computed approximately**. Derive Strang's lemma for the error of the FEM: find  $u_h \in S_h$  such that  $a_h(u_h, v) = F_h(v)$ ,  $\forall v \in S_h$ .



---

<sup>1</sup>Reference: 4.5.3 in Grossman et al

## (First) Solution to (b)

Note that the “statement” of the problem is ambiguous. That is, what does it mean to *derive* a known result?



---

<sup>1</sup>Reference: 4.5.3 in Grossman et al



## (First) Solution to (b)

Note that the “statement” of the problem is ambiguous. That is, what does it mean to *derive* a known result? So let us consider the first solution. Recall the statement to Strang’s First Lemma<sup>1</sup>;

---

<sup>1</sup>Reference: 4.5.3 in Grossman et al



## (First) Solution to (b)

Note that the “statement” of the problem is ambiguous. That is, what does it mean to *derive* a known result? So let us consider the first solution. Recall the statement to Strang’s First Lemma<sup>1</sup>;

### Strang’s First Lemma

Let  $V_h \subset V$  and let the bilinear form  $a_h(\cdot, \cdot)$  be uniformly  $V_h$  – *elliptic*. Then, there exists a constant  $c > 0$  such that

$$\|u - u_h\| \leq c \left[ \inf_{z_h \in V_h} \{\|u - z_h\| + \|a(z_h, \cdot) - a_h(z_h, \cdot)\|_{*,h}\} + \|f - f_h\|_{*,h} \right].$$

---

<sup>1</sup>Reference: 4.5.3 in Grossman et al



**Proof.**

(i) First recall the Galerkin orthogonality property:

$$a_h(u - u_h, v_h) = 0, \quad \text{for all } v_h \in S_h.$$

This holds since  $S_h \subset H^1(\Omega)$ .



## Proof.

(i) First recall the Galerkin orthogonality property:

$$a_h(u - u_h, v_h) = 0, \quad \text{for all } v_h \in S_h.$$

This holds since  $S_h \subset H^1(\Omega)$ .

(ii) Let  $z_h, v_h \in S_h$ . Then we see that

$$a_h(u_h - z_h, v_h) = a_h(u_h, v_h) - a_h(z_h, v_h)$$



## Proof.

(i) First recall the Galerkin orthogonality property:

$$a_h(u - u_h, v_h) = 0, \quad \text{for all } v_h \in S_h.$$

This holds since  $S_h \subset H^1(\Omega)$ .

(ii) Let  $z_h, v_h \in S_h$ . Then we see that

$$\begin{aligned} a_h(u_h - z_h, v_h) &= a_h(u_h, v_h) - a_h(z_h, v_h) \\ &= F_h(v_h) - a_h(z_h, v_h) \end{aligned}$$



## Proof.

(i) First recall the Galerkin orthogonality property:

$$a_h(u - u_h, v_h) = 0, \quad \text{for all } v_h \in S_h.$$

This holds since  $S_h \subset H^1(\Omega)$ .

(ii) Let  $z_h, v_h \in S_h$ . Then we see that

$$\begin{aligned} a_h(u_h - z_h, v_h) &= a_h(u_h, v_h) - a_h(z_h, v_h) \\ &= F_h(v_h) - a_h(z_h, v_h) \\ &= F_h(v_h) - a_h(z_h, v_h) + \underbrace{a(u, v_h) - F(v_h)}_{=0} \\ &\quad + \underbrace{a(z_h, v_h) - a_h(z_h, v_h)}_{=0} \end{aligned}$$



## Proof.

(i) First recall the Galerkin orthogonality property:

$$a_h(u - u_h, v_h) = 0, \quad \text{for all } v_h \in S_h.$$

This holds since  $S_h \subset H^1(\Omega)$ .

(ii) Let  $z_h, v_h \in S_h$ . Then we see that

$$\begin{aligned} a_h(u_h - z_h, v_h) &= a_h(u_h, v_h) - a_h(z_h, v_h) \\ &= F_h(v_h) - a_h(z_h, v_h) \\ &= F_h(v_h) - a_h(z_h, v_h) + \underbrace{a(u, v_h) - F(v_h)}_{=0} \\ &\quad + \underbrace{a(z_h, v_h) - a_h(z_h, v_h)}_{=0} \\ &= a(u - z_h, v_h) + a(z_h, v_h) - a_h(z_h, v_h) + \\ &\quad F_h(v_h) - F(v_h) \end{aligned}$$



(iii) Now let  $v_h = u_h - z_h$ . Then we see that

$$a_h(u_h - z_h, u_h - z_h) = a(u - z_h, u_h - z_h) + a(z_h, u_h - z_h)$$





(iii) Now let  $v_h = u_h - z_h$ . Then we see that

$$\begin{aligned} a_h(u_h - z_h, u_h - z_h) &= a(u - z_h, u_h - z_h) + a(z_h, u_h - z_h) \\ &\quad - a_h(z_h, u_h - z_h) + F_h(v_h) - F(v_h) \end{aligned}$$



(iii) Now let  $v_h = u_h - z_h$ . Then we see that

$$\begin{aligned} a_h(u_h - z_h, u_h - z_h) &= a(u - z_h, u_h - z_h) + a(z_h, u_h - z_h) \\ &\quad - a_h(z_h, u_h - z_h) + F_h(v_h) - F(v_h) \\ &\implies \\ \alpha \|u_h - z_h\|^2 &\leq \|u - z_h\| \|u_h - z_h\| + \end{aligned}$$



(iii) Now let  $v_h = u_h - z_h$ . Then we see that

$$\begin{aligned} a_h(u_h - z_h, u_h - z_h) &= a(u - z_h, u_h - z_h) + a(z_h, u_h - z_h) \\ &\quad - a_h(z_h, u_h - z_h) + F_h(v_h) - F(v_h) \\ &\implies \\ \alpha \|u_h - z_h\|^2 &\leq \|u - z_h\| \|u_h - z_h\| + \\ &\quad |a(z_h, v_h) - a_h(z_h, v_h)| + |F_h(v_h) - F(v_h)| \end{aligned}$$

where we invoked  $V$ -ellipticity and continuity of  $a(\cdot, \cdot)$ .



(iii) Now let  $v_h = u_h - z_h$ . Then we see that

$$\begin{aligned} a_h(u_h - z_h, u_h - z_h) &= a(u - z_h, u_h - z_h) + a(z_h, u_h - z_h) \\ &\quad - a_h(z_h, u_h - z_h) + F_h(v_h) - F(v_h) \\ &\implies \\ \alpha \|u_h - z_h\|^2 &\leq \|u - z_h\| \|u_h - z_h\| + \\ &\quad |a(z_h, v_h) - a_h(z_h, v_h)| + |F_h(v_h) - F(v_h)| \end{aligned}$$

where we invoked  $V$ -ellipticity and continuity of  $a(\cdot, \cdot)$ .

**Recall:**  $\|w\|_{*,h} := \sup_{v_h \in V_h} \frac{|w(v_h)|}{\|v_h\|_h}$  for  $w \in V_h^*$  is the norm in the dual space  $V_h^*$ .



(iv) Now, we divide by  $\alpha\|u_h - z_h\|$  and take the supremum over  $v_h \in V_h$ :

$$\|u_h - z_h\| \leq \frac{C}{\alpha} \|u - z_h\| +$$



(iv) Now, we divide by  $\alpha\|u_h - z_h\|$  and take the supremum over  $v_h \in V_h$ :

$$\begin{aligned}\|u_h - z_h\| &\leq \frac{C}{\alpha}\|u - z_h\| + \\ &\quad \frac{1}{\alpha}\left(\|a(z_h, v_h) - a_h(z_h, v_h)\|_{*,h} + \|F_h(v_h) - F(v_h)\|_{*,h}\right)\end{aligned}$$



(iv) Now, we divide by  $\alpha\|u_h - z_h\|$  and take the supremum over  $v_h \in V_h$ :

$$\begin{aligned}\|u_h - z_h\| &\leq \frac{C}{\alpha}\|u - z_h\| + \\ &\quad \frac{1}{\alpha}\left(\|a(z_h, v_h) - a_h(z_h, v_h)\|_{*,h} + \|F_h(v_h) - F(v_h)\|_{*,h}\right)\end{aligned}$$

(v) Recall by the triangle inequality

$$\|u - u_h\| = \|u - z_h + z_h - u_h\| \leq \|u - z_h\| + \|u_h - z_h\|$$



(iv) Now, we divide by  $\alpha\|u_h - z_h\|$  and take the supremum over  $v_h \in V_h$ :

$$\begin{aligned}\|u_h - z_h\| &\leq \frac{C}{\alpha}\|u - z_h\| + \\ &\quad \frac{1}{\alpha}\left(\|a(z_h, v_h) - a_h(z_h, v_h)\|_{*,h} + \|F_h(v_h) - F(v_h)\|_{*,h}\right)\end{aligned}$$

(v) Recall by the triangle inequality

$$\|u - u_h\| = \|u - z_h + z_h - u_h\| \leq \|u - z_h\| + \|u_h - z_h\|$$

(vi) Thus, we have that

$$\|u - u_h\| \leq \|u - z_h\| + \underbrace{\|u_h - z_h\|}$$





(iv) Now, we divide by  $\alpha\|u_h - z_h\|$  and take the supremum over  $v_h \in V_h$ :

$$\begin{aligned}\|u_h - z_h\| &\leq \frac{C}{\alpha}\|u - z_h\| + \\ &\quad \frac{1}{\alpha}\left(\|a(z_h, v_h) - a_h(z_h, v_h)\|_{*,h} + \|F_h(v_h) - F(v_h)\|_{*,h}\right)\end{aligned}$$

(v) Recall by the triangle inequality

$$\|u - u_h\| = \|u - z_h + z_h - u_h\| \leq \|u - z_h\| + \|u_h - z_h\|$$

(vi) Thus, we have that

$$\begin{aligned}\|u - u_h\| &\leq \|u - z_h\| + \underbrace{\|u_h - z_h\|}_{\leq \frac{C}{\alpha}\|u - z_h\| + \frac{1}{\alpha}(\|a(z_h, v_h) - a_h(z_h, v_h)\|_{*,h} + \|F_h(v_h) - F(v_h)\|_{*,h})} \\ &\leq \left(1 + \frac{C}{\alpha}\right)\|u - z_h\| + \\ &\quad \frac{1}{\alpha}\left(\|a(z_h, v_h) - a_h(z_h, v_h)\|_{*,h} + \|F_h(v_h) - F(v_h)\|_{*,h}\right)\end{aligned}$$



(iv) Now, we divide by  $\alpha \|u_h - z_h\|$  and take the supremum over  $v_h \in V_h$ :

$$\begin{aligned} \|u_h - z_h\| &\leq \frac{C}{\alpha} \|u - z_h\| + \\ &\quad \frac{1}{\alpha} \left( \|a(z_h, v_h) - a_h(z_h, v_h)\|_{*,h} + \|F_h(v_h) - F(v_h)\|_{*,h} \right) \end{aligned}$$

(v) Recall by the triangle inequality

$$\|u - u_h\| = \|u - z_h + z_h - u_h\| \leq \|u - z_h\| + \|u_h - z_h\|$$

(vi) Thus, we have that

$$\begin{aligned} \|u - u_h\| &\leq \|u - z_h\| + \underbrace{\|u_h - z_h\|}_{\leq \frac{C}{\alpha} \|u - z_h\| + \frac{1}{\alpha} (\|a(z_h, v_h) - a_h(z_h, v_h)\|_{*,h} + \|F_h(v_h) - F(v_h)\|_{*,h})} \\ &\leq \left(1 + \frac{C}{\alpha}\right) \|u - z_h\| + \\ &\quad \frac{1}{\alpha} \left( \|a(z_h, v_h) - a_h(z_h, v_h)\|_{*,h} + \|F_h(v_h) - F(v_h)\|_{*,h} \right) \end{aligned}$$



(vii) Then take infimum over  $z_h$  to get final result.

## Alternative solution to (b)

An alternative way to approach this problem is to assume that it is asking to derive a generic error estimate.



## Alternative solution to (b) (technical) version

We say that the problem<sup>2</sup>

Find  $u_h \in v_h \in S_h$  such that

$$a_h(u_h, v_h) = F_h(v_h), \quad \text{for all } v_h \in S_h$$

is **stable** (or well-posed) whenever

$$\inf_{v_h \in S_h} \sup_{w_h \in S_h} \frac{|a_h(v_h, w_h)|}{\|v_h\|_{H^1(\Omega)} \|w_h\|_{H^1(\Omega)}} =: \alpha_h > 0 \quad (1.6)$$

---

<sup>2</sup>See discrete inf-sup condition on Page 91

# Alternative solution to (b) (technical) version

We say that the problem<sup>2</sup>

Find  $u_h \in v_h \in S_h$  such that

$$a_h(u_h, v_h) = F_h(v_h), \quad \text{for all } v_h \in S_h$$

is **stable** (or well-posed) whenever

$$\inf_{v_h \in S_h} \sup_{w_h \in S_h} \frac{|a_h(v_h, w_h)|}{\|v_h\|_{H^1(\Omega)} \|w_h\|_{H^1(\Omega)}} =: \alpha_h > 0 \quad (1.6)$$

(i) Assume the problem is stable. Then, an immediate consequence of (1.6) is: For every  $v_h \in S_h$

$$\sup_{w_h \in S_h} \frac{|a_h(v_h, w_h)|}{\|v_h\|_{H^1(\Omega)} \|w_h\|_{H^1(\Omega)}} \geq \alpha_h \implies \sup_{w_h \in S_h} \frac{|a_h(v_h, w_h)|}{\|w_h\|_{H^1(\Omega)}} \geq \alpha_h \|v_h\|_{H^1(\Omega)}.$$

---

<sup>2</sup>See discrete inf-sup condition on Page 91

(ii) Since  $u_h - v_h \in S_h$ , we have that:

$$\alpha_h \|u_h - v_h\|_{H^1(\Omega)} \leq \sup_{w_h \in S_h} \frac{|a_h(u_h - v_h, w_h)|}{\|w_h\|_{H^1(\Omega)}} \quad (1.7)$$

Note that we call this a **stability** condition.



(ii) Since  $u_h - v_h \in S_h$ , we have that:

$$\alpha_h \|u_h - v_h\|_{H^1(\Omega)} \leq \sup_{w_h \in S_h} \frac{|a_h(u_h - v_h, w_h)|}{\|w_h\|_{H^1(\Omega)}} \quad (1.7)$$

Note that we call this a **stability** condition.

(iii) Using that  $a_h(u_h - v_h, w_h) = F_h(w_h) - a_h(v_h, w_h)$ , we get the following **consistency** term:

$$\alpha_h \|u_h - v_h\|_{H^1(\Omega)} \leq \sup_{w_h \in S_h} \frac{|F_h(w_h) - a_h(v_h, w_h)|}{\|w_h\|_{H^1(\Omega)}} \quad (1.8)$$



(iv) **Triangle inequality.** The triangle inequality implies

$\|u - u_h\|_{H^1(\Omega)} \leq \|u - v_h\|_{H^1(\Omega)} + \|u_h - v_h\|_{H^1(\Omega)}$ . Thus using this and combining the above, we have that:

$$\begin{aligned} \|u - u_h\|_{H^1(\Omega)} &\leq \|u - v_h\|_{H^1(\Omega)} + \frac{1}{\alpha_h} \sup_{w_h \in S_h} \frac{|F_h(w_h) - a_h(v_h, w_h)|}{\|w_h\|_{H^1(\Omega)}} \\ &\leq \inf_{v_h \in S_h} \left( \|u - v_h\|_{H^1(\Omega)} \right. \\ &\quad \left. + \frac{1}{\alpha_h} \sup_{w_h \in S_h} \frac{|F_h(w_h) - a_h(v_h, w_h)|}{\|w_h\|_{H^1(\Omega)}} \right) \end{aligned}$$





## Solution to (c)

Let  $S_h$  be the finite element space of piece-wise **linear** functions. Let all integrals in  $a(u, v)$  and  $F(v)$  be computed using quadratures. Namely, for  $\tau$  and  $e$  being triangle and edge defined by the vertexes  $P_1, P_2, P_3$  and  $P_1, P_2$  respectively,

$$\int_{\tau} w(x) dx \approx \frac{|\tau|}{3} (w(P_1) + w(P_2) + w(P_3)), \quad \int_e w(x) ds \approx \frac{|e|}{2} (w(\alpha) + w(\beta)) \quad (1.9)$$

where  $|\tau|$  is the area of  $\tau$  and  $|e|$  is the length of  $e$ , and  $\alpha$  and  $\beta$  are the Gaussian quadrature nodes. Explain why  $a(w, v) = a_h(w, v)$  for all  $w, v \in S_h$ .



## Solution to (c)

Recall that

$$a(u, v) = \int_{\Omega} \nabla u \cdot \nabla v dx + \int_{\partial\Omega} uv ds.$$

Let  $w, v \in S_h$ .



## Solution to (c)

Recall that

$$a(u, v) = \int_{\Omega} \nabla u \cdot \nabla v dx + \int_{\partial\Omega} uv ds.$$

Let  $w, v \in S_h$ . **Q: What kind of function is  $\nabla w \cdot \nabla v$ ?**



## Solution to (c)

Recall that

$$a(u, v) = \int_{\Omega} \nabla u \cdot \nabla v dx + \int_{\partial\Omega} uv ds.$$

Let  $w, v \in S_h$ . **Q: What kind of function is  $\nabla w \cdot \nabla v$ ?**  $\nabla w \cdot \nabla v$  is a piecewise constant function. Thus, we see that



# Solution to (c)

Recall that

$$a(u, v) = \int_{\Omega} \nabla u \cdot \nabla v dx + \int_{\partial\Omega} uv ds.$$

Let  $w, v \in S_h$ . **Q: What kind of function is  $\nabla w \cdot \nabla v$ ?**  $\nabla w \cdot \nabla v$  is a piecewise constant function. Thus, we see that

$$\begin{aligned} \int_{\Omega} \nabla w \cdot \nabla v dx &= \sum_{\tau \in \mathcal{T}_h} \int_{\tau} \nabla w \cdot \nabla v dx \\ &= \sum_{\tau \in \mathcal{T}_h} (\nabla w \cdot \nabla v) \int_{\tau} dx \end{aligned}$$



## Solution to (c)

Recall that

$$a(u, v) = \int_{\Omega} \nabla u \cdot \nabla v dx + \int_{\partial\Omega} uv ds.$$

Let  $w, v \in S_h$ . **Q: What kind of function is  $\nabla w \cdot \nabla v$ ?**  $\nabla w \cdot \nabla v$  is a piecewise constant function. Thus, we see that

$$\begin{aligned} \int_{\Omega} \nabla w \cdot \nabla v dx &= \sum_{\tau \in \mathcal{T}_h} \int_{\tau} \nabla w \cdot \nabla v dx \\ &= \sum_{\tau \in \mathcal{T}_h} (\nabla w \cdot \nabla v) \int_{\tau} dx \\ &= \sum_{\tau \in \mathcal{T}_h} |\tau| (\nabla w \cdot \nabla v) \end{aligned}$$



# Solution to (c)

Recall that

$$a(u, v) = \int_{\Omega} \nabla u \cdot \nabla v dx + \int_{\partial\Omega} uv ds.$$

Let  $w, v \in S_h$ . **Q: What kind of function is  $\nabla w \cdot \nabla v$ ?**  $\nabla w \cdot \nabla v$  is a piecewise constant function. Thus, we see that

$$\begin{aligned} \int_{\Omega} \nabla w \cdot \nabla v dx &= \sum_{\tau \in \mathcal{T}_h} \int_{\tau} \nabla w \cdot \nabla v dx \\ &= \sum_{\tau \in \mathcal{T}_h} (\nabla w \cdot \nabla v) \int_{\tau} dx \\ &= \sum_{\tau \in \mathcal{T}_h} |\tau| (\nabla w \cdot \nabla v) \\ &= \sum_{\tau \in \mathcal{T}_h} \frac{|\tau|}{3} \left( (\nabla w \cdot \nabla v)(P_1) + (\nabla w \cdot \nabla v)(P_2) \right. \\ &\quad \left. + (\nabla w \cdot \nabla v)(P_3) \right) \end{aligned}$$



Now we want to compute the boundary term  $\int_{\partial\Omega} wvds$  for  $w, v \in S_h$ .





Now we want to compute the boundary term  $\int_{\partial\Omega} wv ds$  for  $w, v \in S_h$ . **Q:**  
**What kind of function is the product  $wv$  on a triangle edge?**



Now we want to compute the boundary term  $\int_{\partial\Omega} wv ds$  for  $w, v \in S_h$ . **Q:**  
**What kind of function is the product  $wv$  on a triangle edge?**  $wv$  is a quadratic one-dimensional polynomial.



Now we want to compute the boundary term  $\int_{\partial\Omega} wv ds$  for  $w, v \in S_h$ . **Q: What kind of function is the product  $wv$  on a triangle edge?**  $wv$  is a quadratic one-dimensional polynomial.

Recall that we are dealing with Gaussian quadrature which are exact for polynomials of degree  $2n - 1$  where  $n$  is the number of points used (2 in our case).



Now we want to compute the boundary term  $\int_{\partial\Omega} wv ds$  for  $w, v \in S_h$ . **Q: What kind of function is the product  $wv$  on a triangle edge?**  $wv$  is a quadratic one-dimensional polynomial.

Recall that we are dealing with Gaussian quadrature which are exact for polynomials of degree  $2n - 1$  where  $n$  is the number of points used (2 in our case).

Also recall that the Gaussian quadrature is defined on the interval  $[-1, 1]$ . Let  $e = [a, b]$  be an arbitrary edge.



Now we want to compute the boundary term  $\int_{\partial\Omega} wv ds$  for  $w, v \in S_h$ . **Q: What kind of function is the product  $wv$  on a triangle edge?**  $wv$  is a quadratic one-dimensional polynomial.

Recall that we are dealing with Gaussian quadrature which are exact for polynomials of degree  $2n - 1$  where  $n$  is the number of points used (2 in our case).

Also recall that the Gaussian quadrature is defined on the interval  $[-1, 1]$ . Let  $e = [a, b]$  be an arbitrary edge. Then, we need the transformation

$$T(t) = \frac{1}{2}(1 - t)a + \frac{1}{2}(t + a)b$$

so that  $T(-1) = a$  and  $T(1) = b$ . Note that  $T'(t) = \frac{1}{2}(b - a) = \frac{1}{2}|e|$ .



Now we can compute the boundary integral:

$$\begin{aligned}\int_{\partial\Omega} wv ds &= \sum_{e \in \partial\Omega} \int_e wv dx \\ &= \sum_{e \in \partial\Omega} \int_{-1}^1 wv \underbrace{\frac{1}{2}|e| dt} \\ &= \sum_{e \in \partial\Omega} \frac{|e|}{2} \left( (wv)(\alpha) + (wv)(\beta) \right)\end{aligned}$$



Now we can compute the boundary integral:

$$\begin{aligned}\int_{\partial\Omega} wv ds &= \sum_{e \in \partial\Omega} \int_e wv dx \\ &= \sum_{e \in \partial\Omega} \int_{-1}^1 wv \underbrace{\frac{1}{2}|e| dt} \\ &= \sum_{e \in \partial\Omega} \frac{|e|}{2} \left( (wv)(\alpha) + (wv)(\beta) \right)\end{aligned}$$

Thus, combining everything, we see that  $a_h(w, v) = a(w, v)$  for all  $w, v \in S_h$ .



## Solution to (d)

Using the estimate of Part (b) estimate the error  $\|u - u_h\|_{H^1}$ .





## Solution to (d)

We just showed that  $a_h(w, v) = a(w, v)$  for all  $w, v \in S_h$ .



## Solution to (d)

We just showed that  $a_h(w, v) = a(w, v)$  for all  $w, v \in S_h$ . Recall by Strang's first Lemma above, we know that

$$\begin{aligned} \|u - u_h\|_{H^1(\Omega)} \leq & \inf_{z_h \in S_h} \left(1 + \frac{C}{\alpha}\right) \|u - z_h\| + \\ & \frac{1}{\alpha} \left( \underbrace{\|a(z_h, v_h) - a_h(z_h, v_h)\|_{*,h}}_{=0} + \|F_h(v_h) - F(v_h)\|_{*,h} \right) \end{aligned}$$



## Solution to (d)

We just showed that  $a_h(w, v) = a(w, v)$  for all  $w, v \in S_h$ . Recall by Strang's first Lemma above, we know that

$$\begin{aligned}\|u - u_h\|_{H^1(\Omega)} &\leq \inf_{z_h \in S_h} \left(1 + \frac{C}{\alpha}\right) \|u - z_h\| + \\ &\quad \frac{1}{\alpha} \underbrace{\left(\|a(z_h, v_h) - a_h(z_h, v_h)\|_{*,h} + \|F_h(v_h) - F(v_h)\|_{*,h}\right)}_{=0} \\ &= \inf_{z_h \in S_h} \left(1 + \frac{C}{\alpha}\right) \|u - z_h\| + \frac{1}{\alpha} \|F_h(v_h) - F(v_h)\|_{*,h}\end{aligned}$$



## Solution to (d)

We just showed that  $a_h(w, v) = a(w, v)$  for all  $w, v \in S_h$ . Recall by Strang's first Lemma above, we know that

$$\begin{aligned}\|u - u_h\|_{H^1(\Omega)} &\leq \inf_{z_h \in S_h} \left(1 + \frac{C}{\alpha}\right) \|u - z_h\| + \\ &\quad \frac{1}{\alpha} \underbrace{\left(\|a(z_h, v_h) - a_h(z_h, v_h)\|_{*,h} + \|F_h(v_h) - F(v_h)\|_{*,h}\right)}_{=0} \\ &= \inf_{z_h \in S_h} \left(1 + \frac{C}{\alpha}\right) \|u - z_h\| + \frac{1}{\alpha} \|F_h(v_h) - F(v_h)\|_{*,h}\end{aligned}$$

**This is justified since  $a_h$  is  $V_h$ -elliptic since  $a$  is  $V$ -elliptic and  $a_h = a$  on  $V_h$ .**



## Solution to (d)

We just showed that  $a_h(w, v) = a(w, v)$  for all  $w, v \in S_h$ . Recall by Strang's first Lemma above, we know that

$$\begin{aligned}\|u - u_h\|_{H^1(\Omega)} &\leq \inf_{z_h \in S_h} \left(1 + \frac{C}{\alpha}\right) \|u - z_h\| + \\ &\quad \frac{1}{\alpha} \underbrace{\left(\|a(z_h, v_h) - a_h(z_h, v_h)\|_{*,h} + \|F_h(v_h) - F(v_h)\|_{*,h}\right)}_{=0} \\ &= \inf_{z_h \in S_h} \left(1 + \frac{C}{\alpha}\right) \|u - z_h\| + \frac{1}{\alpha} \|F_h(v_h) - F(v_h)\|_{*,h}\end{aligned}$$

**This is justified since  $a_h$  is  $V_h$ -elliptic since  $a$  is  $V$ -elliptic and  $a_h = a$  on  $V_h$ .** Since we want an error estimate, we need to do the following:



## Solution to (d)

We just showed that  $a_h(w, v) = a(w, v)$  for all  $w, v \in S_h$ . Recall by Strang's first Lemma above, we know that

$$\begin{aligned}\|u - u_h\|_{H^1(\Omega)} &\leq \inf_{z_h \in S_h} \left(1 + \frac{C}{\alpha}\right) \|u - z_h\| + \\ &\quad \frac{1}{\alpha} \underbrace{\left(\|a(z_h, v_h) - a_h(z_h, v_h)\|_{*,h} + \|F_h(v_h) - F(v_h)\|_{*,h}\right)}_{=0} \\ &= \inf_{z_h \in S_h} \left(1 + \frac{C}{\alpha}\right) \|u - z_h\| + \frac{1}{\alpha} \|F_h(v_h) - F(v_h)\|_{*,h}\end{aligned}$$

**This is justified since  $a_h$  is  $V_h$ -elliptic since  $a$  is  $V$ -elliptic and  $a_h = a$  on  $V_h$ .** Since we want an error estimate, we need to do the following: **1. Define a projection;**



## Solution to (d)

We just showed that  $a_h(w, v) = a(w, v)$  for all  $w, v \in S_h$ . Recall by Strang's first Lemma above, we know that

$$\begin{aligned}\|u - u_h\|_{H^1(\Omega)} &\leq \inf_{z_h \in S_h} \left(1 + \frac{C}{\alpha}\right) \|u - z_h\| + \\ &\quad \frac{1}{\alpha} \underbrace{\left(\|a(z_h, v_h) - a_h(z_h, v_h)\|_{*,h} + \|F_h(v_h) - F(v_h)\|_{*,h}\right)}_{=0} \\ &= \inf_{z_h \in S_h} \left(1 + \frac{C}{\alpha}\right) \|u - z_h\| + \frac{1}{\alpha} \|F_h(v_h) - F(v_h)\|_{*,h}\end{aligned}$$

**This is justified since  $a_h$  is  $V_h$ -elliptic since  $a$  is  $V$ -elliptic and  $a_h = a$  on  $V_h$ .** Since we want an error estimate, we need to do the following: **1. Define a projection; 2. Transform to the reference element;**



## Solution to (d)

We just showed that  $a_h(w, v) = a(w, v)$  for all  $w, v \in S_h$ . Recall by Strang's first Lemma above, we know that

$$\begin{aligned}\|u - u_h\|_{H^1(\Omega)} &\leq \inf_{z_h \in S_h} \left(1 + \frac{C}{\alpha}\right) \|u - z_h\| + \\ &\quad \frac{1}{\alpha} \underbrace{\left(\|a(z_h, v_h) - a_h(z_h, v_h)\|_{*,h} + \|F_h(v_h) - F(v_h)\|_{*,h}\right)}_{=0} \\ &= \inf_{z_h \in S_h} \left(1 + \frac{C}{\alpha}\right) \|u - z_h\| + \frac{1}{\alpha} \|F_h(v_h) - F(v_h)\|_{*,h}\end{aligned}$$

**This is justified since  $a_h$  is  $V_h$ -elliptic since  $a$  is  $V$ -elliptic and  $a_h = a$  on  $V_h$ .** Since we want an error estimate, we need to do the following: **1. Define a projection; 2. Transform to the reference element; 3. Use Bramble-Hilbert to get error on reference element;**





## Solution to (d)

We just showed that  $a_h(w, v) = a(w, v)$  for all  $w, v \in S_h$ . Recall by Strang's first Lemma above, we know that

$$\begin{aligned}\|u - u_h\|_{H^1(\Omega)} &\leq \inf_{z_h \in S_h} \left(1 + \frac{C}{\alpha}\right) \|u - z_h\| + \\ &\quad \frac{1}{\alpha} \underbrace{\left(\|a(z_h, v_h) - a_h(z_h, v_h)\|_{*,h} + \|F_h(v_h) - F(v_h)\|_{*,h}\right)}_{=0} \\ &= \inf_{z_h \in S_h} \left(1 + \frac{C}{\alpha}\right) \|u - z_h\| + \frac{1}{\alpha} \|F_h(v_h) - F(v_h)\|_{*,h}\end{aligned}$$

**This is justified since  $a_h$  is  $V_h$ -elliptic since  $a$  is  $V$ -elliptic and  $a_h = a$  on  $V_h$ .** Since we want an error estimate, we need to do the following: **1. Define a projection; 2. Transform to the reference element; 3. Use Bramble-Hilbert to get error on reference element; 4. Transform back to physical element;**



# Review of affine transformations

For reference, see 4.4.1 of Grossman book pg 219.

- We first define the reference element as the triangle:

$$\hat{\tau} = \left\{ \begin{pmatrix} \hat{x} \\ \hat{y} \end{pmatrix} : \hat{x} \geq 0, \hat{y} \geq 0, \hat{x} + \hat{y} \leq 1 \right\}. \quad (1.9)$$



# Review of affine transformations

For reference, see 4.4.1 of Grossman book pg 219.

- We first define the reference element as the triangle:

$$\hat{\tau} = \left\{ \begin{pmatrix} \hat{x} \\ \hat{y} \end{pmatrix} : \hat{x} \geq 0, \hat{y} \geq 0, \hat{x} + \hat{y} \leq 1 \right\}. \quad (1.9)$$

- Consider the mapping  $F_{\tau} : \hat{\tau} \rightarrow \tau$  that goes from the reference element to the triangle  $\tau$  (physical element) with the vertices  $(x_1, y_1)^T, (x_2, y_2)^T, (x_3, y_3)^T$ :

$$F_{\tau}(\hat{p}) = \underbrace{\begin{pmatrix} x_2 - x_1 & x_3 - x_1 \\ y_2 - y_1 & y_3 - y_1 \end{pmatrix}}_B \begin{pmatrix} \hat{x} \\ \hat{y} \end{pmatrix} + \begin{pmatrix} x_1 \\ y_1 \end{pmatrix} \quad (1.10)$$

**Note that**  $|\det B| = \left| \det(F'_{\tau}(\hat{p})) \right| = \frac{|\tau|}{|\hat{\tau}|} \sim ch^2$ . ( $h$  is defined on the next slide)



- Setting  $v(\hat{p}) = u(F(\hat{p}))$  for  $\hat{p} \in \hat{\tau}$ , we see that each function  $u(\mathbf{x})$  for  $\mathbf{x} \in \tau$  is mapped to a function  $v(\hat{p})$  defined on the reference element.

**Notation: We will interchange  $v(\hat{p}) = u(F(\hat{p}))$  with  $\hat{u}$ .**



- Setting  $v(\hat{p}) = u(F(\hat{p}))$  for  $\hat{p} \in \hat{\tau}$ , we see that each function  $u(\mathbf{x})$  for  $\mathbf{x} \in \tau$  is mapped to a function  $v(\hat{p})$  defined on the reference element.

**Notation: We will interchange  $v(\hat{p}) = u(F(\hat{p}))$  with  $\hat{u}$ .**

- So, if  $F$  is differentiable, then by the chain rule, we have that:

$$\nabla_{\hat{p}} v(\hat{p}) = F'(\hat{p}) \nabla u(F(\hat{p})). \quad (1.11)$$



- Setting  $v(\hat{p}) = u(F(\hat{p}))$  for  $\hat{p} \in \hat{\tau}$ , we see that each function  $u(\mathbf{x})$  for  $\mathbf{x} \in \tau$  is mapped to a function  $v(\hat{p})$  defined on the reference element.

**Notation: We will interchange  $v(\hat{p}) = u(F(\hat{p}))$  with  $\hat{u}$ .**

- So, if  $F$  is differentiable, then by the chain rule, we have that:

$$\nabla_{\hat{p}} v(\hat{p}) = F'(\hat{p}) \nabla u(F(\hat{p})). \quad (1.11)$$

- Let  $h$  be the maximum length of the edges of the triangles in the triangulation of  $\Omega$ . Then there exists  $c > 0$  such that

$$\|F'(\hat{p})\| = \|B\| \leq ch \quad \text{for all } \hat{p} \in \hat{\tau} \quad (1.12)$$



- Setting  $v(\hat{p}) = u(F(\hat{p}))$  for  $\hat{p} \in \hat{\tau}$ , we see that each function  $u(\mathbf{x})$  for  $\mathbf{x} \in \tau$  is mapped to a function  $v(\hat{p})$  defined on the reference element.

**Notation: We will interchange  $v(\hat{p}) = u(F(\hat{p}))$  with  $\hat{u}$ .**

- So, if  $F$  is differentiable, then by the chain rule, we have that:

$$\nabla_{\hat{p}} v(\hat{p}) = F'(\hat{p}) \nabla u(F(\hat{p})). \quad (1.11)$$

- Let  $h$  be the maximum length of the edges of the triangles in the triangulation of  $\Omega$ . Then there exists  $c > 0$  such that

$$\|F'(\hat{p})\| = \|B\| \leq ch \quad \text{for all } \hat{p} \in \hat{\tau} \quad (1.12)$$

- Using the above we see that

$$\int_{\tau} u^2(\mathbf{x}) d\mathbf{x} = \int_{\hat{\tau}} v^2(\hat{p}) \left| \det F'(\hat{p}) \right| d\hat{p} \quad (1.13)$$



- According to Lemma 4.23 in Grossman book:

$$\int_{\tau} |(\nabla u)|^2 d\mathbf{x} \leq c \|B^{-1}\|^2 \int_{\hat{\tau}} |\nabla_{\hat{\rho}} \hat{u}|^2 |\det B| d\hat{\rho} \quad (1.14)$$

---

<sup>3</sup>see pg 222 of Grossman book





- According to Lemma 4.23 in Grossman book:

$$\int_{\tau} |(\nabla u)|^2 d\mathbf{x} \leq c \|B^{-1}\|^2 \int_{\hat{\tau}} |\nabla_{\hat{\rho}} \hat{u}|^2 |\det B| d\hat{\rho} \quad (1.14)$$

- Assuming that the mesh is shape-regular<sup>3</sup> (which we can always do for these error estimates), we have the following bound:

$$\|B^{-1}\| \leq ch^{-1}.$$

---

<sup>3</sup>see pg 222 of Grossman book



# A note on Bramble-Hilbert

For applying Bramble-Hilbert (and any other major result), it is recommended that students first state the result in their solution and then apply it appropriately (by making sure the assumptions are satisfied). You should not just say “by Bramble-Hilbert” in a sub-step and move on.



# A note on Bramble-Hilbert

## Bramble-Hilbert

Let  $B \subset \mathbb{R}^n$  be a domain with a Lipschitz boundary and let  $q$  be a bounded sub-linear functional on  $H^{k+1}(B)$ . Assume that

$$q(w) = 0, \quad \text{for all } w \in P^k.$$

Then there exists a constant  $c = c(B) > 0$ , which depends on  $B$ , such that

$$|q(v)| \leq c |v|_{k+1,B}, \quad \text{for all } v \in H^{k+1}(B).$$



## Back to solution...

We first focus on the term  $\inf_{z_h \in S_h} \left(1 + \frac{c}{\alpha}\right) \|u - z_h\|$ .



## Back to solution...

We first focus on the term  $\inf_{z_h \in S_h} \left(1 + \frac{c}{\alpha}\right) \|u - z_h\|$ . Let

$\Pi_h : H^1(\Omega) \rightarrow S_h$  be the projection onto the finite element subspace.



## Back to solution...

We first focus on the term  $\inf_{z_h \in S_h} \left(1 + \frac{C}{\alpha}\right) \|u - z_h\|$ . Let

$\Pi_h : H^1(\Omega) \rightarrow S_h$  be the projection onto the finite element subspace.

Recall that the projection error bounds this term from above:

$$\left( \inf_{z_h \in S_h} \left(1 + \frac{C}{\alpha}\right) \|u - z_h\| \right)^2 \leq c \|u - \Pi_h u\|_{H^1(\Omega)}^2 \quad (1.15)$$



## Back to solution...

We first focus on the term  $\inf_{z_h \in S_h} \left(1 + \frac{C}{\alpha}\right) \|u - z_h\|$ . Let

$\Pi_h : H^1(\Omega) \rightarrow S_h$  be the projection onto the finite element subspace.

Recall that the projection error bounds this term from above:

$$\left( \inf_{z_h \in S_h} \left(1 + \frac{C}{\alpha}\right) \|u - z_h\| \right)^2 \leq c \|u - \Pi_h u\|_{H^1(\Omega)}^2 \quad (1.15)$$

We want to compute this term on the reference element:

$$\|u - \Pi_h u\|_{H^1(\Omega)}^2 = \sum_{\tau \in \mathcal{T}_h} \|u - \Pi_h u\|_{H^1(\tau)}^2$$



## Back to solution...

We first focus on the term  $\inf_{z_h \in S_h} \left(1 + \frac{C}{\alpha}\right) \|u - z_h\|$ . Let

$\Pi_h : H^1(\Omega) \rightarrow S_h$  be the projection onto the finite element subspace.

Recall that the projection error bounds this term from above:

$$\left( \inf_{z_h \in S_h} \left(1 + \frac{C}{\alpha}\right) \|u - z_h\| \right)^2 \leq c \|u - \Pi_h u\|_{H^1(\Omega)}^2 \quad (1.15)$$

We want to compute this term on the reference element:

$$\begin{aligned} \|u - \Pi_h u\|_{H^1(\Omega)}^2 &= \sum_{\tau \in \mathcal{T}_h} \|u - \Pi_h u\|_{H^1(\tau)}^2 \\ &= \sum_{\tau \in \mathcal{T}_h} \int_{\tau} \left( |u - \Pi_h u|^2 + |\nabla(u - \Pi_h u)|^2 \right) dx \\ &\leq c \sum_{\tau \in \mathcal{T}_h} \int_{\hat{\tau}} \left( \left| \hat{u} - \widehat{(\Pi_h u)} \right|^2 + \right. \end{aligned}$$





## Back to solution...

We first focus on the term  $\inf_{z_h \in S_h} \left(1 + \frac{C}{\alpha}\right) \|u - z_h\|$ . Let

$\Pi_h : H^1(\Omega) \rightarrow S_h$  be the projection onto the finite element subspace.

Recall that the projection error bounds this term from above:

$$\left( \inf_{z_h \in S_h} \left(1 + \frac{C}{\alpha}\right) \|u - z_h\| \right)^2 \leq c \|u - \Pi_h u\|_{H^1(\Omega)}^2 \quad (1.15)$$

We want to compute this term on the reference element:

$$\begin{aligned} \|u - \Pi_h u\|_{H^1(\Omega)}^2 &= \sum_{\tau \in \mathcal{T}_h} \|u - \Pi_h u\|_{H^1(\tau)}^2 \\ &= \sum_{\tau \in \mathcal{T}_h} \int_{\tau} \left( |u - \Pi_h u|^2 + |\nabla(u - \Pi_h u)|^2 \right) dx \\ &\leq c \sum_{\tau \in \mathcal{T}_h} \int_{\hat{\tau}} \left( \left| \hat{u} - \widehat{(\Pi_h u)} \right|^2 + \right. \\ &\quad \left. \|B^{-1}\|^2 \left| \nabla_{\hat{p}}(\hat{u} - \widehat{(\Pi_h u)}) \right|^2 \right) \underbrace{\frac{|\tau|}{|\hat{\tau}|} d\hat{p}} \end{aligned}$$



In this next step, we bound  $\|B^{-1}\| \leq ch^{-1}$  and replace  $\frac{|\tau|}{|\hat{\tau}|}$  by  $ch^2$ :

$$\begin{aligned} \|u - \Pi_h u\|_{H^1(\Omega)}^2 &\leq ch^2 \sum_{\tau \in \mathcal{T}_h} \int_{\hat{\tau}} \left| \hat{u} - \widehat{(\Pi_h u)} \right|^2 + \\ &\quad c \sum_{\tau \in \mathcal{T}_h} \int_{\hat{\tau}} \left| \nabla(\hat{u} - \widehat{(\Pi_h u)}) \right|^2 d\hat{p} \end{aligned}$$



In this next step, we bound  $\|B^{-1}\| \leq ch^{-1}$  and replace  $\left|\frac{\tau}{\widehat{\tau}}\right|$  by  $ch^2$ :

$$\begin{aligned}\|u - \Pi_h u\|_{H^1(\Omega)}^2 &\leq ch^2 \sum_{\tau \in \mathcal{T}_h} \int_{\widehat{\tau}} \left| \widehat{u} - (\widehat{\Pi_h u}) \right|^2 + \\ &\quad c \sum_{\tau \in \mathcal{T}_h} \int_{\widehat{\tau}} \left| \nabla(\widehat{u} - (\widehat{\Pi_h u})) \right|^2 d\widehat{p} \\ &= ch^2 \sum_{\tau \in \mathcal{T}_h} \|(\text{Id} - \widehat{\Pi}_h) \widehat{u}\|^2 + \\ &\quad c \sum_{\tau \in \mathcal{T}_h} \left| (\text{Id} - \widehat{\Pi}_h) \widehat{u} \right|_{H^1(\Omega)}^2\end{aligned}$$



In this next step, we bound  $\|B^{-1}\| \leq ch^{-1}$  and replace  $\frac{|\tau|}{|\hat{\tau}|}$  by  $ch^2$ :

$$\begin{aligned}
 \|u - \Pi_h u\|_{H^1(\Omega)}^2 &\leq ch^2 \sum_{\tau \in \mathcal{T}_h} \int_{\hat{\tau}} \left| \hat{u} - \widehat{(\Pi_h u)} \right|^2 + \\
 &\quad c \sum_{\tau \in \mathcal{T}_h} \int_{\hat{\tau}} \left| \nabla(\hat{u} - \widehat{(\Pi_h u)}) \right|^2 d\hat{p} \\
 &= ch^2 \sum_{\tau \in \mathcal{T}_h} \|(\text{Id} - \hat{\Pi}_h)\hat{u}\|^2 + \\
 &\quad c \sum_{\tau \in \mathcal{T}_h} \left| (\text{Id} - \hat{\Pi}_h)\hat{u} \right|_{H^1(\Omega)}^2
 \end{aligned}$$

Notice that  $\|(\text{Id} - \hat{\Pi}_h)(\cdot)\|_{L^2(\tau)}$  and  $|(\text{Id} - \hat{\Pi}_h)(\cdot)|_{H^1(\tau)}$  are both sublinear functionals defined on  $H^2(\hat{\tau})$  and are zero exactly zero for linear polynomials on  $\tau$ , therefore the Bramble-Hilbert lemma can be applied.



**Recall:**  $\|B\| \leq ch$ ,  $\frac{|\hat{\tau}|}{|\tau|} \rightarrow ch^{-2}$

$$\sum_{\tau \in \mathcal{T}_h} \|u - \Pi_h u\|_{H^1(\tau)}^2 \leq ch^2 \sum_{\tau \in \mathcal{T}_h} |\hat{u}|_{H^2(\hat{\tau})}^2 + c \sum_{\tau \in \mathcal{T}_h} |\hat{u}|_{H^2(\hat{\tau})}^2$$



**Recall:**  $\|B\| \leq ch$ ,  $\frac{|\hat{\tau}|}{|\tau|} \rightarrow ch^{-2}$

$$\begin{aligned} \sum_{\tau \in \mathcal{T}_h} \|u - \Pi_h u\|_{H^1(\tau)}^2 &\leq ch^2 \sum_{\tau \in \mathcal{T}_h} |\hat{u}|_{H^2(\hat{\tau})}^2 + c \sum_{\tau \in \mathcal{T}_h} |\hat{u}|_{H^2(\hat{\tau})}^2 \\ &\leq c(h^2 + 1) \sum_{\tau \in \mathcal{T}_h} \int_{\hat{\tau}} \sum_{|\alpha|=2} |\hat{D}^\alpha \hat{u}|^2 d\hat{p} \end{aligned}$$



**Recall:**  $\|B\| \leq ch$ ,  $\frac{|\hat{\tau}|}{|\tau|} \rightarrow ch^{-2}$

$$\begin{aligned}
 \sum_{\tau \in \mathcal{T}_h} \|u - \Pi_h u\|_{H^1(\tau)}^2 &\leq ch^2 \sum_{\tau \in \mathcal{T}_h} |\hat{u}|_{H^2(\hat{\tau})}^2 + c \sum_{\tau \in \mathcal{T}_h} |\hat{u}|_{H^2(\hat{\tau})}^2 \\
 &\leq c(h^2 + 1) \sum_{\tau \in \mathcal{T}_h} \int_{\hat{\tau}} \sum_{|\alpha|=2} |\hat{D}^\alpha \hat{u}|^2 d\hat{p} \\
 \text{(Lemma 4.23(ii)) } &\leq c(h^2 + 1) \sum_{\tau \in \mathcal{T}_h} \int_{\tau} \sum_{|\alpha|=2} \underbrace{\|B\|^4}_{\substack{\text{ } \\ \text{ }}} |D^\alpha u|^2 \underbrace{\frac{|\hat{\tau}|}{|\tau|}}_{\substack{\text{ } \\ \text{ }}} dx
 \end{aligned}$$



**Recall:**  $\|B\| \leq ch$ ,  $\frac{|\hat{\tau}|}{|\tau|} \rightarrow ch^{-2}$

$$\begin{aligned}
 \sum_{\tau \in \mathcal{T}_h} \|u - \Pi_h u\|_{H^1(\tau)}^2 &\leq ch^2 \sum_{\tau \in \mathcal{T}_h} |\hat{u}|_{H^2(\hat{\tau})}^2 + c \sum_{\tau \in \mathcal{T}_h} |\hat{u}|_{H^2(\hat{\tau})}^2 \\
 &\leq c(h^2 + 1) \sum_{\tau \in \mathcal{T}_h} \int_{\hat{\tau}} \sum_{|\alpha|=2} |\hat{D}^\alpha \hat{u}|^2 d\hat{p} \\
 (\text{Lemma 4.23(ii)}) \quad &\leq c(h^2 + 1) \sum_{\tau \in \mathcal{T}_h} \int_{\tau} \sum_{|\alpha|=2} \underbrace{\|B\|^4}_{\substack{\text{ } \\ \text{ }}} |D^\alpha u|^2 \underbrace{\frac{|\hat{\tau}|}{|\tau|}}_{\substack{\text{ } \\ \text{ }}} dx \\
 &\leq c(h^2 + 1) \sum_{\tau \in \mathcal{T}_h} \int_{\tau} h^2 \sum_{|\alpha|=2} |D^\alpha u|^2 dx \\
 &= c(h^4 + h^2) \sum_{\tau \in \mathcal{T}_h} \int_{\tau} \sum_{|\alpha|=2} |D^\alpha u|^2 dx
 \end{aligned}$$





**Recall:**  $\|B\| \leq ch, \frac{|\hat{\tau}|}{|\tau|} \rightarrow ch^{-2}$

$$\begin{aligned}
 \sum_{\tau \in \mathcal{T}_h} \|u - \Pi_h u\|_{H^1(\tau)}^2 &\leq ch^2 \sum_{\tau \in \mathcal{T}_h} |\hat{u}|_{H^2(\hat{\tau})}^2 + c \sum_{\tau \in \mathcal{T}_h} |\hat{u}|_{H^2(\hat{\tau})}^2 \\
 &\leq c(h^2 + 1) \sum_{\tau \in \mathcal{T}_h} \int_{\hat{\tau}} \sum_{|\alpha|=2} |\hat{D}^\alpha \hat{u}|^2 d\hat{p} \\
 (\text{Lemma 4.23(ii)}) &\leq c(h^2 + 1) \sum_{\tau \in \mathcal{T}_h} \int_{\tau} \sum_{|\alpha|=2} \underbrace{\|B\|^4}_{\leq ch^4} |D^\alpha u|^2 \underbrace{\frac{|\hat{\tau}|}{|\tau|}}_{\leq ch^{-2}} dx \\
 &\leq c(h^2 + 1) \sum_{\tau \in \mathcal{T}_h} \int_{\tau} h^2 \sum_{|\alpha|=2} |D^\alpha u|^2 dx \\
 &= c(h^4 + h^2) \sum_{\tau \in \mathcal{T}_h} \int_{\tau} \sum_{|\alpha|=2} |D^\alpha u|^2 dx \\
 &\leq ch^2 |u|_{H^2(\Omega)}^2.
 \end{aligned}$$

