

Lecture 5

Eric Tovar

Qualifying Prep Course – Numerical

07-09-2020



Outline

- 1 January 2010 Exam
 - Problem 2
- 2 August 2009 Exam
 - Problem 1
- 3 January 2010 Problem 3



Problem 2

Consider the two finite elements (τ, Q_1, Σ) and $(\tau, \tilde{Q}_1, \Sigma)$ where $\tau = [-1, 1]^2$ is the reference square and

$$Q_1 = \text{span}\{1, x, y, xy\} \quad (1.1)$$

$$\tilde{Q}_1 = \text{span}\{1, x, y, x^2 - y^2\}. \quad (1.2)$$

$\Sigma = \{w(1, 0), w(-1, 0), w(0, 1), w(0, -1)\}$ is the set of values of a function $w(x, y)$ at the midpoints of the edges τ .



Problem 2

Consider the two finite elements (τ, Q_1, Σ) and $(\tau, \tilde{Q}_1, \Sigma)$ where $\tau = [-1, 1]^2$ is the reference square and

$$Q_1 = \text{span}\{1, x, y, xy\} \quad (1.1)$$

$$\tilde{Q}_1 = \text{span}\{1, x, y, x^2 - y^2\}. \quad (1.2)$$

$\Sigma = \{w(1, 0), w(-1, 0), w(0, 1), w(0, -1)\}$ is the set of values of a function $w(x, y)$ at the midpoints of the edges τ .

- (a) Which of the two elements is unisolvent? Prove it!
- (b) Show that the unisolvent element leads to a finite element space, which is not H^1 -conforming.



Definition 5.2 (Finite element). Let $d \geq 1$, an integer $n_{\text{sh}} \geq 1$, and the set $\mathcal{N} := \{1:n_{\text{sh}}\}$. A finite element consists of a triple (K, P, Σ) where:

- (i) K is a polyhedron in \mathbb{R}^d or the image of a polyhedron in \mathbb{R}^d by some smooth diffeomorphism. More generally, K could be the closure of a Lipschitz domain in \mathbb{R}^d (see §3.1). K is nontrivial, i.e., $\text{int}(K) \neq \emptyset$.
- (ii) P is a finite-dimensional vector space of functions $p : K \rightarrow \mathbb{R}^q$ for some integer $q \geq 1$ (typically $q \in \{1, d\}$). P is nontrivial, i.e., $P \neq \{0\}$. The members of P are polynomial functions, possibly composed with some smooth diffeomorphism.
- (iii) Σ is a set of n_{sh} linear forms from P to \mathbb{R} , say $\Sigma := \{\sigma_i\}_{i \in \mathcal{N}}$, such that the map $\Phi_\Sigma : P \rightarrow \mathbb{R}^{n_{\text{sh}}}$ defined by $\Phi_\Sigma(p) := (\sigma_i(p))_{i \in \mathcal{N}}$ is an isomorphism. The linear forms σ_i are called degrees of freedom (in short dofs), and the bijectivity of the map Φ_Σ is referred to as unisolvence.

Remark 5.3 (Proving unisolvence). To prove unisolvence, it suffices to show that $\dim P \geq n_{\text{sh}} = \text{card } \Sigma$ and that Φ_Σ is injective, i.e.,

$$[\sigma_i(p) = 0, \forall i \in \mathcal{N}] \implies [p = 0], \quad \forall p \in P. \quad (5.4)$$

Owing to the rank nullity theorem Φ_Σ is then bijective and $\dim P = n_{\text{sh}}$. \square



Solution to (a)

We want to show that either (τ, Q_1, Σ) or $(\tau, \tilde{Q}_1, \Sigma)$ is unisolvent.



Solution to (a)

We want to show that either (τ, Q_1, Σ) or $(\tau, \tilde{Q}_1, \Sigma)$ is unisolvent. Let us start with (τ, Q_1, Σ) . We first note that $\text{card}(\Sigma) = 4 = \dim Q_1$.



Solution to (a)

We want to show that either (τ, Q_1, Σ) or $(\tau, \tilde{Q}_1, \Sigma)$ is unisolvent. Let us start with (τ, Q_1, Σ) . We first note that $\text{card}(\Sigma) = 4 = \dim Q_1$.

We now want to check that $\Sigma \ni \sigma_i(p) = 0 \forall i \in \{1, 2, 3, 4\}$ implies that $p = 0$ for any $p \in Q_1$.



Solution to (a)

We want to show that either (τ, Q_1, Σ) or $(\tau, \tilde{Q}_1, \Sigma)$ is unisolvent. Let us start with (τ, Q_1, Σ) . We first note that $\text{card}(\Sigma) = 4 = \dim Q_1$.

We now want to check that $\Sigma \ni \sigma_i(p) = 0 \forall i \in \{1, 2, 3, 4\}$ implies that $p = 0$ for any $p \in Q_1$. Let $w(x, y)$ be an arbitrary polynomial in Q_1 :

$$w(x, y) = a + bx + cy + dxy$$



Then note that for each σ_i , $i \in \{1, 2, 3, 4\}$:



Then note that for each σ_i , $i \in \{1, 2, 3, 4\}$:

$$\sigma_1(w) = w(1, 0) = a + b = 0 \quad (1.3)$$

$$\sigma_2(w) = w(-1, 0) = a - b = 0 \quad (1.4)$$

$$\sigma_3(w) = w(0, 1) = a + c = 0 \quad (1.5)$$

$$\sigma_4(w) = w(0, -1) = a - c = 0. \quad (1.6)$$



Then note that for each σ_i , $i \in \{1, 2, 3, 4\}$:

$$\sigma_1(w) = w(1, 0) = a + b = 0 \quad (1.3)$$

$$\sigma_2(w) = w(-1, 0) = a - b = 0 \quad (1.4)$$

$$\sigma_3(w) = w(0, 1) = a + c = 0 \quad (1.5)$$

$$\sigma_4(w) = w(0, -1) = a - c = 0. \quad (1.6)$$

Then (1.3) implies $a = -b$; (1.4) implies $b = 0 \implies a = 0$; (1.5) implies $c = 0$.



Then note that for each σ_i , $i \in \{1, 2, 3, 4\}$:

$$\sigma_1(w) = w(1, 0) = a + b = 0 \quad (1.3)$$

$$\sigma_2(w) = w(-1, 0) = a - b = 0 \quad (1.4)$$

$$\sigma_3(w) = w(0, 1) = a + c = 0 \quad (1.5)$$

$$\sigma_4(w) = w(0, -1) = a - c = 0. \quad (1.6)$$

Then (1.3) implies $a = -b$; (1.4) implies $b = 0 \implies a = 0$; (1.5) implies $c = 0$. However, d is a free variable and can be chosen to be any value.



Then note that for each σ_i , $i \in \{1, 2, 3, 4\}$:

$$\sigma_1(w) = w(1, 0) = a + b = 0 \quad (1.3)$$

$$\sigma_2(w) = w(-1, 0) = a - b = 0 \quad (1.4)$$

$$\sigma_3(w) = w(0, 1) = a + c = 0 \quad (1.5)$$

$$\sigma_4(w) = w(0, -1) = a - c = 0. \quad (1.6)$$

Then (1.3) implies $a = -b$; (1.4) implies $b = 0 \implies a = 0$; (1.5) implies $c = 0$. However, d is a free variable and can be chosen to be any value. Thus, $w(x, y)$ is not identically zero and consequently (τ, Q_1, σ) is **not** unisolvent.



We now want to check to see if $(\tau, \tilde{Q}_1, \sigma)$ is unisolvent.



We now want to check to see if $(\tau, \tilde{Q}_1, \sigma)$ is unisolvent. We repeat the same process as above.



We now want to check to see if $(\tau, \tilde{Q}_1, \sigma)$ is unisolvent. We repeat the same process as above. Now, let $w(x, y)$ be an arbitrary polynomial in \tilde{Q}_1 :

$$w(x, y) = a + bx + cy + d(x^2 - y^2)$$



Then, for each σ_i , $i \in \{1, 2, 3, 4\}$ we have that:



Then, for each σ_i , $i \in \{1, 2, 3, 4\}$ we have that:

$$\sigma_1(w) = w(1, 0) = a + b + d = 0 \quad (1.7)$$

$$\sigma_2(w) = w(-1, 0) = a - b + d = 0 \quad (1.8)$$

$$\sigma_3(w) = w(0, 1) = a + c - d = 0 \quad (1.9)$$

$$\sigma_4(w) = w(0, -1) = a - c - d = 0. \quad (1.10)$$



Then, for each σ_i , $i \in \{1, 2, 3, 4\}$ we have that:

$$\sigma_1(w) = w(1, 0) = a + b + d = 0 \quad (1.7)$$

$$\sigma_2(w) = w(-1, 0) = a - b + d = 0 \quad (1.8)$$

$$\sigma_3(w) = w(0, 1) = a + c - d = 0 \quad (1.9)$$

$$\sigma_4(w) = w(0, -1) = a - c - d = 0. \quad (1.10)$$

Then (1.7) implies $a = -(b + d)$;



Then, for each σ_i , $i \in \{1, 2, 3, 4\}$ we have that:

$$\sigma_1(w) = w(1, 0) = a + b + d = 0 \quad (1.7)$$

$$\sigma_2(w) = w(-1, 0) = a - b + d = 0 \quad (1.8)$$

$$\sigma_3(w) = w(0, 1) = a + c - d = 0 \quad (1.9)$$

$$\sigma_4(w) = w(0, -1) = a - c - d = 0. \quad (1.10)$$

Then (1.7) implies $a = -(b + d)$; (1.8) implies

$$-(b + d) + d = 0 \implies b = 0 \implies a = -d;$$



Then, for each σ_i , $i \in \{1, 2, 3, 4\}$ we have that:

$$\sigma_1(w) = w(1, 0) = a + b + d = 0 \quad (1.7)$$

$$\sigma_2(w) = w(-1, 0) = a - b + d = 0 \quad (1.8)$$

$$\sigma_3(w) = w(0, 1) = a + c - d = 0 \quad (1.9)$$

$$\sigma_4(w) = w(0, -1) = a - c - d = 0. \quad (1.10)$$

Then (1.7) implies $a = -(b + d)$; (1.8) implies $-(b + d) + d = 0 \implies b = 0 \implies a = -d$; (1.9) implies $c - 2d = 0$.



Then, for each σ_i , $i \in \{1, 2, 3, 4\}$ we have that:

$$\sigma_1(w) = w(1, 0) = a + b + d = 0 \quad (1.7)$$

$$\sigma_2(w) = w(-1, 0) = a - b + d = 0 \quad (1.8)$$

$$\sigma_3(w) = w(0, 1) = a + c - d = 0 \quad (1.9)$$

$$\sigma_4(w) = w(0, -1) = a - c - d = 0. \quad (1.10)$$

Then (1.7) implies $a = -(b + d)$; (1.8) implies

$$-(b + d) + d = 0 \implies b = 0 \implies a = -d; \text{ (1.9) implies } c - 2d = 0.$$

$$\text{(1.10) implies } -d + (2d) - d = 0 \implies d = 0 \implies c = 0 \implies a = 0.$$



Then, for each σ_i , $i \in \{1, 2, 3, 4\}$ we have that:

$$\sigma_1(w) = w(1, 0) = a + b + d = 0 \quad (1.7)$$

$$\sigma_2(w) = w(-1, 0) = a - b + d = 0 \quad (1.8)$$

$$\sigma_3(w) = w(0, 1) = a + c - d = 0 \quad (1.9)$$

$$\sigma_4(w) = w(0, -1) = a - c - d = 0. \quad (1.10)$$

Then (1.7) implies $a = -(b + d)$; (1.8) implies $-(b + d) + d = 0 \implies b = 0 \implies a = -d$; (1.9) implies $c - 2d = 0$.
(1.10) implies $-d + (2d) - d = 0 \implies d = 0 \implies c = 0 \implies a = 0$.

Thus, $w(x, y)$ is identically zero and consequently $(\tau, \tilde{Q}_1, \sigma)$ is unisolvent.



Solution to (b)

We now want to show that $(\tau, \tilde{Q}_1, \sigma)$ leads to a finite element space which is not H^1 -conforming.



¹For more details see Grossmann book pg 238

Solution to (b)

We now want to show that $(\tau, \tilde{Q}_1, \sigma)$ leads to a finite element space which is not H^1 -conforming.

Let us first recall what it means to be **non-conforming**¹:

¹For more details see Grossmann book pg 238



Solution to (b)

We now want to show that $(\tau, \tilde{Q}_1, \sigma)$ leads to a finite element space which is not H^1 -conforming.

Let us first recall what it means to be **non-conforming**¹: Consider the abstract variational problem: Find $u \in V$ such that $a(u, v) = f(v)$ for all $v \in V$. Any finite element method that is not directly based on the discretization of the weak formulation by $a(u_h, v_h) = f(v_h)$ with $u_h, v_h \in V_h \subset V$ is called a nonconforming finite element method.

¹For more details see Grossmann book pg 238



Solution to (b)

We now want to show that $(\tau, \tilde{Q}_1, \sigma)$ leads to a finite element space which is not H^1 -conforming.

Let us first recall what it means to be **non-conforming**¹: Consider the abstract variational problem: Find $u \in V$ such that $a(u, v) = f(v)$ for all $v \in V$. Any finite element method that is not directly based on the discretization of the weak formulation by $a(u_h, v_h) = f(v_h)$ with $u_h, v_h \in V_h \subset V$ is called a nonconforming finite element method.

Thus we want to construct a finite element space for $(\tau, \tilde{Q}_1, \sigma)$ such that $V_h \not\subset H^1$.

¹For more details see Grossmann book pg 238



We show the above with an example.



We show the above with an example. Let K_1 be the square defined by the vertices $\{(-1, -1), (-1, 1), (1, 1), (1, -1)\}$ and K_2 be the square defined by the vertices $\{(-1, -1), (1, -1), (3, 1), (3, -1)\}$.



We show the above with an example. Let K_1 be the square defined by the vertices $\{(-1, -1), (-1, 1), (1, 1), (1, -1)\}$ and K_2 be the square defined by the vertices $\{(-1, -1), (1, -1), (3, 1), (3, -1)\}$. Let us define our domain to be the rectangle $\Omega = K_1 \cup K_2$.



We show the above with an example. Let K_1 be the square defined by the vertices $\{(-1, -1), (-1, 1), (1, 1), (1, -1)\}$ and K_2 be the square defined by the vertices $\{(-1, -1), (1, -1), (3, 1), (3, -1)\}$. Let us define our domain to be the rectangle $\Omega = K_1 \cup K_2$.

Then, the unisolvent finite element leads to the following finite element space:

$$V_h := \{v : \Omega \rightarrow \mathbb{R} \mid v|_{K_i} \in \tilde{Q}_1 \text{ for } i = 1, 2 \text{ and } v|_{K_1}(1, 0) = v|_{K_2}(1, 0)\}. \quad (1.11)$$



We show the above with an example. Let K_1 be the square defined by the vertices $\{(-1, -1), (-1, 1), (1, 1), (1, -1)\}$ and K_2 be the square defined by the vertices $\{(-1, -1), (1, -1), (3, 1), (3, -1)\}$. Let us define our domain to be the rectangle $\Omega = K_1 \cup K_2$.

Then, the unisolvent finite element leads to the following finite element space:

$$V_h := \{v : \Omega \rightarrow \mathbb{R} \mid v|_{K_i} \in \tilde{Q}_1 \text{ for } i = 1, 2 \text{ and } v|_{K_1}(1, 0) = v|_{K_2}(1, 0)\}. \quad (1.11)$$

We claim that $V_h \not\subset H^1(\Omega)$.



Let $w \in V_h$ and let us define $w_1 := w|_{K_1}$ and $w_2 := w|_{K_2}$ such that

$$w_1(x, y) := x^2 - y^2,$$

$$w_2(x, y) := x.$$



²<https://mathworld.wolfram.com/RemovableDiscontinuity.html>

Let $w \in V_h$ and let us define $w_1 := w|_{K_1}$ and $w_2 := w|_{K_2}$ such that

$$w_1(x, y) := x^2 - y^2,$$

$$w_2(x, y) := x.$$

Then notice that $w_1(1, 0) = w_2(1, 0) = 1$,

²<https://mathworld.wolfram.com/RemovableDiscontinuity.html>



Let $w \in V_h$ and let us define $w_1 := w|_{K_1}$ and $w_2 := w|_{K_2}$ such that

$$w_1(x, y) := x^2 - y^2,$$

$$w_2(x, y) := x.$$

Then notice that $w_1(1, 0) = w_2(1, 0) = 1$, but $w_1(1, y) = 1 - y^2 \neq 1 = w_2(1, y)$ where $0 \neq y \in (-1, 1)$.

²<https://mathworld.wolfram.com/RemovableDiscontinuity.html>



Let $w \in V_h$ and let us define $w_1 := w|_{K_1}$ and $w_2 := w|_{K_2}$ such that


$$w_1(x, y) := x^2 - y^2,$$

$$w_2(x, y) := x.$$

Then notice that $w_1(1, 0) = w_2(1, 0) = 1$, but $w_1(1, y) = 1 - y^2 \neq 1 = w_2(1, y)$ where $0 \neq y \in (-1, 1)$.

Therefore, $w \notin H^1(\Omega)$. Note that this discontinuity is not *removable*². Hence, the finite element space is not H^1 -conforming.

²<https://mathworld.wolfram.com/RemovableDiscontinuity.html>



Outline

- 1 January 2010 Exam
 - Problem 2
- 2 August 2009 Exam
 - Problem 1
- 3 January 2010 Problem 3



Problem 1

Consider the following finite element triple:

- K = a rectangle with vertices $\{a^i\}$, $i = 1, 2, 3, 4$.
- $P = Q^3 = \text{span}\{x_1^i x_2^j : i, j = 0, \dots, 3\}$
- $N = \{p(a^i), p_1(a^i), p_2(a^i), p_{12}(a^i), i = 1, 2, 3, 4\}$. (Here p_i denotes differentiation with respect to x_i .)

- (a) Show that the above finite element is unisolvent.
- (b) What do you need to do to check if the above element with a rectangular mesh results in a C^1 finite element space?
- (c) Does the above element (with a rectangular mesh) result in a C^1 finite element space? (Explain your answer).



Solution to (a)

We want to show that the above finite element is unisolvent.



Solution to (a)

We want to show that the above finite element is unisolvent. Recall that to show unisolvence it suffices to show that

$[N \ni \sigma_i(p) = 0, \forall i \in \{1, 2, \dots, 16\}] \implies [p = 0]$ for any $p \in Q^3$. Note that $\text{card}(N) = 16 = \dim P$.



Solution to (a)

We want to show that the above finite element is unisolvent. Recall that to show unisolvence it suffices to show that

$[N \ni \sigma_i(p) = 0, \forall i \in \{1, 2, \dots, 16\}] \implies [p = 0]$ for any $p \in Q^3$. Note that $\text{card}(N) = 16 = \dim P$.

To simplify our computations, let us use the following notation $x := x_1$ and $y := x_2$.



Solution to (a)

We want to show that the above finite element is unisolvent. Recall that to show unisolvence it suffices to show that

$[N \ni \sigma_i(p) = 0, \forall i \in \{1, 2, \dots, 16\}] \implies [p = 0]$ for any $p \in Q^3$. Note that $\text{card}(N) = 16 = \dim P$.

To simplify our computations, let us use the following notation $x := x_1$ and $y := x_2$. In this notation $P = Q^3 = \text{span}\{1, x, x^2, x^3, y, xy, x^2y, x^3y, y^2, xy^2, x^2y^2, x^3y^2, y^3, xy^3, x^2y^3, x^3y^3\}$.



Solution to (a)

We want to show that the above finite element is unisolvent. Recall that to show unisolvence it suffices to show that

$[N \ni \sigma_i(p) = 0, \forall i \in \{1, 2, \dots, 16\}] \implies [p = 0]$ for any $p \in Q^3$. Note that $\text{card}(N) = 16 = \dim P$.

To simplify our computations, let us use the following notation $x := x_1$ and $y := x_2$. In this notation $P = Q^3 = \text{span}\{1, x, x^2, x^3, y, xy, x^2y, x^3y, y^2, xy^2, x^2y^2, x^3y^2, y^3, xy^3, x^2y^3, x^3y^3\}$.

To further simplify our computations, we will show unisolvence on the unit square $\hat{K} = [0, 1] \times [0, 1]$.



Solution to (a)

We want to show that the above finite element is unisolvent. Recall that to show unisolvence it suffices to show that

$[N \ni \sigma_i(p) = 0, \forall i \in \{1, 2, \dots, 16\}] \implies [p = 0]$ for any $p \in Q^3$. Note that $\text{card}(N) = 16 = \dim P$.

To simplify our computations, let us use the following notation $x := x_1$ and $y := x_2$. In this notation $P = Q^3 = \text{span}\{1, x, x^2, x^3, y, xy, x^2y, x^3y, y^2, xy^2, x^2y^2, x^3y^2, y^3, xy^3, x^2y^3, x^3y^3\}$.

To further simplify our computations, we will show unisolvence on the unit square $\hat{K} = [0, 1] \times [0, 1]$. Note that this is justified since we can show **affine equivalence** of the physical and reference finite elements. (



Solution to (a)

We want to show that the above finite element is unisolvent. Recall that to show unisolvence it suffices to show that

$[N \ni \sigma_i(p) = 0, \forall i \in \{1, 2, \dots, 16\}] \implies [p = 0]$ for any $p \in Q^3$. Note that $\text{card}(N) = 16 = \dim P$.

To simplify our computations, let us use the following notation $x := x_1$ and $y := x_2$. In this notation $P = Q^3 = \text{span}\{1, x, x^2, x^3, y, xy, x^2y, x^3y, y^2, xy^2, x^2y^2, x^3y^2, y^3, xy^3, x^2y^3, x^3y^3\}$.

To further simplify our computations, we will show unisolvence on the unit square $\hat{K} = [0, 1] \times [0, 1]$. Note that this is justified since we can show **affine equivalence** of the physical and reference finite elements. (*Do we need to define said mapping? Not sure*)



Definition 3.11 Affine equivalence of finite elements

Let \mathcal{T}_h be a triangulation of $\Omega \subset \mathbb{R}^d$, $K \in \mathcal{T}_h$ and P_K, Σ_K as in Definition 3.9. Let further $(\hat{K}, \hat{P}_{\hat{K}}, \hat{\Sigma}_{\hat{K}})$ be a **reference element**. Then, the finite elements (K, P_K, Σ_K) , $K \in \mathcal{T}_h$, are said to be **affine equivalent** to the reference element $(\hat{K}, \hat{P}_{\hat{K}}, \hat{\Sigma}_{\hat{K}})$, if there exists an invertible affine mapping $F_K : \mathbb{R}^d \rightarrow \mathbb{R}^d$ such that for all $K \in \mathcal{T}_h$

$$(3.15) \quad K = F_K(\hat{K}) ,$$

$$(3.16) \quad P_K = \{p : K \rightarrow \mathbb{R} \mid p = \hat{p} \circ F_K^{-1} , \hat{p} \in \hat{P}_{\hat{K}}\} ,$$

$$(3.17) \quad \Sigma_K = \{\ell_i : P_K \rightarrow \mathbb{R} \mid \ell_i = \hat{\ell}_i \circ F_K^{-1} , \hat{\ell}_i \in \hat{\Sigma}_{\hat{K}} , 1 \leq i \leq n_K\} .$$



Let $f \in \hat{P} = \{\hat{p} : \hat{K} \rightarrow \mathbb{R} \mid \hat{p} = p \circ F_k, p \in P_K\}$.



Let $f \in \hat{P} = \{\hat{p} : \hat{K} \rightarrow \mathbb{R} \mid \hat{p} = p \circ F_k, p \in P_K\}$.

Let us consider f restricted to the edge $e = [0, 1] \times \{y = 0\}$.



Let $f \in \hat{P} = \{\hat{p} : \hat{K} \rightarrow \mathbb{R} \mid \hat{p} = p \circ F_k, p \in P_K\}$.

Let us consider f restricted to the edge $e = [0, 1] \times \{y = 0\}$. With a slight abuse of notation, $f|_e$ is a third degree polynomial of the form $f|_e(x) = a + bx + cx^2 + dx^3$.



Let $f \in \hat{P} = \{\hat{p} : \hat{K} \rightarrow \mathbb{R} \mid \hat{p} = p \circ F_k, p \in P_K\}$.

Let us consider f restricted to the edge $e = [0, 1] \times \{y = 0\}$. With a slight abuse of notation, $f|_e$ is a third degree polynomial of the form $f|_e(x) = a + bx + cx^2 + dx^3$. Note that the conditions $f|_e(0) = 0, f|_e'(0) = 0, f|_e(1) = 0, f|_e'(1) = 0$ imply that $f|_e$ must be identically zero on this edge



Let $f \in \hat{P} = \{\hat{p} : \hat{K} \rightarrow \mathbb{R} \mid \hat{p} = p \circ F_k, p \in P_K\}$.

Let us consider f restricted to the edge $e = [0, 1] \times \{y = 0\}$. With a slight abuse of notation, $f|_e$ is a third degree polynomial of the form $f|_e(x) = a + bx + cx^2 + dx^3$. Note that the conditions $f|_e(0) = 0, f|_e'(0) = 0, f|_e(1) = 0, f|_e'(1) = 0$ imply that $f|_e$ must be identically zero on this edge (**Why?**).



Let $f \in \hat{P} = \{\hat{p} : \hat{K} \rightarrow \mathbb{R} \mid \hat{p} = p \circ F_k, p \in P_K\}$.

Let us consider f restricted to the edge $e = [0, 1] \times \{y = 0\}$. With a slight abuse of notation, $f|_e$ is a third degree polynomial of the form $f|_e(x) = a + bx + cx^2 + dx^3$. Note that the conditions $f|_e(0) = 0, f|_e'(0) = 0, f|_e(1) = 0, f|_e'(1) = 0$ imply that $f|_e$ must be identically zero on this edge (**Why?**). Similarly, we can also show that $f \equiv 0$ restricted to the other three edges.



Let $f \in \hat{P} = \{\hat{p} : \hat{K} \rightarrow \mathbb{R} \mid \hat{p} = p \circ F_k, p \in P_K\}$.

Let us consider f restricted to the edge $e = [0, 1] \times \{y = 0\}$. With a slight abuse of notation, $f|_e$ is a third degree polynomial of the form $f|_e(x) = a + bx + cx^2 + dx^3$. Note that the conditions $f|_e(0) = 0, f|_e'(0) = 0, f|_e(1) = 0, f|_e'(1) = 0$ imply that $f|_e$ must be identically zero on this edge (**Why?**). Similarly, we can also show that $f \equiv 0$ restricted to the other three edges.

This implies that our function $f(x, y)$ can be written as follows:

$$f(x, y) = x(x - 1)y(y - 1)g(x, y)$$



Let $f \in \hat{P} = \{\hat{p} : \hat{K} \rightarrow \mathbb{R} \mid \hat{p} = p \circ F_k, p \in P_K\}$.

Let us consider f restricted to the edge $e = [0, 1] \times \{y = 0\}$. With a slight abuse of notation, $f|_e$ is a third degree polynomial of the form $f|_e(x) = a + bx + cx^2 + dx^3$. Note that the conditions $f|_e(0) = 0, f|_e'(0) = 0, f|_e(1) = 0, f|_e'(1) = 0$ imply that $f|_e$ must be identically zero on this edge (**Why?**). Similarly, we can also show that $f \equiv 0$ restricted to the other three edges.

This implies that our function $f(x, y)$ can be written as follows:

$$f(x, y) = x(x - 1)y(y - 1)g(x, y)$$

Note that $g(x, y)$ is a polynomial with terms that have at most $(xy)^1$, i.e., $g = a + bx + cy + dxy$



Let $f \in \hat{P} = \{\hat{p} : \hat{K} \rightarrow \mathbb{R} \mid \hat{p} = p \circ F_k, p \in P_K\}$.

Let us consider f restricted to the edge $e = [0, 1] \times \{y = 0\}$. With a slight abuse of notation, $f|_e$ is a third degree polynomial of the form $f|_e(x) = a + bx + cx^2 + dx^3$. Note that the conditions $f|_e(0) = 0, f|_e'(0) = 0, f|_e(1) = 0, f|_e'(1) = 0$ imply that $f|_e$ must be identically zero on this edge (**Why?**). Similarly, we can also show that $f \equiv 0$ restricted to the other three edges.

This implies that our function $f(x, y)$ can be written as follows:

$$f(x, y) = x(x - 1)y(y - 1)g(x, y)$$

Note that $g(x, y)$ is a polynomial with terms that have at most $(xy)^1$, i.e., $g = a + bx + cy + dxy$ (**Why?**)



We now want to consider the mixed partial derivative of f and show that this must vanish at each a_i for $i = 1, 2, 3, 4$. A simple computation yields:

$$\begin{aligned} f_{xy} = & (2x - 1)(2y - 1)g + (2x - 1)(y - 1)g_y \\ & + (x^2 - x)(2y - 1)g_x + (x^2 - x)(y^2 - y)g_{xy} \end{aligned}$$



We now want to consider the mixed partial derivative of f and show that this must vanish at each a_i for $i = 1, 2, 3, 4$. A simple computation yields:

$$\begin{aligned} f_{xy} = & (2x - 1)(2y - 1)g + (2x - 1)(y - 1)g_y \\ & + (x^2 - x)(2y - 1)g_x + (x^2 - x)(y^2 - y)g_{xy} \end{aligned}$$

Note that when evaluating this mixed partial derivative at each a_i for $i = 1, 2, 3, 4$,



We now want to consider the mixed partial derivative of f and show that this must vanish at each a_i for $i = 1, 2, 3, 4$. A simple computation yields:

$$\begin{aligned} f_{xy} = & (2x - 1)(2y - 1)g + (2x - 1)(y - 1)g_y \\ & + (x^2 - x)(2y - 1)g_x + (x^2 - x)(y^2 - y)g_{xy} \end{aligned}$$

Note that when evaluating this mixed partial derivative at each a_i for $i = 1, 2, 3, 4$, the last three terms vanish and only the first one remains.



We now want to consider the mixed partial derivative of f and show that this must vanish at each a_i for $i = 1, 2, 3, 4$. A simple computation yields:

$$\begin{aligned} f_{xy} = & (2x - 1)(2y - 1)g + (2x - 1)(y - 1)g_y \\ & + (x^2 - x)(2y - 1)g_x + (x^2 - x)(y^2 - y)g_{xy} \end{aligned}$$

Note that when evaluating this mixed partial derivative at each a_i for $i = 1, 2, 3, 4$, the last three terms vanish and only the first one remains. Therefore, we must have that $g(a_i) = 0$ for each $i = 1, 2, 3, 4$.



We now want to consider the mixed partial derivative of f and show that this must vanish at each a_i for $i = 1, 2, 3, 4$. A simple computation yields:

$$\begin{aligned} f_{xy} = & (2x - 1)(2y - 1)g + (2x - 1)(y - 1)g_y \\ & + (x^2 - x)(2y - 1)g_x + (x^2 - x)(y^2 - y)g_{xy} \end{aligned}$$

Note that when evaluating this mixed partial derivative at each a_i for $i = 1, 2, 3, 4$, the last three terms vanish and only the first one remains. Therefore, we must have that $g(a_i) = 0$ for each $i = 1, 2, 3, 4$. However, since g contains terms with powers of at most $(xy)^1$, this implies that $g \equiv 0$.



We now want to consider the mixed partial derivative of f and show that this must vanish at each a_i for $i = 1, 2, 3, 4$. A simple computation yields:

$$\begin{aligned} f_{xy} = & (2x - 1)(2y - 1)g + (2x - 1)(y - 1)g_y \\ & + (x^2 - x)(2y - 1)g_x + (x^2 - x)(y^2 - y)g_{xy} \end{aligned}$$

Note that when evaluating this mixed partial derivative at each a_i for $i = 1, 2, 3, 4$, the last three terms vanish and only the first one remains. Therefore, we must have that $g(a_i) = 0$ for each $i = 1, 2, 3, 4$. However, since g contains terms with powers of at most $(xy)^1$, this implies that $g \equiv 0$. Thus, our reference finite element is unisolvent and by affine equivalency, so is the given finite element.



Solution to (b)

What do you need to do to check if the above element with a rectangular mesh results in a C^1 finite element space?



Solution to (b)

To check that our finite element space is C^1 globally, we need to check the continuity of our shape functions and their respective derivatives across the edges of adjacent elements.



Solution to (c)

Does the above element (with a rectangular mesh) result in a C^1 finite element space? (Explain your answer).



Solution to (c)

Let us consider two elements K_1 and K_2 of our triangulation such that these elements share an edge e .



Solution to (c)

Let us consider two elements K_1 and K_2 of our triangulation such that these elements share an edge e . More precisely, let e be the line segment formed by the two vertices a_1 and a_2 which belong to both K_1 and K_2 .



Solution to (c)

Let us consider two elements K_1 and K_2 of our triangulation such that these elements share an edge e . More precisely, let e be the line segment formed by the two vertices a_1 and a_2 which belong to both K_1 and K_2 . Specifically, let $a_1 = (x_0, y_1)$ and $a_2 = (x_0, y_2)$ so that e is a vertical line segment.



Solution to (c)

Let us consider two elements K_1 and K_2 of our triangulation such that these elements share an edge e . More precisely, let e be the line segment formed by the two vertices a_1 and a_2 which belong to both K_1 and K_2 . Specifically, let $a_1 = (x_0, y_1)$ and $a_2 = (x_0, y_2)$ so that e is a vertical line segment.

Let $p \in Q^3$ and $q \in Q^3$ be defined on K_1 and K_2 respectively. Let us define:

$$f = p|_e - q|_e$$

$$g = (p_y)|_e - (q_y)|_e$$



Solution to (c)

Let us consider two elements K_1 and K_2 of our triangulation such that these elements share an edge e . More precisely, let e be the line segment formed by the two vertices a_1 and a_2 which belong to both K_1 and K_2 . Specifically, let $a_1 = (x_0, y_1)$ and $a_2 = (x_0, y_2)$ so that e is a vertical line segment.

Let $p \in Q^3$ and $q \in Q^3$ be defined on K_1 and K_2 respectively. Let us define:

$$\begin{aligned}f &= p|_e - q|_e \\g &= (p_y)|_e - (q_y)|_e\end{aligned}$$

We want to show that f and g are identically zero since this will imply that the piecewise defined functions and their derivatives will be continuous across the edge e .



Note that f and g are functions of y only and are both cubic polynomials



Note that f and g are functions of y only and are both cubic polynomials (**why?**).



Note that f and g are functions of y only and are both cubic polynomials (**why?**). Since our finite element is unisolvent, we know that $p(a_1) = q(a_1)$, $p(a_2) = q(a_2)$ and $p_y(a_1) = q_y(a_1)$, $p_y(a_2) = q_y(a_2)$ which implies that $f(a_1) = f(a_2) = f_y(a_1) = f_y(a_2) = 0$.



Note that f and g are functions of y only and are both cubic polynomials (**why?**). Since our finite element is unisolvent, we know that $p(a_1) = q(a_1)$, $p(a_2) = q(a_2)$ and $p_y(a_1) = q_y(a_1)$, $p_y(a_2) = q_y(a_2)$ which implies that $f(a_1) = f(a_2) = f_y(a_1) = f_y(a_2) = 0$. Thus, this implies that f and its derivative evaluated at two points are zero.



Note that f and g are functions of y only and are both cubic polynomials (**why?**). Since our finite element is unisolvent, we know that $p(a_1) = q(a_1)$, $p(a_2) = q(a_2)$ and $p_y(a_1) = q_y(a_1)$, $p_y(a_2) = q_y(a_2)$ which implies that $f(a_1) = f(a_2) = f_y(a_1) = f_y(a_2) = 0$. Thus, this implies that f and its derivative evaluated at two points are zero. Since f was cubic, it must be that $f \equiv 0$.



Note that f and g are functions of y only and are both cubic polynomials (**why?**). Since our finite element is unisolvent, we know that $p(a_1) = q(a_1)$, $p(a_2) = q(a_2)$ and $p_y(a_1) = q_y(a_1)$, $p_y(a_2) = q_y(a_2)$ which implies that $f(a_1) = f(a_2) = f_y(a_1) = f_y(a_2) = 0$. Thus, this implies that f and its derivative evaluated at two points are zero. Since f was cubic, it must be that $f \equiv 0$. A similar idea can be applied to g to show that $g(a_1) = g(a_2) = g_y(a_1) = g_y(a_2) = 0$ which implies $g \equiv 0$. The same argument can be used to the case of a horizontal line segment e . Thus, the above element will result in a C^1 finite element space.



Outline

- 1 January 2010 Exam
 - Problem 2
- 2 August 2009 Exam
 - Problem 1
- 3 January 2010 Problem 3



Problem 3

Consider the following initial boundary value problem: find $u(x, t)$ such that

$$u_t - u_{xx} + u = 0, \quad 0 < x < 1, \quad t > 0 \quad (3.1)$$

$$u_x(0, t) = u_x(1, t) = 0, \quad t > 0 \quad (3.2)$$

$$u(x, 0) = g(x), \quad 0 < x < 1. \quad (3.3)$$

- (a) Derive the semi-discrete approximation of this problem using linear finite elements over a uniform partition of $(0, 1)$. Write it as a system of linear ordinary differential equations for the coefficient vector.
- (b) Further, derive discretizations in time using backward Euler and Crank-Nicolson methods, respectively.
- (c) Show that both fully discrete schemes are unconditionally stable with respect to the initial data in the spatial $L^2(0, 1)$ -norm.



Solution to (a)

Derive the semi-discrete approximation of this problem using linear finite elements over a uniform partition of $(0, 1)$. Write it as a system of linear ordinary differential equations for the coefficient vector.



Solution to (a)

Q: What does it mean to be semi-discrete?



Solution to (a)

Q: What does it mean to be semi-discrete?

Let us proceed “informally”.



Solution to (a)

Q: What does it mean to be semi-discrete?

Let us proceed “informally”. We first multiply the PDE in our IBVP by a sufficiently smooth test function $v(x)$ in some real vector space V , and integrate over the domain $(0, 1)$:

$$0 = \int_0^1 \left(u_t(x, t)v(x) - u_{xx}(x, t)v(x) + u(x, t)v(x) \right) dx,$$



Solution to (a)

Q: What does it mean to be semi-discrete?

Let us proceed “informally”. We first multiply the PDE in our IBVP by a sufficiently smooth test function $v(x)$ in some real vector space V , and integrate over the domain $(0, 1)$:

$$0 = \int_0^1 \left(u_t(x, t)v(x) - u_{xx}(x, t)v(x) + u(x, t)v(x) \right) dx,$$

$$\stackrel{\text{IBP}}{=} (u_t, v) + \int_0^1 \left(u_x(x, t)v_x(x) + u(x, t)v(x) \right) dx - \cancel{[u_x(x, t)v(x)]_0^1}$$



Solution to (a)

Q: What does it mean to be semi-discrete?

Let us proceed “informally”. We first multiply the PDE in our IBVP by a sufficiently smooth test function $v(x)$ in some real vector space V , and integrate over the domain $(0, 1)$:

$$\begin{aligned} 0 &= \int_0^1 \left(u_t(x, t)v(x) - u_{xx}(x, t)v(x) + u(x, t)v(x) \right) dx, \\ &\stackrel{\text{IBP}}{=} (u_t, v) + \int_0^1 \left(u_x(x, t)v_x(x) + u(x, t)v(x) \right) dx - \cancel{[u_x(x, t)v(x)]_0^1} \\ &= (u_t, v) + a(u, v), \end{aligned}$$

where we define the inner product (\cdot, \cdot) and bilinear form $a(\cdot, \cdot)$ to be:

$$\begin{aligned} (u_t, v) &= \int_0^1 u_t v \, dx \\ a(u, v) &= \int_0^1 (u_x v_x + uv) \, dx. \end{aligned}$$



For the integrals above, it is sufficient to take our function space to be $H^1(0, 1)$.



For the integrals above, it is sufficient to take our function space to be $H^1(0, 1)$. Now let V_h be the space of piecewise linear functions defined over the uniform partition of $(0, 1)$.



For the integrals above, it is sufficient to take our function space to be $H^1(0, 1)$. Now let V_h be the space of piecewise linear functions defined over the uniform partition of $(0, 1)$. More precisely,

$$V_h := \{v \in H^1(0, 1) : v|_{[x_i, x_{i+1}]} \in \mathbb{P}_1, i = 0, \dots, N-1\}.$$



For the integrals above, it is sufficient to take our function space to be $H^1(0, 1)$. Now let V_h be the space of piecewise linear functions defined over the uniform partition of $(0, 1)$. More precisely,

$$V_h := \{v \in H^1(0, 1) : v|_{[x_i, x_{i+1}]} \in \mathbb{P}_1, i = 0, \dots, N-1\}.$$

Thus, our weak formulation in the finite element discretization becomes: Find $u_h(t) \in V_h$ such that $(u_{h,t}, v_h) + a(u_h, v_h) = 0$ for all $v_h \in V_h$. Note, $u_h(t) \in V_h$ for each $t > 0$.



Now, let us consider the standard tent functions as our basis for V_h : $\text{span}\{\phi_i\}_{i=0}^N$. We set $v = \phi_j(x)$ and express u_h in terms of the basis functions

$$u_h(x, t) = \sum_{i=0}^N u_i(t) \phi_i(x),$$

where $\{u_i\}_{i=0}^N$ are the unknown coefficients.



Now, let us consider the standard tent functions as our basis for V_h : $\text{span}\{\phi_i\}_{i=0}^N$. We set $v = \phi_j(x)$ and express u_h in terms of the basis functions

$$u_h(x, t) = \sum_{i=0}^N u_i(t) \phi_i(x),$$

where $\{u_i\}_{i=0}^N$ are the unknown coefficients. Substituting these into our weak formulation yields:



Now, let us consider the standard tent functions as our basis for V_h : $\text{span}\{\phi_i\}_{i=0}^N$. We set $v = \phi_j(x)$ and express u_h in terms of the basis functions

$$u_h(x, t) = \sum_{i=0}^N u_i(t) \phi_i(x),$$

where $\{u_i\}_{i=0}^N$ are the unknown coefficients. Substituting these into our weak formulation yields:

$$\int_0^1 \sum_{i=0}^N u_i'(t) \phi_i(x) \phi_j(x) dx + \int_0^1 \left(\sum_{i=0}^N u_i(t) \phi_i'(x) \phi_j'(x) + \sum_{i=0}^N u_i(t) \phi_i(x) \phi_j(x) \right) dx,$$

for $j = 0, \dots, N$.



Using the vector/matrix notation, $\mathbf{U}(t) = [u_i(t)]_{i=1}^N$, $\mathbf{M} = [(\phi_i, \phi_j)]_{i,j=1}^N$,
and $\mathbf{A} = [a(\phi_i, \phi_j)]_{i,j=1}^N$,



Using the vector/matrix notation, $\mathbf{U}(t) = [u_i(t)]_{i=1}^N$, $\mathbf{M} = [(\phi_i, \phi_j)]_{i,j=1}^N$, and $\mathbf{A} = [a(\phi_i, \phi_j)]_{i,j=1}^N$, we can rewrite our semi-discrete variational equation as,

$$\begin{cases} \mathbf{M}\mathbf{U}'(t) + \mathbf{A}\mathbf{U}(t) = \mathbf{0}, \\ \mathbf{U}(0) = \mathbf{G}, \end{cases}$$

where $\mathbf{G} = [g(x_i)]_{i=1}^N$.



Using the vector/matrix notation, $\mathbf{U}(t) = [u_i(t)]_{i=1}^N$, $\mathbf{M} = [(\phi_i, \phi_j)]_{i,j=1}^N$, and $\mathbf{A} = [a(\phi_i, \phi_j)]_{i,j=1}^N$, we can rewrite our semi-discrete variational equation as,

$$\begin{cases} \mathbf{M}\mathbf{U}'(t) + \mathbf{A}\mathbf{U}(t) = \mathbf{0}, \\ \mathbf{U}(0) = \mathbf{G}, \end{cases}$$

where $\mathbf{G} = [g(x_i)]_{i=1}^N$. This is our system of linear ordinary differential equations.

