

07-30-2020

Eric Tovar

Qualifying Prep Course – Numerical

07-30-2020



Outline

- 1 August 2017 Exam
 - Problem 3
- 2 August 2018 Exam
 - Problem 2
- 3 Hermite polynomials
- 4 Elliptic projection



A note on problem 2 boundary conditions

(open PDF)



Problem 3

Let $u(x, t)$ be a smooth solution satisfying

$$\partial_t u + \beta \partial_x u = 0, \quad x \in \Omega := (0, 1), \quad t > 0 \quad \text{and} \quad u(0, x) = \phi(x), \quad x \in \Omega,$$

where $\beta \in \mathbb{R}$ and ϕ is a given smooth function. In addition, we assume that $u(x, t)$ satisfies the periodic boundary condition $u(0, t) = u(1, t), t > 0$. Let $\mathbb{V} = \{v \in H^1(\Omega) : v(0) = v(1)\}$.



(b) Show that the Finite Element approximation $u_h(t)$ satisfies

$$\frac{d}{dt} \sum_{i=0}^N u_h(t, x_i)^2 = 0.$$



(b) Show that the Finite Element approximation $u_h(t)$ satisfies

$$\frac{d}{dt} \sum_{i=0}^N u_h(t, x_i)^2 = 0.$$

(c) Show that

$$c^{-1} \int_{\Omega} u_h^2(t, x) dx \leq h \sum_{i=0}^N u_h(t, x_i)^2 \leq c \int_{\Omega} u_h^2(t, x) dx$$

and deduce the estimate

$$\int_{\Omega} u_h^2(t, x) dx \leq C \int_{\Omega} \phi_h^2(0, x) dx.$$

Here c and C are constants independent of h .



Solution to (b)

First recall that for a smooth solution u , we have $u\partial_t u = \frac{1}{2}\partial_t(u^2)$ and $u\partial_x u = \frac{1}{2}\partial_x(u^2)$.



Solution to (b)

First recall that for a smooth solution u , we have $u\partial_t u = \frac{1}{2}\partial_t(u^2)$ and $u\partial_x u = \frac{1}{2}\partial_x(u^2)$.

Let us take $v_h(x) := u_h(t, x)$ for some fixed t .



Solution to (b)

First recall that for a smooth solution u , we have $u\partial_t u = \frac{1}{2}\partial_t(u^2)$ and $u\partial_x au = \frac{1}{2}\partial_x(u^2)$.

Let us take $v_h(x) := u_h(t, x)$ for some fixed t . Substituting this into the discrete yields the following

$$\frac{h}{2} \sum_{i=0}^N \left(\frac{1}{2} \partial_t (u_h(t, x_{i+1}))^2 + \frac{1}{2} \partial_t (u_h(t, x_i))^2 \right) + \beta \int_{\Omega} \frac{1}{2} \partial_x (u_h(t, x))^2 dx = 0.$$



Solution to (b)

First recall that for a smooth solution u , we have $u\partial_t u = \frac{1}{2}\partial_t(u^2)$ and $u\partial_x au = \frac{1}{2}\partial_x(u^2)$.

Let us take $v_h(x) := u_h(t, x)$ for some fixed t . Substituting this into the discrete yields the following

$$\frac{h}{2} \sum_{i=0}^N \left(\frac{1}{2} \partial_t (u_h(t, x_{i+1}))^2 + \frac{1}{2} \partial_t (u_h(t, x_i))^2 \right) + \beta \int_{\Omega} \frac{1}{2} \partial_x (u_h(t, x))^2 dx = 0.$$

Note that the integral term will vanish because we have periodic boundary conditions.



Solution to (b)

First recall that for a smooth solution u , we have $u\partial_t u = \frac{1}{2}\partial_t(u^2)$ and $u\partial_x a u = \frac{1}{2}\partial_x(u^2)$.

Let us take $v_h(x) := u_h(t, x)$ for some fixed t . Substituting this into the discrete yields the following

$$\frac{h}{2} \sum_{i=0}^N \left(\frac{1}{2} \partial_t (u_h(t, x_{i+1}))^2 + \frac{1}{2} \partial_t (u_h(t, x_i))^2 \right) + \beta \int_{\Omega} \frac{1}{2} \partial_x (u_h(t, x))^2 dx = 0.$$

Note that the integral term will vanish because we have periodic boundary conditions. Then, the remaining summation term can be written as follows

$$\frac{h}{2} \frac{d}{dt} \left[\sum_{i=1}^N u_h(t, x_i)^2 + \frac{1}{2} u_h(t, x_0) + \frac{1}{2} u_h(t, x_{N+1}) \right] = 0$$



Solution to (b)

First recall that for a smooth solution u , we have $u\partial_t u = \frac{1}{2}\partial_t(u^2)$ and $u\partial_x u = \frac{1}{2}\partial_x(u^2)$.

Let us take $v_h(x) := u_h(t, x)$ for some fixed t . Substituting this into the discrete yields the following

$$\frac{h}{2} \sum_{i=0}^N \left(\frac{1}{2} \partial_t (u_h(t, x_{i+1}))^2 + \frac{1}{2} \partial_t (u_h(t, x_i))^2 \right) + \beta \int_{\Omega} \frac{1}{2} \partial_x (u_h(t, x))^2 dx = 0.$$

Note that the integral term will vanish because we have periodic boundary conditions. Then, the remaining summation term can be written as follows

$$\frac{h}{2} \frac{d}{dt} \left[\sum_{i=1}^N u_h(t, x_i)^2 + \frac{1}{2} u_h(t, x_0) + \frac{1}{2} u_h(t, x_{N+1}) \right] = 0$$

But, because we have periodic boundary conditions, we know that $u_h(t, x_{N+1}) = u_h(t, x_0)$. Thus we can combine these terms with the summation:

$$\frac{d}{dt} \sum_{i=0}^N u_h(t, x_i)^2 = 0.$$



Definition A.5 (Equivalent norms). *Two norms $\|\cdot\|_{V,1}$ and $\|\cdot\|_{V,2}$ are said to be equivalent on V if there exists a positive real number c such that*

$$c\|v\|_{V,2} \leq \|v\|_{V,1} \leq c^{-1}\|v\|_{V,2}, \quad \forall v \in V. \quad (\text{A.1})$$

Whenever (A.1) holds true, V is a Banach space for the norm $\|\cdot\|_{V,1}$ if and only if it is a Banach space for the norm $\|\cdot\|_{V,2}$.



We want to show the first string of inequalities in part (c).



We want to show the first string of inequalities in part (c). Let u_h be an element of our finite dimensional space.



We want to show the first string of inequalities in part (c). Let u_h be an element of our finite dimensional space. Let us consider $\|u_h\|_{L^2(\Omega)}^2$:

$$\|u_h\|_{L^2(\Omega)}^2 = \int_{\Omega} u_h^2(t, x) dx$$



We want to show the first string of inequalities in part (c). Let u_h be an element of our finite dimensional space. Let us consider $\|u_h\|_{L^2(\Omega)}^2$:

$$\begin{aligned}\|u_h\|_{L^2(\Omega)}^2 &= \int_{\Omega} u_h^2(t, x) dx \\ &= \sum_{T \in \mathcal{T}_h} \int_T u_h^2(t, x) dx\end{aligned}$$



We want to show the first string of inequalities in part (c). Let u_h be an element of our finite dimensional space. Let us consider $\|u_h\|_{L^2(\Omega)}^2$:

$$\begin{aligned}\|u_h\|_{L^2(\Omega)}^2 &= \int_{\Omega} u_h^2(t, x) dx \\ &= \sum_{T \in \mathcal{T}_h} \int_T u_h^2(t, x) dx \\ &= \sum_{T \in \mathcal{T}_h} \frac{|T|}{|\hat{T}|} \int_{\hat{T}} (\hat{u}_h)^2(t, \hat{x}) d\hat{x}\end{aligned}$$



We want to show the first string of inequalities in part (c). Let u_h be an element of our finite dimensional space. Let us consider $\|u_h\|_{L^2(\Omega)}^2$:

$$\begin{aligned}
 \|u_h\|_{L^2(\Omega)}^2 &= \int_{\Omega} u_h^2(t, x) dx \\
 &= \sum_{T \in \mathcal{T}_h} \int_T u_h^2(t, x) dx \\
 &= \sum_{T \in \mathcal{T}_h} \frac{|T|}{|\hat{T}|} \int_{\hat{T}} (\hat{u}_h)^2(t, \hat{x}) d\hat{x} \\
 &= \sum_{T \in \mathcal{T}_h} \frac{|T|}{|\hat{T}|} \|\hat{u}_h\|_{L^2(\hat{T})}^2,
 \end{aligned}$$

where $\hat{u}_h = u_h \circ F_T$ where $F_T : \hat{T} \rightarrow T$ is the affine map from the reference element to the physical element T .



Note that since our mesh is uniform (can also be just shape-regular),



¹We can only invoke norm equivalence ON THE REFERENCE ELEMENT.

Note that since our mesh is uniform (can also be just shape-regular), we have that $c_1 h \leq |T| \leq c_2 h$. Also note that $\hat{u}_h \in \mathbb{P}_1$ ($\dim \mathbb{P}_1 = 2$).



¹We can only invoke norm equivalence ON THE REFERENCE ELEMENT.

Note that since our mesh is uniform (can also be just shape-regular), we have that $c_1 h \leq |T| \leq c_2 h$. Also note that $\hat{u}_h \in \mathbb{P}_1$ ($\dim \mathbb{P}_1 = 2$). Since we are working on the reference element AND with a finite dimensional space, we can invoke the following norm equivalence¹:

¹We can only invoke norm equivalence ON THE REFERENCE ELEMENT.



Note that since our mesh is uniform (can also be just shape-regular), we have that $c_1 h \leq |T| \leq c_2 h$. Also note that $\hat{u}_h \in \mathbb{P}_1$ ($\dim \mathbb{P}_1 = 2$). Since we are working on the reference element AND with a finite dimensional space, we can invoke the following norm equivalence¹:

$$c' \left(u_h(t, x_i)^2 + u_h(t, x_{i+1})^2 \right) \leq \|\hat{u}_h\|_{L^2(\hat{T})}^2 \leq c \left(u_h(t, x_i)^2 + u_h(t, x_{i+1})^2 \right),$$

¹We can only invoke norm equivalence ON THE REFERENCE ELEMENT.



Note that since our mesh is uniform (can also be just shape-regular), we have that $c_1 h \leq |T| \leq c_2 h$. Also note that $\hat{u}_h \in \mathbb{P}_1$ ($\dim \mathbb{P}_1 = 2$). Since we are working on the reference element AND with a finite dimensional space, we can invoke the following norm equivalence¹:

$$c' \left(u_h(t, x_i)^2 + u_h(t, x_{i+1})^2 \right) \leq \|\hat{u}_h\|_{L^2(\hat{T})}^2 \leq c \left(u_h(t, x_i)^2 + u_h(t, x_{i+1})^2 \right),$$

for positive constants c, c' where we used the norm

$$\|u_h\|_E := \sqrt{u_h(t, x_i)^2 + u_h(t, x_{i+1})^2} \text{ (Verify that this is indeed a norm).}$$

¹We can only invoke norm equivalence ON THE REFERENCE ELEMENT.



Note that since our mesh is uniform (can also be just shape-regular), we have that $c_1 h \leq |T| \leq c_2 h$. Also note that $\hat{u}_h \in \mathbb{P}_1$ ($\dim \mathbb{P}_1 = 2$). Since we are working on the reference element AND with a finite dimensional space, we can invoke the following norm equivalence¹:

$$c' \left(u_h(t, x_i)^2 + u_h(t, x_{i+1})^2 \right) \leq \|\hat{u}_h\|_{L^2(\hat{T})}^2 \leq c \left(u_h(t, x_i)^2 + u_h(t, x_{i+1})^2 \right),$$

for positive constants c, c' where we used the norm

$$\|u_h\|_E := \sqrt{u_h(t, x_i)^2 + u_h(t, x_{i+1})^2} \text{ (Verify that this is indeed a norm).}$$

Note that we have used the fact that we can write $\hat{u}_h(t, 0) = u_h(x_i)$ and $\hat{u}_h(t, 1) = u_h(x_{i+1})$ via the mapping F_T .

¹We can only invoke norm equivalence ON THE REFERENCE ELEMENT.

Then using these inequalities, we have that



Then using these inequalities, we have that

$$\|u_h\|_{L^2(\Omega)}^2 = \sum_{T \in \mathcal{T}_h} \frac{|T|}{|\hat{T}|} \|\hat{u}_h\|_{L^2(\hat{T})}^2$$



Then using these inequalities, we have that

$$\begin{aligned}\|u_h\|_{L^2(\Omega)}^2 &= \sum_{T \in \mathcal{T}_h} \frac{|T|}{|\hat{T}|} \|\hat{u}_h\|_{L^2(\hat{T})}^2 \\ &\leq \sum_{T \in \mathcal{T}_h} \frac{c_2 h}{|\hat{T}|} \|\hat{u}_h\|_{L^2(\hat{T})}^2\end{aligned}$$



Then using these inequalities, we have that

$$\begin{aligned}\|u_h\|_{L^2(\Omega)}^2 &= \sum_{T \in \mathcal{T}_h} \frac{|T|}{|\hat{T}|} \|\hat{u}_h\|_{L^2(\hat{T})}^2 \\ &\leq \sum_{T \in \mathcal{T}_h} \frac{c_2 h}{|\hat{T}|} \|\hat{u}_h\|_{L^2(\hat{T})}^2 \\ &\leq \sum_{T \in \mathcal{T}_h} \frac{c_2 h}{|\hat{T}|} c \left(u_h(t, x_i)^2 + u_h(t, x_{i+1})^2 \right)\end{aligned}$$



Then using these inequalities, we have that

$$\begin{aligned}\|u_h\|_{L^2(\Omega)}^2 &= \sum_{T \in \mathcal{T}_h} \frac{|T|}{|\hat{T}|} \|\hat{u}_h\|_{L^2(\hat{T})}^2 \\ &\leq \sum_{T \in \mathcal{T}_h} \frac{c_2 h}{|\hat{T}|} \|\hat{u}_h\|_{L^2(\hat{T})}^2 \\ &\leq \sum_{T \in \mathcal{T}_h} \frac{c_2 h}{|\hat{T}|} c \left(u_h(t, x_i)^2 + u_h(t, x_{i+1})^2 \right) \\ &\leq 2cc_2 h \sum_{i=1}^N u_h(t, x_i)^2\end{aligned}$$



Then using these inequalities, we have that

$$\begin{aligned}\|u_h\|_{L^2(\Omega)}^2 &= \sum_{T \in \mathcal{T}_h} \frac{|T|}{|\hat{T}|} \|\hat{u}_h\|_{L^2(\hat{T})}^2 \\ &\leq \sum_{T \in \mathcal{T}_h} \frac{c_2 h}{|\hat{T}|} \|\hat{u}_h\|_{L^2(\hat{T})}^2 \\ &\leq \sum_{T \in \mathcal{T}_h} \frac{c_2 h}{|\hat{T}|} c \left(u_h(t, x_i)^2 + u_h(t, x_{i+1})^2 \right) \\ &\leq 2cc_2 h \sum_{i=1}^N u_h(t, x_i)^2\end{aligned}$$

The other direction can be shown by using the other side of the inequalities.



We now want to prove the last inequality of part (c).



We now want to prove the last inequality of part (c). First, note that by part (b), the sum $\sum_{i=1}^N u_h(t, x_i)^2$ is constant in time so we can say that $\sum_{i=1}^N u_h(t, x_i)^2 = \sum_{i=1}^N u_h(0, x_i)^2 = \sum_{i=1}^N \phi_h(x_i)^2$.



We now want to prove the last inequality of part (c). First, note that by part (b), the sum $\sum_{i=1}^N u_h(t, x_i)^2$ is constant in time so we can say that $\sum_{i=1}^N u_h(t, x_i)^2 = \sum_{i=1}^N u_h(0, x_i)^2 = \sum_{i=1}^N \phi_h(x_i)^2$. Then, using the inequality we just proved, we see that:

$$c^{-1} \int_{\Omega} u_h^2(t, x) dx \leq h \sum_{i=1}^N u_h(t, x_i)^2$$



We now want to prove the last inequality of part (c). First, note that by part (b), the sum $\sum_{i=1}^N u_h(t, x_i)^2$ is constant in time so we can say that $\sum_{i=1}^N u_h(t, x_i)^2 = \sum_{i=1}^N u_h(0, x_i)^2 = \sum_{i=1}^N \phi_h(x_i)^2$. Then, using the inequality we just proved, we see that:

$$\begin{aligned} c^{-1} \int_{\Omega} u_h^2(t, x) dx &\leq h \sum_{i=1}^N u_h(t, x_i)^2 \\ &= h \sum_{i=1}^N u_h(0, x_i)^2 \end{aligned}$$



We now want to prove the last inequality of part (c). First, note that by part (b), the sum $\sum_{i=1}^N u_h(t, x_i)^2$ is constant in time so we can say that $\sum_{i=1}^N u_h(t, x_i)^2 = \sum_{i=1}^N u_h(0, x_i)^2 = \sum_{i=1}^N \phi_h(x_i)^2$. Then, using the inequality we just proved, we see that:

$$\begin{aligned} c^{-1} \int_{\Omega} u_h^2(t, x) dx &\leq h \sum_{i=1}^N u_h(t, x_i)^2 \\ &= h \sum_{i=1}^N u_h(0, x_i)^2 \\ &= h \sum_{i=1}^N \phi_h(x_i)^2 \end{aligned}$$



We now want to prove the last inequality of part (c). First, note that by part (b), the sum $\sum_{i=1}^N u_h(t, x_i)^2$ is constant in time so we can say that $\sum_{i=1}^N u_h(t, x_i)^2 = \sum_{i=1}^N u_h(0, x_i)^2 = \sum_{i=1}^N \phi_h(x_i)^2$. Then, using the inequality we just proved, we see that:

$$\begin{aligned}
 c^{-1} \int_{\Omega} u_h^2(t, x) dx &\leq h \sum_{i=1}^N u_h(t, x_i)^2 \\
 &= h \sum_{i=1}^N u_h(0, x_i)^2 \\
 &= h \sum_{i=1}^N \phi_h(x_i)^2 \\
 &\leq c \int_{\Omega} \phi_h^2(x) dx
 \end{aligned}$$



Outline

- 1 August 2017 Exam
 - Problem 3
- 2 August 2018 Exam
 - Problem 2
- 3 Hermite polynomials
- 4 Elliptic projection



Problem 2

Let T be the unit triangle in \mathbb{R}^2 , with vertices $v_1 = (0, 0)$, $v_2 = (1, 0)$, and $v_3 = (0, 1)$ and edges $e_1 = v_1 v_2$, $e_2 = v_2 v_3$, and $e_3 = v_3 v_1$. Let $RT_0 = \{(a + cx, b + cy) : a, b, c \in \mathbb{R}\}$ (so that members of RT_0 are vector functions over T , and $[\mathbb{P}_0]^2 \subsetneq RT_0 \subsetneq [\mathbb{P}_1]^2$). Finally, let $\sigma_i(\vec{u}) = \int_{e_i} \vec{u} \cdot \vec{n}_i$ where \vec{n}_i is the outward pointing unit normal vector to T on e_i , and let $\Sigma = \{\sigma_1, \sigma_2, \sigma_3\}$.



Problem 2

Let T be the unit triangle in \mathbb{R}^2 , with vertices $v_1 = (0, 0)$, $v_2 = (1, 0)$, and $v_3 = (0, 1)$ and edges $e_1 = v_1 v_2$, $e_2 = v_2 v_3$, and $e_3 = v_3 v_1$. Let $RT_0 = \{(a + cx, b + cy) : a, b, c \in \mathbb{R}\}$ (so that members of RT_0 are vector functions over T , and $[\mathbb{P}_0]^2 \subsetneq RT_0 \subsetneq [\mathbb{P}_1]^2$). Finally, let $\sigma_i(\vec{u}) = \int_{e_i} \vec{u} \cdot \vec{n}_i$ where \vec{n}_i is the outward pointing unit normal vector to T on e_i , and let $\Sigma = \{\sigma_1, \sigma_2, \sigma_3\}$.

(a) Show that (T, RT_0, Σ) is unisolvent.



Problem 2

Let T be the unit triangle in \mathbb{R}^2 , with vertices $v_1 = (0, 0)$, $v_2 = (1, 0)$, and $v_3 = (0, 1)$ and edges $e_1 = v_1 v_2$, $e_2 = v_2 v_3$, and $e_3 = v_3 v_1$. Let $RT_0 = \{(a + cx, b + cy) : a, b, c \in \mathbb{R}\}$ (so that members of RT_0 are vector functions over T , and $[\mathbb{P}_0]^2 \subsetneq RT_0 \subsetneq [\mathbb{P}_1]^2$). Finally, let $\sigma_i(\vec{u}) = \int_{e_i} \vec{u} \cdot \vec{n}_i$ where \vec{n}_i is the outward pointing unit normal vector to T on e_i , and let $\Sigma = \{\sigma_1, \sigma_2, \sigma_3\}$.

- (a) Show that (T, RT_0, Σ) is unisolvent.
- (b) Find a basis $\{\vec{\varphi}_1, \vec{\varphi}_2, \vec{\varphi}_3\}$ for RT_0 that is dual to Σ , i.e. $\sigma_i(\vec{\varphi}_j) = \delta_{ij}$ with $\delta_{ij} = 1$ if $i = j$ and $\delta_{ij} = 0$ otherwise.



Solution to (a)

We want to show that the finite element (T, RT_0, Σ) is unisolvent.



Solution to (a)

We want to show that the finite element (T, RT_0, Σ) is unisolvent. To simplify our computations, we will show that the reference finite element is unisolvent.




Solution to (a)

We want to show that the finite element (T, RT_0, Σ) is unisolvent. To simplify our computations, we will show that the reference finite element is unisolvent. Note that this is justified by the affine equivalence of the physical and reference element.




Solution to (a)

We want to show that the finite element (T, RT_0, Σ) is unisolvent. To simplify our computations, we will show that the reference finite element is unisolvent. Note that this is justified by the affine equivalence of the physical and reference element.



Solution to (a)

We want to show that the finite element (T, RT_0, Σ) is unisolvent. To simplify our computations, we will show that the reference finite element is unisolvent. Note that this is justified by the affine equivalence of the physical and reference element.




First note that $\text{card}(\Sigma) = 3 = \dim RT_0$.



Solution to (a)

We want to show that the finite element (T, RT_0, Σ) is unisolvent. To simplify our computations, we will show that the reference finite element is unisolvent. Note that this is justified by the affine equivalence of the physical and reference element.



First note that $\text{card}(\Sigma) = 3 = \dim RT_0$. We now want to verify that $\sigma(\vec{p}) = 0$ for all $\sigma \in \Sigma$ implies that $\vec{p} = 0$ for any $\vec{p} \in RT_0$.



Let $\vec{p} \in RT_0$ be arbitrary.



Let $\vec{p} \in RT_0$ be arbitrary. Assume that $\sigma_i(\vec{p}) = 0$ for $i = 1, 2, 3$.



Let $\vec{p} \in RT_0$ be arbitrary. Assume that $\sigma_i(\vec{p}) = 0$ for $i = 1, 2, 3$. Then we have,

$$\sigma_1(\vec{p}) = \int_{e_1} (a + cx, b + cy) \cdot (0, -1) \, ds$$



Let $\vec{p} \in RT_0$ be arbitrary. Assume that $\sigma_i(\vec{p}) = 0$ for $i = 1, 2, 3$. Then we have,

$$\begin{aligned}\sigma_1(\vec{p}) &= \int_{e_1} (a + cx, b + cy) \cdot (0, -1) \, ds \\ &= \int_0^1 -b \, dx\end{aligned}$$



Let $\vec{p} \in RT_0$ be arbitrary. Assume that $\sigma_i(\vec{p}) = 0$ for $i = 1, 2, 3$. Then we have,

$$\begin{aligned}\sigma_1(\vec{p}) &= \int_{e_1} (a + cx, b + cy) \cdot (0, -1) \, ds \\ &= \int_0^1 -b \, dx \\ &= 0.\end{aligned}$$

This implies that $b = 0$.



Let $\vec{p} \in RT_0$ be arbitrary. Assume that $\sigma_i(\vec{p}) = 0$ for $i = 1, 2, 3$. Then we have,

$$\begin{aligned}\sigma_1(\vec{p}) &= \int_{e_1} (a + cx, b + cy) \cdot (0, -1) \, ds \\ &= \int_0^1 -b \, dx \\ &= 0.\end{aligned}$$

This implies that $b = 0$. Next, we have,

$$\sigma_3(\vec{p}) = \int_{e_3} (a + cx, cy) \cdot (-1, 0) \, ds$$



Let $\vec{p} \in RT_0$ be arbitrary. Assume that $\sigma_i(\vec{p}) = 0$ for $i = 1, 2, 3$. Then we have,

$$\begin{aligned}\sigma_1(\vec{p}) &= \int_{e_1} (a + cx, b + cy) \cdot (0, -1) \, ds \\ &= \int_0^1 -b \, dx \\ &= 0.\end{aligned}$$

This implies that $b = 0$. Next, we have,

$$\begin{aligned}\sigma_3(\vec{p}) &= \int_{e_3} (a + cx, cy) \cdot (-1, 0) \, ds \\ &= \int_0^1 -a \, dy\end{aligned}$$



Let $\vec{p} \in RT_0$ be arbitrary. Assume that $\sigma_i(\vec{p}) = 0$ for $i = 1, 2, 3$. Then we have,

$$\begin{aligned}\sigma_1(\vec{p}) &= \int_{e_1} (a + cx, b + cy) \cdot (0, -1) \, ds \\ &= \int_0^1 -b \, dx \\ &= 0.\end{aligned}$$

This implies that $b = 0$. Next, we have,

$$\begin{aligned}\sigma_3(\vec{p}) &= \int_{e_3} (a + cx, cy) \cdot (-1, 0) \, ds \\ &= \int_0^1 -a \, dy \\ &= 0.\end{aligned}$$

Hence $a = 0$.



Finally for the last edge, we have,

$$\sigma_2(\vec{p}) = \int_{e_2} (cx, cy) \cdot (1/\sqrt{2}, 1/\sqrt{2}) ds$$



Finally for the last edge, we have,

$$\begin{aligned}\sigma_2(\vec{p}) &= \int_{e_2} (cx, cy) \cdot (1/\sqrt{2}, 1/\sqrt{2}) \, ds \\ &= \int_0^1 c(1-t, t) \cdot (1/\sqrt{2}, 1/\sqrt{2}) \underbrace{\sqrt{2}} \, dt\end{aligned}$$



Finally for the last edge, we have,

$$\begin{aligned}\sigma_2(\vec{p}) &= \int_{e_2} (cx, cy) \cdot (1/\sqrt{2}, 1/\sqrt{2}) \, ds \\ &= \int_0^1 c(1-t, t) \cdot (1/\sqrt{2}, 1/\sqrt{2}) \underbrace{\sqrt{2}} \, dt \\ &= \int_0^1 c \, dt\end{aligned}$$



Finally for the last edge, we have,

$$\begin{aligned}\sigma_2(\vec{p}) &= \int_{e_2} (cx, cy) \cdot (1/\sqrt{2}, 1/\sqrt{2}) \, ds \\ &= \int_0^1 c(1-t, t) \cdot (1/\sqrt{2}, 1/\sqrt{2}) \underbrace{\sqrt{2}} \, dt \\ &= \int_0^1 c \, dt \\ &= 0,\end{aligned}$$

where we have parameterized the line segment e_2 by $\gamma(t) := (1-t, t)$ for $t \in [0, 1]$.



Finally for the last edge, we have,

$$\begin{aligned}\sigma_2(\vec{p}) &= \int_{e_2} (cx, cy) \cdot (1/\sqrt{2}, 1/\sqrt{2}) \, ds \\ &= \int_0^1 c(1-t, t) \cdot (1/\sqrt{2}, 1/\sqrt{2}) \underbrace{\sqrt{2}} \, dt \\ &= \int_0^1 c \, dt \\ &= 0,\end{aligned}$$

where we have parameterized the line segment e_2 by $\gamma(t) := (1-t, t)$ for $t \in [0, 1]$. Thus, we have $c = 0$ and therefore $\vec{p} = \vec{0}$. We have unisolvence on the reference element and thus (T, RT_0, Σ) is unisolvent.



Solution to (b)

We now want to find three elements $\vec{\varphi}_i$ of RT_0 such that $\sigma_i(\vec{\varphi}_j) = \delta_{i,j}$.



Solution to (b)

We now want to find three elements $\vec{\varphi}_i$ of RT_0 such that $\sigma_i(\vec{\varphi}_j) = \delta_{i,j}$. Following the work from part (a)



Solution to (b)

We now want to find three elements $\vec{\varphi}_i$ of RT_0 such that $\sigma_i(\vec{\varphi}_j) = \delta_{i,j}$. Following the work from part (a), we see that if $\sigma_1(\vec{\varphi}_1) = 1$ and $\sigma_2(\vec{\varphi}_1) = \sigma_3(\vec{\varphi}_1) = 0$, we will have that $b = -1$ and $a = 0$.



Solution to (b)

We now want to find three elements $\vec{\varphi}_i$ of RT_0 such that $\sigma_i(\vec{\varphi}_j) = \delta_{i,j}$. Following the work from part (a), we see that if $\sigma_1(\vec{\varphi}_1) = 1$ and $\sigma_2(\vec{\varphi}_1) = \sigma_3(\vec{\varphi}_1) = 0$, we will have that $b = -1$ and $a = 0$. Finding c requires a little bit more work:

$$\begin{aligned}\sigma_3(\vec{\varphi}_1) &= \int_{e_3} (cx, -1 + cy) \cdot (1/\sqrt{2}, 1/\sqrt{2}) ds \\ &= \int_0^1 (c(1-t), -1 + ct) \cdot (1, 1) dt \\ &= \int_0^1 c - 1 dt \\ &= 0.\end{aligned}$$

This implies that $c = 1$.



Solution to (b)

We now want to find three elements $\vec{\varphi}_i$ of RT_0 such that $\sigma_i(\vec{\varphi}_j) = \delta_{i,j}$. Following the work from part (a), we see that if $\sigma_1(\vec{\varphi}_1) = 1$ and $\sigma_2(\vec{\varphi}_1) = \sigma_3(\vec{\varphi}_1) = 0$, we will have that $b = -1$ and $a = 0$. Finding c requires a little bit more work:

$$\begin{aligned}\sigma_3(\vec{\varphi}_1) &= \int_{e_3} (cx, -1 + cy) \cdot (1/\sqrt{2}, 1/\sqrt{2}) ds \\ &= \int_0^1 (c(1-t), -1 + ct) \cdot (1, 1) dt \\ &= \int_0^1 c - 1 dt \\ &= 0.\end{aligned}$$

This implies that $c = 1$. Thus $\vec{\varphi}_1 = (x, y - 1)$.



We can repeat the same process for the other basis functions. Doing so, we find,

$$\vec{\varphi}_1(x, y) = (x, y - 1)$$

$$\vec{\varphi}_2(x, y) = (x, y)$$

$$\vec{\varphi}_3(x, y) = (x - 1, y).$$



Outline

- 1 August 2017 Exam
 - Problem 3
- 2 August 2018 Exam
 - Problem 2
- 3 Hermite polynomials**
- 4 Elliptic projection



Hermite polynomials

Consider the finite elements $(K_i, \mathbb{P}^3, \Sigma_i)$ where $K_i = [x_{i-1}, x_i]$ for $i = 1, \dots, N$ and $x_i - x_{i-1} = h$ and \mathbb{P}^3 is the space of cubic polynomials. The degrees of freedom are defined to be $\Sigma_i = \{\sigma_{i-1}, \sigma'_{i-1}, \sigma_i, \sigma'_i\}$, for $i = 1, \dots, N$ where

$$\sigma_{i-1}(f) = f(x_{i-1})$$

$$\sigma'_{i-1}(f) = f'(x_{i-1})$$

$$\sigma_i(f) = f(x_i)$$

$$\sigma'_i(f) = f'(x_i).$$

Note that one can show that the finite element is unisolvent. For some arbitrary finite element approximation, consider the finite dimensional space

$$V_h = \{v \in C^1(\Omega) : v|_{K_i} \in \mathbb{P}^3, i = 1, \dots, N, v(0) = v(1) = 0\}.$$



A basis for this space is given by $\cup_{i=1}^{N-1} \{\phi_i\} \cup \cup_{i=0}^N \{\psi_i\}$, where ϕ_i and ψ_i are the cubic Hermite polynomials (note that we remove ϕ_0 and ϕ_N because of the boundary conditions). Specifically, ϕ_i and ψ_i are defined by the following conditions,

$$\begin{aligned} \sigma_k(\psi_j) &= 0, \quad \sigma'_k(\psi_j) = \delta_{kj}, & \text{for } j, k = 0, \dots, N \\ \sigma_k(\phi_j) &= \delta_{kj}, \quad \sigma'_k(\phi_j) = 0, & \text{for } j, k = 1, \dots, N-1 \end{aligned}$$

Precisely speaking, ψ_i and ϕ_i are defined as

$$\begin{aligned} \psi_i(x) &= \begin{cases} \frac{1}{h^2}(x - x_i)(x - x_{i-1})^2 & \text{for } x \in [x_{i-1}, x_i], \\ \frac{1}{h^2}(x - x_{i+1})^2(x - x_i) & \text{for } x \in [x_i, x_{i+1}], \\ 0 & \text{otherwise,} \end{cases} \\ \phi_i(x) &= \begin{cases} \frac{1}{h^2}(x - x_{i-1})^2\left(\frac{2}{h}(x_i - x) + 1\right) & \text{for } x \in [x_{i-1}, x_i], \\ \frac{1}{h^2}(x_{i+1} - x)^2\left(\frac{2}{h}(x - x_i) + 1\right) & \text{for } x \in [x_i, x_{i+1}], \\ 0 & \text{otherwise.} \end{cases} \end{aligned}$$



Recall that Hermite cubic polynomials are useful when solving fourth-order boundary value problems. See: **January 2009, Problem 1, January 2017, Problem 2, January 2020, Problem 2**



Exercise 1: Let $K := [0, 1]$, $P := \mathbb{P}_3$, and $\Sigma := \{\sigma_1, \sigma_2, \sigma_3, \sigma_4\}$ be the linear forms on P such that

$\sigma_1(p) := p(0)$, $\sigma_2 := p'(0)$, $\sigma_3(p) := p(1)$, $\sigma_4(p) := p'(1)$ for all $p \in P$.

- (a) Show that (K, P, Σ) is a finite element.
- (b) Compute the shape functions for this finite element.
- (c) Indicate possible choices for the domain $V(K)$ of the canonical interpolation operator.

Exercise 2: Assume we have a conforming finite dimensional space $V_h \subset V$ and everything is “well-defined”. Here V_h is the space defined two slides above. Let $\pi_h : V \rightarrow V_h$ be the canonical interpolation operator. What is:

$$\|u - \pi_h u\|_{L^2(\Omega)} \leq ?$$

$$\|u - \pi_h u\|_{H^1(\Omega)} \leq ?$$

$$\|u - \pi_h u\|_{H^2(\Omega)} \leq ?$$

$$\|u - \pi_h u\|_{H^3(\Omega)} \leq ?$$



Outline

- 1 August 2017 Exam
 - Problem 3
- 2 August 2018 Exam
 - Problem 2
- 3 Hermite polynomials
- 4 Elliptic projection



Elliptic projection

