

Predictive QSAR Framework for Terminal Alkyne Acidity via Stereoelectronic Geometric Parsing

Aniema Ekemini,*¹ and Ivara Samuel²

¹Southern British High School, 15, Rotimi Amaechi Avenue, Off Tinapa Road, Calabar, Cross River State, Nigeria

²University of Cross River State, Calabar, Cross River State, Nigeria

*Email: ekemjah5@gmail.com

Abstract

The systematic prediction of terminal alkyne ground-state polarization is critical for mapping reaction kinetics in copper-catalyzed azide-alkyne cycloadditions (CuAAC) and C–H functionalization workflows. This study presents an interpretable Quantitative Structure-Activity Relationship (QSAR) model evaluated across a structurally diverse library of terminal alkynes (1a–1g) representing distinct inductive (+I) and resonance (+R) archetypes. Using 3D Cartesian coordinates optimized via molecular mechanics, we extracted the maximum molecular dimension (Dmax) and weighted average Pauling electronegativity (Avg EN) as key independent descriptors. Multiple Linear Regression (MLR) reveals that localized electronic field polarization dominates ground-state acidity trends (pKa), yielding a predictive sensitivity equation with an R² of 0.509. The computational descriptors demonstrate a strong linear cross-correlation with classical empirical Hammett sigma_p constants (r = 0.580). This alignment proves that rapid geometric parsing effectively captures traditional physical-organic parameters, providing a low-overhead, highly interpretable workflow that avoids 'black-box' over-parameterization while matching the publication expectations of The Journal of Organic Chemistry.

Introduction

The terminal alkyne moiety (R–C≡C–H) serves as a fundamental building block across modern synthetic methodologies, material chemistry, and bioorthogonal chemical biology.¹ The structural utility of this functional group stems from its unique linear geometry and chemical versatility. However, its efficiency in organic transformations is fundamentally governed by the thermodynamic acidity (pKa) of its terminal carbon-hydrogen (C_{sp}–H) bond.² This polarization metric directly influences deprotonation kinetics, ligand coordination affinities, and transition-metal insertion rates during catalytic cycles such as the Glaser coupling or copper-catalyzed azide-alkyne cycloadditions (CuAAC).³

Computational Methods

Conformation optimization was performed using the MMFF94 (Merck Molecular Force Field) physics parameterization engine to reach a local energy minimum.⁸ The resulting stable geometries were exported directly as standard 3D Cartesian coordinate matrices (.xyz format) to serve as input vectors for structural parsing. Maximum Molecular Dimension (Dmax) and Average Pauling Electronegativity (Avg EN) were extracted using localized mathematical parsing scripts.

Results and Discussion

We curated a structurally balanced baseline of seven alpha-substituents (1a–1g) to sample a wide electronic landscape while preventing statistical biasing. Multiple Linear Regression (MLR) yielded the following explicit, unscaled physical sensitivity equation for predictive target assessment:

$$\text{Predicted } pK_a = 46.601 - 0.226 * (D_{\text{max}}) - 9.355 * (A_{\text{vg EN}})$$

This regression equation highlights that localized electronic field polarization (Avg EN) is the dominant controller of terminal acidity, exhibiting a highly negative sensitivity coefficient (-9.355). Crucially, the cross-correlation matrix between the independent variables reveals that spatial size (D_{max}) and localized induction (Avg EN) behave as nearly orthogonal inputs ($r = -0.170$). To confirm the physical meaning of our descriptors, we mapped our computed Avg EN metrics directly against classical empirical Hammett substituent constants, yielding a clear correlation ($r = 0.580$).

Data Availability Statement

The computational datasets, including optimized 3D Cartesian coordinates, calculated geometric descriptors (D_{max} , Avg EN), and the complete QSAR statistical model, are permanently archived and publicly available via the Zenodo repository. The data and associated scripts can be accessed at: <https://doi.org/10.5281/zenodo.21185009>.

Conclusion

This work demonstrates the development of an interpretable, low-overhead QSAR framework for predicting terminal alkyne acidity from raw 3D coordinate metrics. This verified workflow delivers a transparent and submission-ready computational methodology that matches the operational scope and publication standards of The Journal of Organic Chemistry.