

Quantum Expanders and Geometry of Operator Spaces

by

Gilles Pisier*

Texas A&M University
College Station, TX 77843, U. S. A.

and

Université Paris VI
Inst. Math. Jussieu, Equipe d'Analyse, Case 186, 75252
Paris Cedex 05, France

April 18, 2019

Abstract

We show that there are well separated families of quantum expanders with asymptotically the maximal cardinality allowed by a known upper bound. This has applications to the “growth” of certain operator spaces: It implies asymptotically sharp estimates for the growth of the multiplicity of M_N -spaces needed to represent (up to a constant $C > 1$) the M_N -version of the n -dimensional operator Hilbert space OH_n as a direct sum of copies of M_N . We show that, when C is close to 1, this multiplicity grows as $\exp \beta n N^2$ for some constant $\beta > 0$. The main idea is to relate quantum expanders with “smooth” points on the matricial analogue of the Euclidean unit sphere. This generalizes to operator spaces a classical geometric result on n -dimensional Hilbert space (corresponding to $N=1$). In an appendix, we give a quick proof of an inequality (related to Hastings’s previous work) on random unitary matrices that is crucial for this paper.

The term “Quantum Expander” is used by Hastings in [11] and by Ben-Aroya and Ta-Shma in [2] to designate a sequence $\{U^{(N)} \mid N \geq 1\}$ of n -tuples $U^{(N)} = (U_1^{(N)}, \dots, U_n^{(N)})$ of $N \times N$ unitary matrices such that there is an $\varepsilon > 0$ satisfying the following “spectral gap” condition:

$$(0.1) \quad \forall N \forall x \in M_N \quad \left\| \sum_1^n U_j^{(N)}(x - N^{-1} \text{tr}(x)) U_j^{(N)*} \right\|_2 \leq n(1 - \varepsilon) \|x - N^{-1} \text{tr}(x)\|_2,$$

where $\|\cdot\|_2$ denotes the Hilbert-Schmidt norm on M_N . More generally, the term is extended to the case when this is only defined for infinitely many N ’s, and also to n -tuples of matrices satisfying merely $\sum U_j^{(N)} U_j^{(N)*} = \sum U_j^{(N)*} U_j^{(N)} = nI$.

We will say that an n -tuple $U^{(N)}$ satisfying (0.1) is a ε -quantum expander. We refer the reader to the survey [3] for more information and references on quantum expanders.

In analogy with the classical expanders (see below), one seeks to exhibit (and hopefully to construct explicitly) sequences $\{U^{(N_m)} \mid m \geq 1\}$ of n -tuples of $N_m \times N_m$ unitary matrices that are ε -quantum expanders with $N_m \rightarrow \infty$ while n and $\varepsilon > 0$ remain fixed.

When G is a finite group generated by $S = \{t_1, \dots, t_n\}$ the associated Cayley graph $\mathcal{G}(G, S)$ is said to have a spectral gap if the left regular representation λ_G satisfies

$$(0.2) \quad \left\| \sum \lambda_G(t_j)_{|\mathbb{I}^\perp} \right\| < n(1 - \varepsilon)$$

*Partially supported by ANR-2011-BS01-008-01.

where \mathbb{I} denotes the constant function 1 on G . Obviously, this is equivalent to the condition that the unitaries $U_j = \lambda_G(t_j)$ satisfy (0.1) when restricted to diagonal matrices x (here $N = |G|$). In this light, quantum expanders appear as a non-commutative version of the classical ones. More precisely, (0.2) holds iff the unitaries $U_j = \lambda_G(t_j)$ satisfy (0.1) for all x in the orthogonal complement of right translation operators. This is easy to deduce from the decomposition into irreducibles of $\lambda_G \otimes \bar{\lambda}_G$, in which the component of the trivial representation corresponds to the restriction to right translation operators.

In addition, for any irreducible representation π of G , (0.2) implies that the unitaries $(\pi(t_j))$ satisfy (0.1) because the non trivial irreducible components of the representation $\pi \otimes \bar{\pi}$ are contained in λ_G . See Remark 1.6 for more on this.

A sequence of Cayley graphs $\mathcal{G}(G^{(m)}, S^{(m)})$ constitutes an expander in the usual sense if (0.2) is satisfied with $\varepsilon > 0$ and n fixed while $|G^{(m)}| \rightarrow \infty$.

Expanders (equivalently expanding graphs) have been extremely useful, especially (in the applied direction) since Margulis and Lubotzky-Phillips-Sarnak obtained explicit constructions (as opposed to random ones). We refer to [17, 12] for more information and references.

They have also been used with great success for operator algebras and in operator theory (see e.g. [35, 6, 13] see also [27, 5]). In [13], is crucially used the fact that when the dimensions N, N' are suitably different, say if N is much larger than N' , and $U^{(N)}$ satisfies (0.1) then $U^{(N)}$ and $U^{(N')}$ are separated in the sense that there is a fixed $\delta = \delta(\varepsilon) > 0$ such that $\forall x \in M_{N \times N'} \quad \|\sum U_j^{(N)} x U_j^{(N')*}\|_2 \leq n(1 - \delta)\|x\|_2$ (see Remark 1.13 for more on this).

Motivated by operator theory considerations, it is natural to wonder what happens when $N = N'$. We will say that two n -tuples $u = (u_j)$ and $v = (v_j)$ of $N \times N$ unitary matrices are δ -separated if

$$\forall x \in M_N \quad \|\sum_1^n u_j x v_j^*\|_2 \leq n(1 - \delta)\|x\|_2.$$

Equivalently this means that

$$\|\sum_1^n u_j \otimes \bar{v}_j\| \leq n(1 - \delta)$$

where \bar{v}_j denotes the complex conjugate of the matrix v_j , and the norm is the operator norm on $\ell_2^N \otimes \overline{\ell_2^N}$. This can be interpreted in operator space theory as a rough sort of orthogonality related to the “operator space Hilbert space OH”.

Note for example that when (0.2) holds then, for any pair of inequivalent irreducible representations π, σ on G , the n -tuples $(\pi(t_j))$ and $(\sigma(t_j))$ are ε -separated.

Let $U(N) \subset M_N$ denote the group of unitary matrices. The main result of §1 asserts that for any $0 < \delta < 1$ there is a constant $\beta = \beta_\delta > 0$ such that for each $0 < \varepsilon < 1$, for all sufficiently large integer n (i.e. $n \geq n_0(\varepsilon, \delta)$), for any integer N there is a δ -separated family $\{u(t) \mid t \in T\} \subset U(N)^n$ of ε -quantum expanders such that

$$|T| \geq \exp \beta n N^2.$$

Thus we can “pack” as many as $m = \exp \beta n N^2$ δ -separated ε -quantum expanders inside $U(N)^n$. This number m is remarkably large. In fact, in some sense it is as large as can be. Indeed, it is known ([37, 10], see also Remark 1.5) that the maximal m is at most $\exp \beta' n N^2$ for some constant β' .

In §2, we use quantum expanders to investigate the analogue for operator spaces of a well known geometric property of Euclidean space: The unit sphere in a Hilbert space is smooth. Equivalently all its points admit a unique norming functional. In our extension of this, “norming” will be with respect to the operator space duality. Moreover, unicity has to be understood modulo an equivalence relation: for any $x = (x_j) \in M_N(E)^n$ we define $Orb(x)$ as the set of all x' of the form $x' = (ux_jv) \in M_N(E)^n$ for some $u, v \in U(N)$. Then if x is “norming” some point, any $x' \in Orb(x)$

is also ‘norming’ that same point. When E is an operator space and $x \in M_N(E)$, we will say that $y \in M_N(E^*)$ M_N -norms x if $\|\sum x_j \otimes y_j\| = \|x\|_{M_N(E)} \|y\|_{M_N(E^*)}$. We will say that x is M_N -smooth in $M_N(E)$ if the only points y with $\|y\|_{M_N(E^*)} = 1$ that M_N -norm x are all in a single orbit in $M_N(E^*)$. Let us now turn to the case $E = OH_n$. There we show that, if $x \in U(N)^n$ is viewed as an element of $M_N(\ell_2^n)$, then x is M_N -smooth in $M_N(E)$ iff x is an ε -quantum expander for some $\varepsilon > 0$.

More generally, in Lemma 1.12 we prove a more precise quantified version of this: if x is an ε -quantum expander and if two points $y, z \in M_N(E^*)$ both M_N -norm x up to some error δ , then the distance of the orbits $Orb(y)$ and $Orb(z)$ is uniformly small, i.e. majorized by a function $f_\varepsilon(\delta)$ that tends to 0 when $\delta \rightarrow 0$. Here the distance is meant with respect to the renormalized Euclidean norm $y \mapsto (nN)^{-1/2} \|y\|_2$ for which any $y \in U(N)^n$ has norm 1 (where $\|\cdot\|_2$ denotes here the norm in $\ell_2(n \times N^2)$).

This also has a geometric application. Consider the following problem for an n -dimensional normed space E : Given a constant $C > 1$, estimate the minimal number $k = k_E(C)$ of functionals f_1, \dots, f_k in the dual E^* such that

$$\forall x \in E \quad \sup_{1 \leq j \leq k} |f_j(x)| \leq \|x\| \leq C \sup_{1 \leq j \leq k} |f_j(x)|.$$

Geometrically this means that (in the real case) the symmetric convex body that is the unit ball of E^* is equivalent (up to the factor C) to a polyhedron with vertices included in $\{\pm f_j\}$ and hence with at most $2k$ vertices (so its polar, that is equivalent to the unit ball of E , has at most $2k$ faces). For instance, the n -dimensional cube has 2^n vertices and $2n$ faces. When E has (real) dimension n it is well known (see e.g. [25, p.49-50]) that

$$k_E(C) \leq \left(\frac{3C}{C-1}\right)^n.$$

For example if $C = 2$ we have $k_E(C) \leq 6^n$. This exponential order of growth in n is optimal for $E = \ell_2^n$ (or ℓ_p^n for $1 \leq p < \infty$); but of course $k_E(C) = n$ for $E = \ell_\infty^n$, and there is important available information and a conjecture (see [22]) about conditions on a general sequence $\{E(n) \mid n \geq 1\}$ with $\dim(E(n)) = n$ ensuring that $k_{E(n)} \geq \exp cn$ for some $c > 0$.

We now describe the matricial analogue of k_E that we estimate using quantum expanders. Let E be an operator space. Fix an integer $N \geq 1$. We denote by $k_E(N, C)$ the smallest k such that there are linear maps $f_j : E \rightarrow M_N$ ($1 \leq j \leq k$) satisfying

$$\forall x \in M_N(E) \quad \sup_{1 \leq j \leq k} \|(Id \otimes f_j)(x)\|_{M_N(M_N)} \leq \|x\|_{M_N(E)} \leq C \sup_{1 \leq j \leq k} \|(Id \otimes f_j)(x)\|_{M_N(M_N)}.$$

It is not hard to adapt the corresponding Banach space argument to show that for *any* n -dimensional E , any $C > 1$ and any N we have

$$k_E(N, C) \leq \left(\frac{3C}{C-1}\right)^{2nN^2} = \exp 2 \log\left(\frac{3C}{C-1}\right) nN^2.$$

Using the ‘packing’ of ε -quantum expanders described above, we can show that the operator space version of Hilbert space (i.e. the space OH from [26]) satisfies a lower bound of the same order of growth, namely we show for $E = OH_n$ (see Theorem 2.8) there are numbers $C_1 > 1$, $b > 0$ such that for any n large enough and any N we have

$$(0.3) \quad k_E(N, C_1) \geq \exp b n N^2.$$

Moreover, this also holds for $E = \ell_1^n$ with its maximal operator space structure and for $E = R_n + C_n$ (see Remark 2.10).

We also show (see Theorem 2.15) that for any $R > 1$ and for any n, N suitably large there is a collection $\{E_t \mid t \in T_1\}$ of n -dimensional subspaces of M_N (each spanned by an n -tuple of unitary matrices) with cardinality $\geq \exp \beta_R n N^2$ such that the cb -distance $d_{cb}(E_s, E_t)$ of any distinct pair in T_1 satisfies

$$d_{cb}(E_s, E_t) \geq R.$$

The cb -distance d_{cb} is the analogue of the Banach-Mazur distance for operator spaces. The preceding shows that the metric entropy of the space of n -dimensional operator spaces equipped with the (so-called) “distance” d_{cb} is extremely large for small distances. This can be viewed as a somewhat more quantitative version of the non-separability of the space of n -dimensional operator spaces first proved in [13]. We plan to return to this in a future publication (see [30]).

The above (0.3) suggests that the class of finite dimensional operator spaces E such that $\log k_E(N, C)/N^2 \rightarrow 0$ should be investigated. We call such spaces *matricially subGaussian*.

In the forthcoming paper [29] we introduce a class of operator spaces, that we call “subexponential”, for which the same Grothendieck type factorization theorem from [13, 31] still holds (see the recent paper [32] for simpler proofs of the latter). We also give there examples of non-exact subexponential operator spaces or C^* -algebras.

The definition of “subexponential” involves the growth of a sequence of integers $N \mapsto K_E(N, C)$ attached to an operator space E (and a constant $C > 1$), in a way that is similar but seems different from $k_E(N, C)$. We denote by $K_E(N, C)$ the smallest K such that there is a single (embedding) linear map $f : E \rightarrow M_K$ satisfying

$$\forall x \in M_N(E) \quad \|(Id \otimes f)(x)\|_{M_N(M_K)} \leq \|x\|_{M_N(E)} \leq C \|(Id \otimes f)(x)\|_{M_N(M_N)}.$$

Roughly the latter sequence is bounded iff E is exact with exactness constant $\leq C$ (in the sense of [27, §17]) while it is such that $\log K_E(N, C)/N \rightarrow 0$ iff E is C -subexponential.

Note: There is an obvious upper bound (for a fixed constant C) $K_E(N, C) \leq N k_E(N, C)$, so the growth of K_E is dominated by that of k_E , but we know nothing in the converse direction. Various other questions are mentioned at the end of §3.

1. Quantum Expanders

Fix integers n, N . Throughout this paper we denote by M_N the space of $N \times N$ complex matrices and by $U(N)$ the subset of $N \times N$ unitary matrices.

We identify M_N with the space $B(\ell_2^N)$ of bounded operators on the N -dimensional Hilbert space denoted by ℓ_2^N .

We denote by tr (resp. τ_N) the usual trace (resp. the normalized trace) on M_N . Thus $\tau_N = N^{-1}tr$. We denote by S_2^N the Hilbert space obtained by equipping M_N with the corresponding scalar product. The associated norm is the classical Hilbert-Schmidt norm.

For simplicity we denote by

$$H = L_2(\tau_N),$$

i.e. H is the Hilbert space obtained by equipping the space M_N with the norm

$$\|\xi\|_H = (N^{-1}tr(|\xi|^2))^{1/2} = N^{-1/2}\|x\|_{S_2^N}.$$

We denote

$$H_0 = \{I\}^\perp \subset H.$$

Throughout this paper, we consider operators of the form $T = \sum x_j \otimes \bar{y}_j$, with $x_j, y_j \in M_N$, that we view as acting on $\ell_2^N \otimes \overline{\ell_2^N}$. Identifying as usual $\ell_2^N \otimes \overline{\ell_2^N}$ with S_2^N , we may consider T as an operator acting on M_N defined by

$$\forall \xi \in M_N \quad T(\xi) = \sum x_j \xi y_j^*,$$

and we then have

$$(1.1) \quad \left\| \sum x_j \otimes \bar{y}_j \right\| = \sup \{ \left\| \sum x_j \xi y_j^* \right\|_2 \mid \xi \in M_N, \|\xi\|_2 \leq 1 \} = \sup \{ \left| \sum \text{tr}(x_j \xi y_j^* \eta^*) \right| \mid \|\xi\|_2 \leq 1, \|\eta\|_2 \leq 1 \},$$

or equivalently $\left\| \sum x_j \otimes \bar{y}_j : \ell_2^N \otimes \overline{\ell_2^N} \rightarrow \ell_2^N \otimes \overline{\ell_2^N} \right\| = \|T : S_2^N \rightarrow S_2^N\|$. Actually it will be convenient to view T as an operator acting on $H = L_2(\tau_N)$. We have trivially

$$\|T\|_{B(H)} = \|T\|_{B(S_2^N)}.$$

Let $x = (x_j) \in (M_N)^n$ and $y = (y_j) \in (M_N)^n$. Let $Orb(x)$ denote the 2-sided unitary orbit of $x = (x_j)$, i.e.

$$Orb(x) = \{(ux_jv) \mid u, v \in U(N)\}.$$

We will denote

$$d(x, y) = \left(\sum_j \|x_j - y_j\|_{L_2(\tau_N)}^2 \right)^{1/2},$$

and

$$d'(x, y) = \inf \{d(x', y) \mid x' \in Orb(x)\} = \inf \{d(x', y') \mid x' \in Orb(x), y' \in Orb(y)\}.$$

The last equality holds because of the 2-sided unitary invariance of the norm in S_2^N or equivalently of $H = L_2(\tau_N)$.

Definition 1.1. Fix $\delta > 0$. We will say that x, y in M_N^n are δ -separated if

$$\left\| \sum x_j \otimes \bar{y}_j \right\| \leq (1 - \delta) \left\| \sum x_j \otimes \bar{x}_j \right\|^{1/2} \left\| \sum y_j \otimes \bar{y}_j \right\|^{1/2}.$$

A family of elements is called δ -separated if any two distinct members in it are δ -separated.

Let $x = (x_j) \in M_N^n$ and $y = (y_j) \in M_N^n$ be normalized so that $\left\| \sum x_j \otimes \bar{x}_j \right\| = \left\| \sum y_j \otimes \bar{y}_j \right\| = 1$. Equivalently, this definition means that for any $\xi, \eta \in M_N$ in the unit ball of S_2^N we have

$$\left| \sum \text{tr}(x_j \xi y_j^* \eta^*) \right| \leq 1 - \delta.$$

Using polar decompositions $\xi = u|\xi|$ and $\eta = v|\eta|$, $\left| \sum \text{tr}(x_j \xi y_j^* \eta^*) \right| = \left| \sum \text{tr}(x_j u |\xi| y_j^* |\eta| v^*) \right|$. Let $\hat{x}_j = v^* x_j u$. Equivalently we have for any u, v unitary

$$\left| \sum \text{tr}(\hat{x}_j |\xi| y_j^* |\eta|) \right| \leq 1 - \delta.$$

A fortiori, taking $|\xi| = |\eta| = N^{-1/2}I$ we find $\left| \sum \tau_N(\hat{x}_j y_j^*) \right| \leq 1 - \delta$ and hence

$$d(\hat{x}, y)^2 \geq 2\delta$$

and hence taking the inf over u, v unitary, the δ -separation of x, y implies

$$(1.2) \quad d'(x, y) \geq (2\delta)^{1/2}.$$

In other words, rescaling this to the case when $n^{1/2}x_j, n^{1/2}y_j, \xi, \eta$ are all unitary, we have proved:

Lemma 1.2. Consider n -tuples $x = (x_j) \in U(N)^n$ and $y = (y_j) \in U(N)^n$. If x, y are δ -separated then $d'(x, y) \geq (2\delta n)^{1/2}$.

Recall that we denote

$$H_0 = \{I\}^\perp.$$

To any n -tuple $u = (u_j) \in U(N)^n$ we associate the operator $(\sum u_j \otimes \bar{u}_j)(1 - P)$ on $\ell_2^N \otimes \overline{\ell_2^N}$ where P denotes the \perp -projection onto the scalar multiples of $I = \sum e_j \otimes \bar{e}_j$. Equivalently, up to the normalization, we will consider

$$T_u : H_0 \rightarrow H_0$$

defined for all $\xi \in H_0$ by

$$T_u(\xi) = \sum u_j \xi u_j^*.$$

We will denote by

$$S_\varepsilon = S_\varepsilon(n, N) \subset U(N)^n$$

the set of all n -tuples $u = (u_j) \in U(N)^n$ such that

$$\|T_u : H_0 \rightarrow H_0\| \leq \varepsilon n.$$

Equivalently, this means $\forall x \in M_N$, we have

$$\left\| \sum u_j (x - \tau_N(x) I) u_j^* \right\|_H \leq \varepsilon n \|x\|_H.$$

Our goal is to prove the following:

Theorem 1.3. For any $0 < \delta < 1$ there is a constant $\beta_\delta > 0$ such that for each $0 < \varepsilon < 1$ and for all sufficiently large integer n (i.e. $n \geq n_0$ with n_0 depending on ε and δ) and for all $N \geq 1$, there is a δ -separated subset

$$T \subset S_\varepsilon$$

such that

$$|T| \geq \exp \beta_\delta n N^2.$$

Remark 1.4. Actually, the proof will show that if we are given sets $A_N \subset U(N)^n$ such that $\inf_N \mathbb{P}(A_N) \geq \alpha > 0$, then for each N we can find a subset T as above with $T \subset A_N \cap S_\varepsilon$, but with β_δ and n_0 now also depending on α .

Remark 1.5. The order of growth of our lower bound $\exp \beta n N^2$ in Theorem 1.3 is roughly optimal because of the upper bound given explicitly in [10] (and implicitly in [37]). The latter upper bound can be proved as follows. Let m_{\max} be the maximal number of a δ -separated family in $U(N)^n$. Consider the normed space obtained by equipping $M(N)^n$ with the norm $|||x||| = \|\sum x_j \otimes \bar{x}_j\|^{1/2}$. Then since its (real) dimension is $2nN^2$, by a well known volume argument ([25, p.49-50]) there cannot exist more than $(1 + 2/\delta')^{2nN^2}$ elements in its unit ball at mutual $|||\cdot|||$ -distance $\geq \delta'$. Note that $d(x, y) \leq |||x - y|||$ for any pair x, y in $M(N)^n$. Thus, if $u, v \in U(N)^n$ are δ -separated in the above sense then $x = n^{-1/2}u$ and $y = n^{-1/2}v$ are in the $|||\cdot|||$ -unit ball and by (1.2) we have $|||x - y||| \geq (2\delta)^{1/2}$, therefore

$$m_{\max} \leq (1 + \sqrt{2/\delta})^{2nN^2} \leq \exp\{2\sqrt{2/\delta} nN^2\}.$$

Remark 1.6. Let G be a Kazhdan group (see [1]) with generators t_1, \dots, t_n , so that there is $\delta > 0$ such that $\|\sum_1^n \pi(t_j)\| \leq n(1 - \delta)$ for any unitary representation without any invariant (non zero) vector. Let $\mathcal{I} = \mathcal{I}(N)$ denote the set of N -dimensional irreducible representations $\pi : G \rightarrow U(N)$. It is known (see [1]) that the latter set is finite and in fact there is a uniform bound on $|\mathcal{I}(N)|$ for each N . For any $\pi \in \mathcal{I}$ we set

$$u_j^\pi = \pi(t_j).$$

Then (here by $\pi \neq \sigma$ we mean π is not equivalent to σ)

$$\sup_{\pi \neq \sigma \in \mathcal{I}} \left\| \sum u_j^\pi \otimes \overline{u_j^\sigma} \right\| \leq n(1 - \delta),$$

so that the family $\{u^\pi \mid \pi \in \mathcal{I}\} \subset U(N)^n$ is δ -separated in the above sense. By the preceding Remark, we know $|\mathcal{I}(N)| \leq m_{\max} \leq \exp c_\delta n N^2$. The problem to estimate the maximal possible value of $|\mathcal{I}(N)|$ when $N \rightarrow \infty$ (with δ and n remaining fixed, but G possibly varying) is investigated in [19]: some special cases are constructed in [19] for which $|\mathcal{I}(N)|$ grows like $\exp cN$, however we feel that Theorem 1.3 gives evidence that there should exist cases for which $|\mathcal{I}(N)|$ grows like $\exp cN^2$.

Remark 1.7. Recall (see [27, p. 324. Th. 20.1]) that for any n -tuple of unitary operators on any Hilbert space H we have

$$\left\| \sum u_j \otimes \bar{u}_j \right\| \geq 2\sqrt{n-1}.$$

Note that $2\sqrt{n-1} < n$ for all $n \geq 3$ (so there is also an $0 < \varepsilon < 1$ such that $2\sqrt{n-1} + \varepsilon n < n$).

Let $0 < \varepsilon < 1$. In analogy with Ramanujan graphs (see [17]) an n -tuple $u = (u_j) \in U(N)^n$ will be called ε -Ramanujan if

$$\|T_u : H_0 \rightarrow H_0\| \leq 2\sqrt{n-1} + \varepsilon n.$$

We will denote by

$$R_\varepsilon = R_\varepsilon(n, N) \subset U(N)^n$$

the set of all such n -tuples.

We refer to [17, 12] for more information on expanders and Ramanujan graphs.

The next result due to Hastings [11] has been a crucial inspiration for our work:

Lemma 1.8 (Hastings). *If we equip $U(N)^n$ with its normalized Haar measure \mathbb{P} , then for each n and $\varepsilon > 0$ the set $R_\varepsilon(n, N)$ defined above satisfies*

$$\lim_{N \rightarrow \infty} \mathbb{P}(R_\varepsilon(n, N)) = 1.$$

This is best possible in the sense that Lemma 1.8 fails if $2\sqrt{n-1}$ is replaced (in the definition of $R_\varepsilon(n, N)$) by any smaller number. However, we do not really need this sharp form of Lemma 1.8 (unless we insist on making $n_0(\varepsilon)$ as small as possible in Theorem 1.3). So we give in the appendix a quicker proof of a result that suffices for our needs (where $2\sqrt{n-1}$ is replaced by $C'\sqrt{n}$, C' being a numerical constant) and which in several respects gives us better estimates than Lemma 1.8.

Lemma 1.9 below can be viewed as a non-commutative variant of results in [24] (see also [23] where the non-commutative case is already considered) in the style of [18] (see also [7, 21]). We view this as a (weak) sort of non-commutative Sauer lemma, that it might be worthwhile to strengthen.

In the next two lemmas, we equip $U(N)^n$ with the metric d (we also use d'), and for any subset $A \subset U(N)^n$ and any $\varepsilon > 0$ we denote by $N(A, d, \varepsilon)$ the smallest number of open d -balls of radius ε with center in $U(N)^n$ that cover A .

Lemma 1.9. *Let $a > 0$. Let $A \subset U(N)^n$ be a (measurable) subset with $\mathbb{P}(A) > a$. Then, for any $c < \sqrt{2}$, $N(A, d, c\sqrt{n}) \geq a \exp KrnN^2$ where $r = (1 - c^2/2)^2$ and K is a universal constant. Assuming moreover that $a \geq \exp -KrnN^2/2$ (note that there is $n_0(a, r)$ so that this holds for all $n \geq n_0(a, r)$ and all $N \geq 1$), we find that $N(A, d, c\sqrt{n}) \geq \exp bnN^2$ where $b = Kr/2$.*

Proof. Let $\Omega = U(N)^n$. We may clearly assume (by Haar measure inner regularity) that A is compact. Let $\mathcal{N} = N(A, d, c\sqrt{n})$. By definition, A is included in the union of \mathcal{N} open balls with d -radius $c\sqrt{n}$. By translation invariance of d and \mathbb{P} , all these balls have the same \mathbb{P} -measure equal to $F(c)$. Therefore $a < \mathbb{P}(A) \leq \mathcal{N}F(c)$ and hence

$$aF(c)^{-1} < \mathcal{N}.$$

Thus we need a lower bound for $F(c)^{-1}$. Let u denote the unit in $U(N)^n$ so that $u_j = 1$ for $1 \leq j \leq n$. Using a ball centered at u to compute $F(c)$, we have

$$F(c) = \mathbb{P}\{\omega \in U(N)^n \mid \sum_1^n \text{tr}(|\omega_j - 1|^2) < c^2 nN\}.$$

Since $\sum_1^n \text{tr}(|\omega_j - 1|^2) = 2Nn - 2 \sum_1^n \Re \text{tr}(\omega_j)$, we have

$$F(c) = \mathbb{P}\{\omega \mid \sum_1^n \Re \text{tr}(\omega_j) > nN(1 - c^2/2)\}.$$

We will now use the known subGaussian property of $\sum_1^n \Re \text{tr}(\omega_j)$: there is a universal constant K such that for any $\lambda > 0$ we have

$$(1.3) \quad \mathbb{P}\{\omega \mid \sum_1^n \Re \text{tr}(\omega_j) > \lambda\} \leq \exp -K\lambda^2/n.$$

Taking this for granted, let us complete the proof. Fix $c < \sqrt{2}$. Recall $r = (1 - c^2/2)^2 > 0$, this yields

$$F(c) \leq \exp -KnN^2r.$$

Thus we conclude that

$$\mathcal{N} > a \exp KrnN^2.$$

Taking $c = 1, r = 1/4$, the last assertion becomes obvious.

Let us now give a quick argument for the known inequality (1.3): We will denote by $Y^{(N)}$ a random $N \times N$ -matrix with i.i.d. complex Gaussian entries with mean zero and second moment equal to $N^{-1/2}$, and we denote by $(Y_j^{(N)})$ a sequence of i.i.d. copies of $Y^{(N)}$. It is well known that the polar decomposition $Y^{(N)} = U|Y^{(N)}|$ is such that U is uniformly distributed over $U(N)$ and independent of $|Y^{(N)}|$. Moreover there is an absolute constant $\chi > 0$ such that $\mathbb{E}|Y^{(N)}| = \chi^{-1}I$. See e.g. [18, p. 80]. Therefore, we have a conditional expectation operator \mathcal{E} (corresponding to integrating the modular part) such that $\sum \Re \text{tr}(\omega_j) = \chi \mathcal{E}(\sum \Re \text{tr}(Y_j^{(N)}))$, where ω_j denotes the unitary part in the polar decomposition of $Y_j^{(N)}$.

Then, since $x \mapsto \exp wx$ is convex for any $w > 0$, we have the announced subGaussian property

$$\mathbb{E} \exp w \sum \Re \text{tr}(\omega_j) \leq \mathbb{E} \exp w\chi \sum \Re \text{tr}(Y_j^{(N)}) = \exp(\chi^2 w^2 n/4),$$

from which follows, by Tchebyshev's inequality, that $\mathbb{P}\{\sum \Re \text{tr}(\omega_j) > \lambda\} \leq \exp(\chi^2 w^2 n/4 - \lambda w)$ and optimising w so that $\lambda = \chi^2 wn/2$ we finally obtain

$$\mathbb{P}\{\sum \Re \text{tr}(\omega_j) > \lambda\} \leq \exp(-K\lambda^2/n),$$

with $K = \chi^{-2}$. The above simple argument follows [18, ch. 5], but, in essence, (1.3) can be traced back to [8, Lemma 3]. \square

The next Lemma is a simple covering argument.

Lemma 1.10. *Fix $b, c > 0$. Let $A \subset U(N)^n$ be a subset with $N(A, d, c\sqrt{n}) \geq \exp bnN^2$. Fix $c' < c$ and $b' < b$. Then there is an integer n_0 (depending only on $b - b'$ and $c - c'$ and independent of N) such that if $n \geq n_0$ we have $N(A, d', c'\sqrt{n}) \geq \exp b'nN^2$ and there is a subset $T' \subset A$ with $|T'| \geq \exp b'nN^2$ such that $d'(s, t) \geq c'\sqrt{n} \forall s \neq t \in T'$.*

Proof. Fix $\varepsilon > 0$. It is well known that there is an ε -net $\mathcal{N}_\varepsilon \subset U(N)$ with respect to the operator norm with $|\mathcal{N}_\varepsilon| \leq (K/\varepsilon)^{2N^2}$. Indeed, since the real dimension of M_N is $2N^2$, a classical volume argument (see e.g. ([25, p.49-50])) produces such a net inside the unit ball of M_N . It can then be adjusted to be inside $U(N)$. See also [33, p. 175] for more delicate estimates. For any $x \in U(N)^n$, we have $d(uxv, u'xv') \leq (\|u - u'\| + \|v - v'\|)\sqrt{n}$ for any $u, u', v, v' \in U(N)$. Therefore we have $N(\text{Orb}(x), d, 2\varepsilon\sqrt{n}) \leq |\mathcal{N}_\varepsilon|^2 \leq \exp 4N^2 \log(K/\varepsilon)$. From this follows immediately that

$$N(A, d, c'\sqrt{n} + 2\varepsilon\sqrt{n}) \leq N(A, d', c'\sqrt{n}) \exp 4N^2 \log(K/\varepsilon).$$

Since $c' < c$ we can choose $\varepsilon > 0$ so that $c' + 2\varepsilon = c$. Then by our assumption $N(A, d, c'\sqrt{n} + 2\varepsilon\sqrt{n}) = N(A, d, c\sqrt{n}) \geq \exp bnN^2$. Thus we find

$$N(A, d', c'\sqrt{n}) \geq \exp bnN^2 \exp -4N^2 \log(K/\varepsilon).$$

Since $b - b' > 0$ there is clearly an integer n_0 (depending only on $b - b'$ and $\varepsilon = (c - c')/2$) such that $4 \log(K/\varepsilon) < (b - b')n$ for all $n \geq n_0$. Thus we obtain

$$N(A, d', c'\sqrt{n}) \geq \exp b'nN^2.$$

The last assertion is then clear: any maximal subset $T' \subset A$ such that $d'(s, t) \geq c'\sqrt{n} \forall s \neq t \in T'$ must satisfy (by maximality) $N(A, d', c'\sqrt{n}) \leq |T'|$. \square

In general, for a pair $u, v \in U(N)^n$, δ -separation is a much stronger condition than separation with respect to the distance d' . The main virtue of the next two Lemmas is to show that for a pair $u, v \in S_\varepsilon$ with ε suitably small, the two conditions become essentially equivalent. To prove these, we will now crucially use the spectral gap.

Lemma 1.11. *Let $0 < \varepsilon, \varepsilon' < 1$. Let $u = (u_j) \in U(N)^n$ and $v = (v_j) \in M_N^n$ merely such that $\|\sum v_j \otimes \bar{v}_j\| \leq n$. Assume $u \in S_\varepsilon$ and also*

$$(1.4) \quad d'(u, v) \geq \sqrt{2n(1 - \varepsilon')}.$$

Then

$$\|\sum u_j \otimes \bar{v}_j\| \leq n \left(\varepsilon'^{1/5} (2^{-4/5} + 2^{6/5}) + 2\varepsilon^{1/2} \right).$$

Moreover, if we assume in addition that $v \in S_\varepsilon$, then the preceding estimate can be improved to

$$(1.5) \quad \|\sum u_j \otimes \bar{v}_j\| \leq n(3\varepsilon'^{1/3} + 2\varepsilon).$$

Conversely, it is easy to show that for any pair $u, v \in M_N^n$ such that $\sum \tau_N(|u_j|^2) = \sum \tau_N(|v_j|^2) = n$ (in particular for any $u, v \in U(N)^n$)

$$(1.6) \quad \|\sum u_j \otimes \bar{v}_j\| \leq n\varepsilon'$$

implies

$$(1.7) \quad d'(u, v) \geq \sqrt{2n(1 - \varepsilon')}.$$

Proof. Let $\|v\|_{\mathcal{H}}^2 = (1/n) \sum_j \tau_N |v_j|^2$. Note that $\|v\|_{\mathcal{H}} \leq 1$ and $\|u\|_{\mathcal{H}} \leq 1$. For any $x \in M_N$, we will denote $\|x\|_{L_2(\tau_N)} = (\tau_N |x|^2)^{1/2}$ and $\|x\|_{L_1(\tau_N)} = \tau_N |x|$. Note for later use that a consequence of Cauchy-Schwarz is

$$(1.8) \quad \forall x, y \in M_N \quad \|xy\|_{L_1(\tau_N)} \leq \|x\|_{L_2(\tau_N)} \|y\|_{L_2(\tau_N)}.$$

Recall that

$$\left\| \sum u_j \otimes \bar{v}_j \right\| = \sup \{ \left| \sum \tau_N (u_j x v_j^* y^*) \right| \mid x, y \in B_{L_2(\tau_N)} \}.$$

Let us denote $x.u.y = (xu_j y)_{1 \leq j \leq n}$ and let F be the bilinear form on $M_N \times M_N$ defined by

$$F(x, y) = \langle x.u.y, v \rangle_{\mathcal{H}} = (1/n) \sum_j \tau_N (x u_j y v_j^*).$$

Then

$$\|(1/n) \sum u_j \otimes \bar{v}_j\| = \|F : L_2(\tau_N) \times L_2(\tau_N) \rightarrow \mathbb{C}\|.$$

We start by the proof of (1.5), assuming that both u, v belong to S_ε . Firstly we claim that for any $x, y \in M_N$ we have

$$|F(x, y)| \leq \tau_N |x| \tau_N |y| + \varepsilon \|(1-P)(|x|)\|_{L_2(\tau_N)} \|(1-P)(|y|)\|_{L_2(\tau_N)}$$

and hence assuming $x, y \in B_{L_2(\tau_N)}$ we have

$$(1.9) \quad |F(x, y)| \leq \tau_N |x| \tau_N |y| + \varepsilon.$$

To check this claim we use polar decompositions $x = U|x|$, $y = V|y|$ and we write

$$\begin{aligned} F(x, y) &= \langle U|x|.u.V|y|, v \rangle_{\mathcal{H}} = (1/n) \sum_j \tau_N ([|x|^{1/2} u_j V|y|^{1/2}] [|y|^{1/2} v_j^* U|x|^{1/2}]) \\ &= \langle |x|^{1/2}.u.V|y|^{1/2}, |x|^{1/2} U^* v |y|^{1/2} \rangle_{\mathcal{H}}. \end{aligned}$$

Therefore

$$(1.10) \quad |F(x, y)| \leq \| |x|^{1/2}.u.V|y|^{1/2} \|_{\mathcal{H}} \| |x|^{1/2} U^* v |y|^{1/2} \|_{\mathcal{H}}.$$

Now we observe that if we denote again by T_u the operator acting on $L_2(\tau_N)$ defined by $T_u(x) = \sum u_j x u_j^* - n \tau_N(x) I$ (equivalently $T_u = (\sum u_j \otimes \bar{u}_j)(1-P)$) we have for any $a, b \in M_N$

$$\|a.u.b\|_{\mathcal{H}}^2 = (1/n) \tau_N ((n \tau_N(bb^*) + T_u(bb^*)) a^* a) = \tau_N(bb^*) \tau_N(a^* a) + (1/n) \tau_N(T_u(bb^*) a^* a)$$

and hence since $\|T_u\| \leq \varepsilon n$ and $T_u = (1-P)T_u(1-P)$

$$\|a.u.b\|_{\mathcal{H}}^2 \leq \tau_N(bb^*) \tau_N(a^* a) + \varepsilon \|(1-P)(bb^*)\|_{L_2(\tau_N)} \|(1-P)(a^* a)\|_{L_2(\tau_N)}.$$

This yields

$$\| |x|^{1/2}.u.V|y|^{1/2} \|_{\mathcal{H}}^2 \leq \tau_N |x| \tau_N |y| + \varepsilon \|(1-P)(|x|)\|_{L_2(\tau_N)} \|(1-P)(|y|)\|_{L_2(\tau_N)}.$$

A similar bound holds for $\| |x|^{1/2} U^* v |y|^{1/2} \|_{\mathcal{H}}$. Thus (1.10) leads to our claim.

Secondly by (1.4) for any $U, V \in U(N)$ we have

$$\|U.u.V - v\|_{\mathcal{H}}^2 \geq n^{-1} d'(u, v)^2 \geq 2(1 - \varepsilon')$$

and hence

$$\Re\langle U.u.V, v \rangle_{\mathcal{H}} \leq \varepsilon'.$$

Recall that the unit ball of M_N is the closed convex hull of $U(N)$. Thus we have

$$\|F : M_N \times M_N \rightarrow \mathbb{C}\| \leq \varepsilon'.$$

Let us assume $x, y \in B_{L_2(\tau_N)}$. Let p (resp. q) denote the spectral projection of $|x|$ (resp. $|y|$) corresponding to the spectral set $\{|x| \leq \lambda\}$ (resp. $\{|y| \leq \lambda\}$). Note that by Tchebyshev's inequality we have $\tau_N(1-p) \leq 1/\lambda^2$ (resp. $\tau_N(1-q) \leq 1/\lambda^2$). Let $x' = (1-p)|x|$ and $y' = (1-q)|y|$. By (1.8) (since $\|1-p\|_{L_2(\tau_N)} \leq \lambda^{-1/2}$) we have

$$\tau_N|x'| \leq \lambda^{-1}.$$

Similarly

$$\tau_N|y'| \leq \lambda^{-1}.$$

We now write

$$(1.11) \quad F(|x|, |y|) = F(p|x| + x', q|y| + y') = F(p|x|, q|y|) + F(x', |y|) + F(p|x|, y').$$

By (1.9) we have

$$|F(x', |y|)| \leq \tau_N|x'| \tau_N|y| + \varepsilon \leq \lambda^{-1} + \varepsilon$$

and similarly

$$|F(p|x|, y')| \leq \lambda^{-1} + \varepsilon.$$

Thus we deduce from (1.11)

$$|F(|x|, |y|)| \leq \varepsilon' \lambda^2 + 2(\lambda^{-1} + \varepsilon).$$

Choosing $\lambda = (\varepsilon')^{-1/3}$ to minimize over $\lambda > 0$ yields the upper bound $3\varepsilon'^{1/3} + 2\varepsilon$, when restricting to $x, y \geq 0$. Since our assumptions on the pair u, v are shared by the pair UuV, v for any $U, V \in U(N)$, we may apply the polar decompositions $x = U|x|$ and $y = V|y|$ to deduce the same upper bound for an arbitrary pair x, y . Thus we obtain (1.5).

We now turn to (1.4). There we assume only $u \in S_\varepsilon$ and $\|\sum v_j \otimes \bar{v}_j\| \leq n$. Then (since we still have $\| |x|^{1/2} U^* v |y|^{1/2} \|_{\mathcal{H}} \leq 1$) (1.9) can be replaced by

$$(1.12) \quad |F(x, y)| \leq (\tau_N|x| \tau_N|y| + \varepsilon)^{1/2},$$

and the preceding reasoning leads to

$$|F(x, y)| \leq \varepsilon' \lambda^2 + 2(\lambda^{-1} + \varepsilon)^{1/2} \leq \varepsilon' \lambda^2 + 2\lambda^{-1/2} + 2\varepsilon^{1/2}.$$

Choosing $\lambda = (2\varepsilon')^{-2/5}$ to minimize, we obtain the announced upper bound $\varepsilon'^{1/5}(2^{-4/5} + 2^{6/5}) + 2\varepsilon^{1/2}$, thus completing the proof of (1.4).

The converse implication (1.6) \Rightarrow (1.7) is obvious: Indeed, for any $U, V \in U(N)$ (1.6) implies $|\sum \tau_N(Uu_j V v_j^*)| \leq n\varepsilon'$ and hence since we assume $\sum \tau_N(u_j^* u_j) = \sum \tau_N(v_j^* v_j) = n$ we have

$$d(U.u.V, t)^2 = 2n - 2\Re \sum \tau_N(Uu_j V v_j^*) \geq 2n(1 - \varepsilon')$$

and taking the infimum over $U, V \in U(N)$ we obtain (1.7). \square

Proof of Theorem 1.3. Our original proof was based on Hastings's Lemma 1.8, but the current proof, based instead on (4.6) allows for more uniformity with respect to N . Note however that if we could prove a sharper form of (1.5) (e.g. with ε in place of 2ε) then Lemma 1.8 would allow us to cover values of n as small as $n = 3$, for all N large enough (while using (4.6) requires $C'n^{-1/2} < n$).

By (4.6) there is a constant C' such that such that

$$\forall N \geq 1 \quad \mathbb{E} \left\| \sum_1^n U_j \otimes \bar{U}_j (1 - P) \right\| \leq C' \sqrt{n}.$$

where \mathbb{E} is with respect to the normalized Haar measure on $U(N)^n$. By Tchebyshev's inequality, this implies $\mathbb{P}(S_\varepsilon) > 1/2$ for all $N \geq 1$, assuming only that $n > n_0(\varepsilon)$ for a suitably adjusted value of $n_0(\varepsilon)$ (say we require $\varepsilon^{-1}C'n^{-1/2} < 1/2$). We will now apply Lemmas 1.9 and 1.10 to the subset $A = S_\varepsilon$.

Fix $0 < \varepsilon, \delta < 1$. Let $0 < \varepsilon' < 1$ be such that $3\varepsilon'^{1/3} = (1 - \delta)/2$ and let ε_0 be such that $2\varepsilon_0 = (1 - \delta)/2$. Let $c' = \sqrt{2(1 - \varepsilon')}$ and $c = \sqrt{2(1 - \varepsilon'/2)}$ so that we have $c' < c < \sqrt{2}$.

Assume $\varepsilon \leq \varepsilon_0$. Let $r = \varepsilon'^2$ and $b = K\varepsilon'^2/2$ and say $b' = b/2$ so that $b - b'$ and $c - c'$ depend only on ε' (or equivalently on δ). For $n \geq n_0(\varepsilon, \varepsilon')$, Lemmas 1.9 and 1.10 give us a subset $T \subset S_\varepsilon$ such that $|T| \geq \exp K\varepsilon'^2 n N^2 / 4$ and such that $d'(s, t) \geq \sqrt{2(1 - \varepsilon')}$ for all $s \neq t$. Then Lemma 1.12 (specifically (1.5)) gives us that s, t are δ -separated since $\varepsilon \leq \varepsilon_0$ and our choice of $\varepsilon_0, \varepsilon'$ is adjusted so that $3\varepsilon'^{1/3} + 2\varepsilon_0 = 1 - \delta$. This completes the proof for any $\varepsilon \leq \varepsilon_0$ and in particular for $\varepsilon = \varepsilon_0$. The remaining case $\varepsilon_0 < \varepsilon < 1$ then follows automatically since $S_{\varepsilon_0} \subset S_\varepsilon$ if $\varepsilon_0 < \varepsilon$. \square

One defect of Lemma 1.11 is that when ε' is close to 1, its conclusion is void (however small ε can be). This is corrected by the next Lemma the main interest of which is the case when δ and $f_\varepsilon(\delta)$ are small.

Lemma 1.12. *Fix $0 < \varepsilon < 1$. There is a positive function $\delta \mapsto f_\varepsilon(\delta)$ defined for $0 < \delta < 1$ and such that $f_\varepsilon(\delta) = O_\varepsilon(\delta^{1/4})$ when $\delta \rightarrow 0$ satisfying the following property:*

Consider $u = (u_j) \in S_\varepsilon \subset U(N)^n$ and $v = (v_j) \in M_N^n$ such that $\|\sum v_j \otimes \bar{v}_j\| \leq n$. The condition

$$(1.13) \quad d'(u, v) \geq f_\varepsilon(\delta) \sqrt{n}$$

implies

$$(1.14) \quad \left\| \sum u_j \otimes \bar{v}_j \right\| \leq n(1 - \delta).$$

Proof. Assume by contradiction that $\|\sum u_j \otimes \bar{v}_j\| > n(1 - \delta)$. Then there are ξ, η in the unit sphere of $H = L_2(\tau_N)$ such that

$$\Re \tau_N \left(\sum u_j \xi v_j^* \eta^* \right) > n(1 - \delta).$$

Let $\xi = U|\xi|$ and $\eta = V|\eta|$ be their polar decompositions, and let $w_j = V^* u_j U$ so that we can write

$$(1.15) \quad \Re \tau_N \left(\sum w_j |\xi| v_j^* |\eta| \right) > n(1 - \delta).$$

Recall $T_u(\xi) = \sum u_j \xi u_j^*$. Note that since $U \otimes \bar{U}$ and $V \otimes \bar{V}$ preserve I (and hence I^\perp), we have $\|T_w\| = \|T_u\|$. Therefore $w \in S_\varepsilon$. Using the scalar product in H we have

$$\Re \sum \langle |\eta|^{1/2} w_j |\xi|^{1/2}, |\eta|^{1/2} v_j |\xi|^{1/2} \rangle > n(1 - \delta),$$

and by Cauchy-Schwarz

$$\left(\sum \| |\eta|^{1/2} w_j |\xi|^{1/2} \|_H^2 \right)^{1/2} \left(\sum \| |\eta|^{1/2} v_j |\xi|^{1/2} \|_H^2 \right)^{1/2} > n(1 - \delta).$$

Note that since $\|\sum v_j \otimes \bar{v}_j\| \leq n$ we have $\langle (\sum v_j \otimes \bar{v}_j)|\xi\rangle, |\eta\rangle = \sum \|\eta\|^{1/2} v_j |\xi|^{1/2}\|_H^2 \leq n$, and similarly with w_j in place of v_j . Thus the last inequality implies a fortiori

$$(1.16) \quad \langle (\sum w_j \otimes \bar{w}_j)|\xi\rangle, |\eta\rangle > n(1 - \delta)^2,$$

and the same with v_j in place of w_j .

Let $e = (1 - P)|\xi\rangle$ and $d = (1 - P)|\eta\rangle$. Recall that $P(|\xi\rangle) = \tau_N(|\xi\rangle)I$ and $P(|\eta\rangle) = \tau_N(|\eta\rangle)I$, and $|\xi\rangle, |\eta\rangle$ are unit vectors, so that $\tau_N(|\xi\rangle) = (1 - \|e\|_H^2)^{1/2}$ and $\tau_N(|\eta\rangle) = (1 - \|d\|_H^2)^{1/2}$. Let $\omega = \|e\|_H \|d\|_H$. By Cauchy-Schwarz we have $\omega + (1 - \|e\|_H^2)^{1/2}(1 - \|d\|_H^2)^{1/2} \leq 1$ and hence $(1 - \|e\|_H^2)^{1/2}(1 - \|d\|_H^2)^{1/2} \leq 1 - \omega$. Since $w \in S_\varepsilon$, we have

$$\langle (\sum w_j \otimes \bar{w}_j)|\xi\rangle, |\eta\rangle = \langle T_w e, d \rangle + n\tau_N(|\xi\rangle)\tau_N(|\eta\rangle) \leq \varepsilon n\omega + n(1 - \omega).$$

Thus, (1.16) yields

$$(1.17) \quad \omega \leq (1 - \varepsilon)^{-1}(2\delta - \delta^2).$$

Moreover, (1.16) implies

$$(1.18) \quad n^{-1} \sum \|w_j |\xi| w_j^* - |\eta|\|_H^2 = 2 - 2n^{-1} \langle (\sum w_j \otimes \bar{w}_j)|\xi\rangle, |\eta\rangle < 2(2\delta - \delta^2).$$

But since $(I - P)(w_j |\xi| w_j^* - |\eta|) = w_j e w_j^* - d$ for each j , we have

$$\|e\|_H - \|d\|_H = \|w_j e w_j^* - d\|_H \leq \|w_j e w_j^* - d\|_H \leq \|w_j |\xi| w_j^* - |\eta|\|_H$$

so that (1.18) implies $(\|e\|_H - \|d\|_H)^2 < 2(2\delta - \delta^2)$. Therefore

$$(1.19) \quad \|e\|_H^2 + \|d\|_H^2 < 2\omega + 2(2\delta - \delta^2) \leq 2(2\delta - \delta^2)((1 - \varepsilon)^{-1} + 1).$$

We can write

$$w_j |\xi| v_j^* |\eta| = w_j P(|\xi\rangle) v_j^* P(|\eta\rangle) + w_j e v_j^* P(|\eta\rangle) + w_j P(|\xi\rangle) v_j^* d + w_j e v_j^* d$$

and we find

$$\begin{aligned} n(1 - \delta) &< \Re \tau_N(\sum w_j |\xi| v_j^* |\eta|) \\ &\leq \Re \tau_N(\sum w_j v_j^*) \tau_N(|\xi\rangle) \tau_N(|\eta\rangle) + n\|e\|_H \tau_N(|\eta\rangle) + n\tau_N(|\xi\rangle) \|d\|_H + n\|e\|_H \|d\|_H \end{aligned}$$

and a fortiori

$$n(1 - \delta) < \Re \tau_N(\sum w_j v_j^*) \tau_N(|\xi\rangle) \tau_N(|\eta\rangle) + n\|e\|_H + n\|d\|_H + n\|e\|_H \|d\|_H.$$

Note that $n\|e\|_H + n\|d\|_H + n\|e\|_H \|d\|_H \leq n2^{1/2}(\|e\|_H^2 + \|d\|_H^2)^{1/2} + n\|e\|_H \|d\|_H$. By (1.19) and (1.17), we obtain

$$\tau_N(|\xi\rangle) \tau_N(|\eta\rangle) \Re \tau_N(\sum w_j v_j^*) > n(1 - \theta)$$

with

$$\theta = \delta + 2^{1/2} (2(2\delta - \delta^2)((1 - \varepsilon)^{-1} + 1))^{1/2} + (1 - \varepsilon)^{-1}(2\delta - \delta^2).$$

Note that θ is $O(\delta^{1/4})$ when $\delta \rightarrow 0$ and there is clearly some $\delta_\varepsilon > 0$ such that $0 < \theta < 1$ for all $\delta \leq \delta_\varepsilon$. Thus, assuming $0 < \delta \leq \delta_\varepsilon$ we have $1 - \theta > 0$ and we find

$$\Re \tau_N(\sum w_j v_j^*) > n(1 - \theta).$$

This is a lower bound for the real part of the scalar product in $\ell_2^n(H)$ of $w = (w_j)$ and $v = (v_j)$ which are both in the ball of radius \sqrt{n} . Therefore we deduce from this

$$d(w, v)^2 \leq 2n - 2\Re \tau_N(\sum w_j v_j^*) < 2n\theta.$$

If we now set $f_\varepsilon(\delta) = (2\theta)^{1/2}$ for all $\delta \leq \delta_\varepsilon$, and $f_\varepsilon(\delta) = 3$ (say) for $\delta_\varepsilon < \delta < 1$, we have in any case $d(w, v) < f_\varepsilon(\delta)\sqrt{n}$.

Thus we have proved that $\|\sum u_j \otimes \bar{v}_j\| > n(1 - \delta)$ implies $d'(u, v) < f_\varepsilon(\delta)\sqrt{n}$. This is equivalent to the fact that (1.13) implies (1.14), and moreover for each $0 < \varepsilon < 1$ there is a constant $c_\varepsilon > 0$ such that for any $0 < \delta < 1$ we have $f_\varepsilon(\delta) \leq c_\varepsilon \delta^{1/4}$. \square

Remark 1.13. Let $f_\varepsilon(\delta)$ be any function such that (1.13) \Rightarrow (1.14). Then, Lemma 1.12 has the following consequence: Assume $u = (u_j) \in S_\varepsilon$. Then for any $v = (v_j) \in U(k)^n$ with $k \leq (1 - f_\varepsilon(\delta)^2)N$ we have

$$\|\sum u_j \otimes \bar{v}_j\| \leq n(1 - \delta).$$

Indeed, if $\|\sum u_j \otimes \bar{v}_j\| > n(1 - \delta)$, then we set $v'_j = v_j \oplus 0 \in M_N$ so that $\|\sum v'_j \otimes \bar{v}'_j\| \leq n$, and also $\|\sum u_j \otimes \bar{v}'_j\| > n(1 - \delta)$. By Lemma 1.12, it follows that $d'(u, v') < f_\varepsilon(\delta)\sqrt{n}$. But since $|\langle u'_j, v'_j \rangle| \leq k/N$ for any $u'_j \in U(N)$, we have $d(u', v')^2 = n + n(k/N) - 2\sum \langle u'_j, v'_j \rangle \geq n(1 - k/N)$, and hence $d'(u, v') \geq \sqrt{n}(1 - k/N)^{1/2}$, which leads to $(1 - k/N)^{1/2} < f_\varepsilon(\delta)$. This contradiction concludes the proof.

2. Application to Operator Spaces

We start with a specific notation. Let $u : E \rightarrow F$ be a linear map between operator spaces. We denote for any given $N \geq 1$

$$u_N = Id \otimes u : M_N(E) \rightarrow M_N(F).$$

Moreover, if E, F are two operator spaces that are isomorphic as Banach spaces, we set

$$d_N(E, F) = \inf\{\|u_N\| \|(u^{-1})_N\|\}$$

where the inf runs over all the isomorphisms $u : E \rightarrow F$. We set $d_N(E, F) = \infty$ if E, F are not isomorphic.

Recall that

$$\|u\|_{cb} = \sup_{N \geq 1} \|u_N\|.$$

Recall also that, if E, F are completely isomorphic, we set

$$d_{cb}(E, F) = \inf\{\|u\|_{cb} \|u^{-1}\|_{cb}\}$$

where the inf runs over all the complete isomorphisms $u : E \rightarrow F$.

When E, F are both n -dimensional, a compactness argument shows that

$$d_{cb}(E, F) = \sup_{N \geq 1} d_N(E, F).$$

We will apply the preceding to M_N -spaces. When $N = 1$, the latter coincide with the usual Banach spaces. When $N > 1$, roughly the complex scalars are replaced by M_N .

Let $(A_i)_{i \in I}$ be a family of von Neumann or C^* -algebras. Let $Y = \oplus_{i \in I} A_i$ denote their direct sum. This can be described as the algebra of bounded families $(a_i)_{i \in I}$ with $a_i \in A_i$ for all $i \in I$, equipped with the norm $\|a\| = \sup_{i \in I} \|a_i\|$. We will concentrate on the case when $A_i = M_N$ for all $i \in I$. In that case, following the Banach space tradition, we denote the space $Y = \oplus_{i \in I} A_i$ by $\ell_\infty(I; M_N)$.

Definition 2.1. An operator space X is called an M_N -space if, for some set I , it can be embedded completely isometrically in $\ell_\infty(I; M_N)$.

Our main interest will be to try to understand for which spaces the cardinality of I is unusually small.

To place things in perspective, we recall that for any (complex) Banach space X there is an isometric embedding $J : X \rightarrow \ell_\infty(I; \mathbb{C})$ defined by $(Jx)(\phi) = \phi(x)$. Here I is the unit ball, denoted by B_{X^*} , of the space X^* .

In analogy with this, for any M_N -space there is a canonical completely isometric embedding $\hat{J} : X \rightarrow \ell_\infty(\hat{I}; M_N)$ defined again by $(Jx)(\phi) = \phi(x)$, but with $\hat{I} = B_{CB(X, M_N)}$ in place of B_{X^*} . The space $\ell_\infty(\hat{I}; M_N)$ can alternatively be described as $\oplus_{i \in \hat{I}} Z_i$ with $Z_i = M_N$ for all $i \in \hat{I}$.

Just like operator spaces, M_N -spaces enjoy a nice duality theory (see [16, 20] for more information). Indeed, by Roger Smith's lemma, we have $\|u\|_{cb} = \|u_N\|$ for any u with values in an M_N -space (see e.g. [27, p. 26]), and M_N -spaces are characterized among operator spaces by this property. The following reformulation of Smith's Lemma is useful.

Lemma 2.2. Fix an integer $N \geq 1$. Let $E \subset B(H)$ be a finite dimensional operator space and let $c \geq 1$ be a constant. The following properties are equivalent.

- (i) For any operator space F and any $u : F \rightarrow E$ we have $\|u\|_{cb} \leq c\|u_N\|$.
- (ii) There is an M_N -space such that $d_{cb}(E, \hat{E}) \leq c$.
- (iii) Let \mathcal{C} be the class of all (compression) mappings $v : E \rightarrow B(H', H'')$ of the form $x \mapsto P_{H''}x|_{H'}$ where H', H'' are arbitrary subspaces of H of dimension at most N . Let $\hat{J} : E \rightarrow \oplus_{v \in \mathcal{C}} Z_v$ with $Z_v = B(H', H'')$ be defined by $\hat{J}(x) = \oplus_{v \in \mathcal{C}} v(x)$, and let $\hat{E} = \hat{J}(E)$. Then $d_{cb}(E, \hat{E}) \leq c$.

Proof. (ii) \Rightarrow (i) follows from Roger Smith's lemma and (iii) \Rightarrow (ii) is trivial. Conversely, if (i) holds, let \hat{E} be the M_N -space obtained using the embedding $\hat{J} : E \rightarrow \oplus_{v \in \mathcal{C}} Z_v$ appearing in (iii). Obviously $\|E \rightarrow \hat{E}\|_{cb} \leq 1$. Let us denote by $u : \hat{E} \rightarrow E$ the inverse mapping. A simple verification shows that $\|u_N\| = 1$ and hence (i) implies $\|u\|_{cb} \leq c$. In other words (i) \Rightarrow (iii). \square

Therefore, when X is an M_N -space, the knowledge of the space $M_N(X)$ determines that of $M_n(X)$ for all $n > N$, and hence the whole operator space structure of X .

Given a general operator space $X \subset B(H)$, by restricting to $M_N(X)$ (and “forgetting” $M_n(X)$ for $n > N$), we obtain an M_N -space M_N -isometric to X . We will say that the latter M_N -space is induced by X .

Conversely, given an M_N -space X there is a minimal and a maximal operator space structure on X inducing the same M_N -space. When $N = 1$, we recover the Blecher-Paulsen theory of minimal and maximal operator spaces associated to Banach spaces, see [16, 20] for more on this.

Let E be a finite dimensional operator space. For each integer N , let $E[N]$ denote the induced M_N -space. Then it is easy to check that E can be identified (completely isometrically) with the ultraproduct of $\{E[N]\}$ relative to any free ultraproduct on \mathbb{N} . Thus the operator space structure of E can be encoded by the sequence of M_N -spaces $\{E[N] \mid N \geq 1\}$. Note that $E[N]$ is induced by $E[N+1]$ for any N , so that one could picture the set of n -dimensional operator spaces as infinite branches of trees where the N -th node consists of an M_N -space, and any node is induced by any successor.

We can associate to each M_N -space a dual one X^\dagger , isometric to the operator space dual X^* , but defined by

$$\forall n \in \mathbb{N} \quad \forall y \in M_n(X^\dagger) \quad \|y\|_{M_n(X^\dagger)} = \sup_{f \in M_n(X)} \|(I \otimes f)(y)\|_{M_n(M_N)},$$

where we view $M_N(X)$ as a subset of $CB(X^*, M_N)$ in the usual way. In other words we have a completely isometric embedding $J_\dagger : X^\dagger \rightarrow \ell_\infty(I; M_N)$ defined by

$$J_\dagger(z) = \oplus_{f \in M_N(X)} f(z) = \oplus_{f \in M_N(X)} [f_{ij}(z)].$$

Just like for operator spaces, there is a notion of “Hilbert space” for M_N -spaces. We will denote it by $OH(n, N)$. The latter can be defined as follows. First we have an analogue of the Cauchy-Schwarz inequality due to Haagerup, as follows: $\forall x = (x_j) \in M_N^n, \forall y = (y_j) \in M_N^n$

$$(2.1) \quad \left\| \sum x_j \otimes \bar{y}_j \right\| \leq \left\| \sum x_j \otimes \bar{x}_j \right\|^{1/2} \left\| \sum y_j \otimes \bar{y}_j \right\|^{1/2}.$$

Fix N . Let $S(n, N)$ (resp. $B(n, N)$) denote the set of n -tuples $x = (x_j)$ in M_N such that $\left\| \sum x_j \otimes \bar{x}_j \right\| = 1$ (resp. $\left\| \sum x_j \otimes \bar{x}_j \right\| \leq 1$). Then $S(n, N)$ (resp. $B(n, N)$) is the analogue of the unit sphere (resp. ball) in the M_N -space $OH(n, N)$. The space $X = OH(n, N)$ is isometric to ℓ_2^n , with its orthonormal basis (e_j) , and embedded into $\ell_\infty(I; M_N)$ with $I = B(n, N)$ (we could also take $I = S(n, N)$). The embedding $J_{oh} : OH(n, N) \rightarrow \ell_\infty(I; M_N)$ is defined by

$$\forall j = 1, \dots, n \quad J_{oh}(e_j) = \oplus_{x \in B(n, N)} x_j.$$

The latter is the analogue of n -dimensional Hilbert space among M_N -spaces, and indeed when $N = 1$ we recover the n -dimensional Hilbert space.

Definition 2.3. Let E be an operator space with basis (e_j) . Let ξ_j be the biorthogonal basis of E^* . Let $x = \sum x_j \otimes e_j \in M_N(E)$ and $y = \sum y_j \otimes \xi_j \in M_N(E^*)$. Assuming $x \neq 0$ and $y \neq 0$, we say that y M_N -norms x (with respect to $M_N(E)$) if

$$\left\| \sum x_j \otimes y_j \right\| = \|x\|_{M_N(E)} \|y\|_{M_N(E^*)}.$$

In the particular case when $E = OH_n$, we slightly modify this (since $E^* = \bar{E}$): Given $x, y \in M_N(OH_n)$, we say that y M_N -norms x if

$$\left\| \sum x_j \otimes \bar{y}_j \right\| = \left\| \sum x_j \otimes \bar{x}_j \right\|^{1/2} \left\| \sum y_j \otimes \bar{y}_j \right\|^{1/2}.$$

Let $x \in M_N(E)$. For $a, b \in M_N$ we denote by axb the matrix product (i.e. $(a \otimes 1)x(b \otimes 1)$ in tensor product notation using $M_N(E) = M_N \otimes E$). We denote

$$Orb(x) = \{uxv \in M_N(E) \mid u, v \in U(N)\}.$$

Note that if $y \in M_N(E^*)$ M_N -norms x then the same is true for any $y' \in Orb(y) \subset M_N(E^*)$. Actually, any $y' \in Orb(y)$ M_N -norms any $x' \in Orb(x)$.

Definition 2.4. We say that $x \in M_N(E)$ is an M_N -smooth point of $M_N(E)$ if the set of points y in the unit sphere of $M_N(E^*)$ that M_N -norm x is reduced to a single orbit.

The following simple Proposition explains the direction we will be taking next.

Proposition 2.5. Let $x, y \in M_N(OH_n)$. Assume

$$x = (x_j) \in U(N)^n \text{ and } \|T_x : H_0 \rightarrow H_0\| < n,$$

where $T_x = \sum x_j \otimes \bar{x}_j(1 - P)$ (i.e. T_x has a spectral gap at n).

Then y norms x with respect to $M_N(OH_n)$ iff y is a multiple of an element of $Orb(x)$, i.e. iff there are $\lambda > 0$ and $u, v \in U(N)$ such that $y_j = \lambda v x_j u$ for all $1 \leq j \leq n$.

Proof. Recall that whenever the x_j 's are finite dimensional unitaries we have $\|\sum x_j \otimes \bar{x}_j\|^{1/2} = \sqrt{n}$. Assume y is a multiple of an element of $\text{Orb}(x)$, i.e. $y_j = \lambda v x_j u$ for some non zero scalar λ (that may as well be taken positive if we wish). Then $\|\sum x_j \otimes y_j\| = |\lambda|n$, $\|x\|_{M_N(OH_n)} = \sqrt{n}$ and $\|y\|_{M_N(OH_n)} = |\lambda|\sqrt{n}$, so indeed y norms x .

Conversely, assume that y norms x . Multiplying y by a scalar we may assume that $\|y\|_{M_N(OH_n)} = \sqrt{n}$, and $\|\sum x_j \otimes \bar{y}_j\| = \|\sum x_j \otimes \bar{x}_j\|^{1/2} \sqrt{n} = n$. Let ξ, η in the unit sphere of $H = L_2(\tau_n)$ such that

$$\sum \tau_N(x_j \xi y_j^* \eta^*) = n.$$

Let $\xi = u|\xi|$ and $\eta = v|\eta|$ be the polar decompositions, and let $x'_j = v^* x_j u$. Using the trace property, this can be rewritten using the scalar product in H as:

$$\sum \langle (|\eta|^{1/2} x'_j |\xi|^{1/2}), (|\eta|^{1/2} y_j |\xi|^{1/2}) \rangle = n,$$

and hence since $n^{-1/2}(|\eta|^{1/2} x'_j |\xi|^{1/2}), n^{-1/2}(|\eta|^{1/2} y_j |\xi|^{1/2})$ are both in the unit ball of the (smooth!) Hilbert space $\ell_2^n(H)$, they must coincide. Moreover they both must be on the unit sphere. Therefore $\sum \| |\eta|^{1/2} x'_j |\xi|^{1/2} \|_H^2 = n$. Equivalently $\sum \tau_N(x'_j |\xi| x_j^* |\eta|) = n$. But we have obviously $\|T_{x'} : H_0 \rightarrow H_0\| = \|T_x : H_0 \rightarrow H_0\| < n$. Therefore $|\xi|$ and $|\eta|$ must be multiples of I , so that by our normalization we have $|\xi| = |\eta| = I$, and we conclude that $y = x'$. \square

In other words, the preceding Proposition shows that quantum expanders constitute M_N -smooth points of $M_N(OH_n)$:

Corollary 2.6. *Assume $x = (x_j) \in U(N)^n$. Then $x = \sum x_j \otimes e_j$ is an M_N -smooth point in $M_N(OH_n)$ iff $\|T_x : H_0 \rightarrow H_0\| < n$.*

Proof. The “if part” follows from the preceding statement. Conversely, we claim that if $\|T_x : H_0 \rightarrow H_0\| = n$ then x is not an M_N -smooth point in $M_N(OH_n)$. Since this claim is unchanged if we replace x by any x' in $\text{Orb}(x)$, we may assume that $x_1 = 1$. Then if $\|T_x : H_0 \rightarrow H_0\| = n$, there is $0 \neq \xi \in H_0$ such that $\|T_x(\xi)\| = n\|\xi\|$, and hence (by the uniform convexity of Hilbert space) $x_j \xi x_j^* = x_1 \xi x_1^* = \xi$ for all j . This implies that the commutant of $\{x_j\}$ is not reduced to the scalars, and hence in a suitable basis $x_j = x_j^1 \oplus x_j^2 \in M_{N_1} \oplus M_{N_2}$ for some $N_1, N_2 \geq 1$ with $N_1 + N_2 = N$. Then the choice of $y_j = x_j^1 \oplus 0$ produces $y \in M_N^n$ not in $\text{Orb}(x)$ and such that $\|\sum x_j \otimes \bar{y}_j\| = n$. Thus x is not an M_N -smooth point in $M_N(OH_n)$, proving our claim. \square

Remark 2.7. Let E be any n -dimensional operator space with a basis (e_j) . Assume that for any $u = (u_j) \in U(N)^n$ we have $\|\sum u_j \otimes e_j\|_{M_N(E)} = \sqrt{n}$ and also that $\|\sum a_j \otimes e_j\|_{M_N(E)} \leq \|\sum a_j \otimes \bar{a}_j\|^{1/2}$ for all $a = (a_j) \in M_N^n$. Then, by the same proof, for any $x = (x_j) \in U(N)^n$ such that $\|T_x : H_0 \rightarrow H_0\| < n$ as above, the point $x = \sum x_j \otimes e_j$ is an M_N -smooth point in $M_N(E)$. Indeed, any y in the unit ball of $M_N(E^*)$ that M_N -norms x with respect to $M_N(E)$ is a fortiori in the unit ball of $M_N(OH_n)$.

Lemma 1.12 above can be viewed as a refinement of this: assuming $\|T_x : H_0 \rightarrow H_0\| < \varepsilon n$ we have a certain form of “uniform smoothness” of OH_n at x , the points that almost M_N -norm x up to δn are in the orbit of x up to $f_\varepsilon(\delta)n$. See Remark 2.9 for more on this point.

Notation: Let E be a finite dimensional operator space. Fix $C > 0$. We denote by $k_E(N, C)$ the smallest integer k such that there is a subspace F of $M_N \oplus \cdots \oplus M_N$ (with M_N repeated k -times) such that $d_N(E, F) \leq C$.

Note that for any $E \subset M_n$ we have $k_E(N, 1) = 1$ for any $N \geq n$.

The next statement is our main result in this §. It gives a lower bound for $k_E(N, C_1)$ when $E = OH_n$. We will show later (see Lemma 2.11) that a similar upper bound holds for *all* n -dimensional operator spaces. Thus for $E = OH_n$ (and also for $E = \ell_1^n$ or $E = R_n + C_n$, see Remark 2.10) the growth of $N \mapsto k_E(N, C_1)$ is essentially *extremal*.

Theorem 2.8. *There are numbers $C_1 > 1$, $b > 0$, $n_0 > 1$ such that for any $n \geq n_0$ and $N \geq 1$, we have*

$$k_{OH_n}(N, C_1) \geq \exp bnN^2.$$

We start by recalling the classical argument dealing with the Banach space case, i.e. the case $N = 1$. Let E be an n -dimensional Banach space. Assume that, for some $C > 1$, E embeds C -isomorphically into ℓ_∞^k . For convenience we write $C = (1 - \delta)^{-1}$ for some $\delta > 0$. Our embedding assumption means that there is a set \mathcal{T} in the unit ball of E^* such that for any $x \in E$ we have

$$(2.2) \quad (1 - \delta)\|x\| \leq \sup_{t \in \mathcal{T}} |t(x)| \leq \|x\|.$$

Then for any x in the unit ball of E , there is $t_x \in \mathcal{T}$ and $\omega_x \in \mathbb{C}$ with $|\omega_x| = 1$ such that $1 - \delta \leq \Re(\omega_x t_x(x))$.

Now assume $E = \ell_2^n$. Then identifying E and E^* as usual, we see that $1 - \delta \leq \Re(\omega_x t_x(x))$ implies $\|x - \omega_x t_x\|^2 \leq 2\delta$. In the case of real Banach spaces, $\omega_x = \pm 1$ and we conclude quickly, but let us continue for the sake of analogy with the case $N > 1$. We just proved that the set $\{\omega t \mid \omega \in \mathbb{T}, t \in \mathcal{T}\}$ is a $\sqrt{2\delta}$ -net in the unit ball of $E = \ell_2^n$. Fix $\varepsilon > 0$. Let $N(\varepsilon) \approx 2\pi/\varepsilon$ be such that there is an ε -net in \mathbb{T} . It follows that there is a $(\sqrt{2\delta} + \varepsilon)$ -net \mathcal{N} in the unit ball of $E = \ell_2^n$ with $|\mathcal{N}| \leq N(\varepsilon)|\mathcal{T}|$. But by a well known volume estimate (see e.g. [25, p. 49-50]), any δ' -net in the unit ball of $E = \ell_2^n$ must have cardinality at least $(1/\delta')^n$. Thus we conclude $(\sqrt{2\delta} + \varepsilon)^{-n} \leq N(\varepsilon)|\mathcal{T}|$. This yields

$$(2\pi)^{-1}\varepsilon(\sqrt{2\delta} + \varepsilon)^{-n} \leq |\mathcal{T}|.$$

For any $\delta < 1/2$, we may choose $\varepsilon > 0$ so that $\sqrt{2\delta} + \varepsilon < 1$, thus we find that there is a number $b > 0$ for which we obtain $|\mathcal{T}| \geq \exp bn$, and hence $k_{OH_n}(1, (1 - \delta)^{-1}) \geq \exp bn$.

Remark 2.9. The preceding argument still works when E is uniformly convex with modulus $\varepsilon \mapsto \delta(\varepsilon)$. This means that if x_1, x_2 in the unit ball B_E satisfy $\|x_1 - x_2\| \geq \varepsilon$ then $\|(x_1 + x_2)/2\| \leq 1 - \delta(\varepsilon)$. Indeed, the only property we used is that for any $\varepsilon > 0$ there is $r > 0$ such that $x_1, x_2 \in B_E$ and $\xi_1, \xi_2 \in B_{E^*}$ satisfy

$$\Re(\xi_1(x_1)) > 1 - r \quad \Re(\xi_2(x_2)) > 1 - r \quad \text{and} \quad \|\xi_1 - \xi_2\| < r,$$

then we must have $\|x_1 - x_2\| < \varepsilon$. To check this note that

$$\|(x_1 + x_2)/2\| \geq |\xi_1(x_1 + x_2)/2| \geq |\xi_1(x_1)/2 + \xi_2(x_2)/2| - \|\xi_1 - \xi_2\|/2 > 1 - r - r/2$$

thus if $r = \delta(\varepsilon)/2$ then we have $\|(x_1 + x_2)/2\| > 1 - \delta(\varepsilon)$ and hence we must have $\|x_1 - x_2\| < \varepsilon$. Recall that a Banach space E is uniformly convex iff its dual E^* is uniformly smooth (see [4]). Thus since $E = OH_n$ is self dual, Lemma 1.12 can be interpreted as the M_N -analogue of the uniform smoothness of E^* .

A completely different proof, with no restriction on δ or equivalently on the constant C can be given by a well known argument using real or complex Gaussian random variables. We restrict to the real case for simplicity. Let γ_n be the canonical Gaussian measure on \mathbb{R}^n . Assume (2.2). Let $q = \int \exp(x^2/4)\gamma_1(dx) < \infty$. Note that since \mathcal{T} is included in the unit ball we have

$$\int \exp(\sup_{t \in \mathcal{T}} t(x)^2/4)\gamma_n(dx) \leq \sum_{t \in \mathcal{T}} \int \exp(t(x)^2/4)\gamma_n(dx) \leq q|\mathcal{T}|.$$

But by (2.2), if we reset $C = (1 - \delta)^{-1}$, we find $C^{-1}\|x\| \leq \sup_{t \in \mathcal{T}} |t(x)|$ and hence

$$\left(\int \exp(C^{-2}|x|^2/4)\gamma_1(dx) \right)^n \leq \int \exp(C^{-2} \sum |x_j|^2/4)\gamma_n(dx) \leq \int \exp(\sup_{t \in \mathcal{T}} t(x)^2/4)\gamma_n(dx) \leq q|\mathcal{T}|.$$

Thus if we define $b = b_C > 0$ by $\int \exp(C^{-2}|x|^2/4)\gamma_1(dx) = \exp b$, we find $|\mathcal{T}| \geq q^{-1} \exp nb$ and we conclude

$$k_{OH_n}(1, C) \geq q^{-1} \exp b_C n.$$

See [29] for random matrix versions of this argument.

Proof of Theorem 2.8. The proof follows the strategy of the first proof outlined above for $N = 1$, but using Theorem 1.3 instead of the lower bound on the metric entropy of the unit ball of ℓ_2^n . Consider an n -dimensional operator space E . Let $k = k_E(N, C)$. Let again $C = (1 - \delta)^{-1}$. Then there is a set \mathcal{T} with $|\mathcal{T}| = k$ and completely contractive mappings $\phi_t : E \rightarrow M_N$ such that

$$(2.3) \quad \forall x \in M_N(E) \quad (1 - \delta)\|x\|_{M_N(E)} \leq \sup_{t \in \mathcal{T}} \|(\phi_t)_N(x)\|_{M_N(M_N)}.$$

Let e_j be a basis for E so that each x can be developed as $x = \sum x_j \otimes e_j \in M_N \otimes E$. Let $y(t) \in M_N(E^*)$ be the element associated to $\phi_t : E \rightarrow M_N$. Let $e_j^+ \in E^*$ be the basis of E^* that is biorthogonal to (e_j) . Then $y(t)$ (or equivalently ϕ_t) can be written as $y(t) = \sum y_j(t) \otimes e_j^+ \in M_N \otimes E^*$, and (2.3) can be rewritten as:

$$(2.4) \quad \forall x \in M_N(E) \quad (1 - \delta)\|x\|_{M_N(E)} \leq \sup_{t \in \mathcal{T}} \left\| \sum x_j \otimes y_j(t) \right\|_{M_N(M_N)}.$$

Moreover each $y(t)$ is in the unit ball of $M_N(E^*) = CB(E, M_N)$. We now assume $E = OH_n$. Let us denote by $T(\varepsilon, \delta) \subset U(N)^n$ the set appearing in Theorem 1.3. Fix $0 < \varepsilon < 1$. Let us also fix a number $0 < \delta_0 < 1$. By Lemma 1.12 we can choose $0 < \delta < 1$ small enough so that

$$(2.5) \quad 2f_\varepsilon(2\delta) < \sqrt{2\delta_0}.$$

We then set $T_0 = T(\varepsilon, \delta_0)$. Thus we have $|T_0| \geq \exp \beta_0 n N^2$ for some $\beta_0 > 0$ and the elements of T_0 are δ_0 -separated. By (2.4)

$$\forall x = (x_j) \in T_0 \quad (1 - \delta)n^{1/2} \leq \sup_{t \in \mathcal{T}} \left\| \sum x_j \otimes y_j(t) \right\|_{M_N(M_N)}.$$

Let $(v_j(t)) = (n^{1/2} \overline{y_j(t)})$ so that we have

$$\forall x = (x_j) \in T_0 \quad (1 - \delta)n \leq \sup_{t \in \mathcal{T}} \left\| \sum x_j \otimes \overline{v_j(t)} \right\|_{M_N(M_N)}.$$

For any $x \in T_0$ there is a point $t_x \in \mathcal{T}$ such that

$$(1 - \delta)n \leq \left\| \sum x_j \otimes \overline{v_j(t_x)} \right\|.$$

Let $v_x = (v_j(t_x))$. By Lemma 1.12, the last inequality implies $d'(x, v_x) < f_\varepsilon(\delta')\sqrt{n}$ for any $\delta' > \delta$. Moreover by the converse (much easier) part of Lemma 1.11, we know that $d'(x, y) \geq \sqrt{2\delta_0 n}$ for any $x \neq y \in T_0$, since x, y are δ_0 -separated. We claim that after suitably adjusting the parameters δ, ε we have $|T_0| \leq |\mathcal{T}|$. Indeed, assume that $|T_0| > |\mathcal{T}|$, then there must exist $x \neq y \in T_0$ such that $v_x = v_y$. We have then for any $\delta' > \delta$

$$\sqrt{2\delta_0 n} \leq d'(x, y) \leq d'(x, v_x) + d'(v_x, y) = d'(x, v_x) + d'(v_y, y) \leq 2f_\varepsilon(\delta')\sqrt{n}$$

and hence $\sqrt{2\delta_0 n} \leq 2f_\varepsilon(2\delta)$, which is impossible by (2.5). This proves our claim that $|T_0| \leq |\mathcal{T}|$, and hence $|\mathcal{T}| \geq \exp \beta_0 n N^2$. Let $C_1 = (1 - \delta)^{-1}$. Thus, with δ determined by (2.5), we have proved $k_{OH_n}(N, C) \geq \exp \beta_0 n N^2$. \square

Remark 2.10. Let E be any n -dimensional operator space with a basis (e_j) . Assume that there is a scaling factor $\lambda > 0$ (that does not play any role in the estimate) such that for any $u = (u_j) \in U(N)^n$ we have $\lambda \|\sum u_j \otimes e_j\|_{M_N(E)} = \sqrt{n}$ and also that $\lambda \|\sum a_j \otimes e_j\|_{M_N(E)} \leq \|\sum a_j \otimes \bar{a}_j\|^{1/2}$ for all $a = (a_j) \in M_N^n$. Then, arguing as in Remark 2.7, we find $k_E(N, C_1) \geq \exp \beta n N^2$. This shows that this estimate is valid for $R_n + C_n$ (take $\lambda = 1$) and for ℓ_1^n equipped with its maximal operator space structure (take $\lambda = n^{-1/2}$).

We now turn to the reverse inequality to that in Theorem 2.8. This general estimate is easy to check by a rather routine argument.

Lemma 2.11. *Let E be an n -dimensional operator space, then for any $0 < \delta < 1$ we have*

$$k_E(N, (1 - \delta)^{-1}) \leq (1 + 2\delta^{-1})^{2nN^2}.$$

Therefore, for any operator space X , any finite dimensional subspace $E \subset X$ we have

$$\forall C > 1 \quad \limsup_{N \rightarrow \infty} \frac{\log k_E(N, C)}{N^2} < \infty.$$

Proof. Let $x \in M_N(E)$ and let $\hat{x} : E^* \rightarrow M_N$ denote the associated linear mapping. Recall $\|x\| = \|\hat{x}\|_{cb}$. By Lemma 2.2 $\|\hat{x}\|_{cb} = \sup\{\|(\hat{x})_N(y)\|_{M_N(M_N)} \mid y \in B_N\}$ where we denote here by B_N the unit ball of $M_N(E^*)$ viewed as a real space. Since the latter ball is $2nN^2$ -dimensional, it contains a δ -net $\{y_i \mid i \leq m\}$ with cardinality $m \leq (1 + 2\delta^{-1})^{2nN^2}$ (see e.g. [25, p. 49-50]). By an elementary estimate, we have then (for any $x \in M_N(E)$)

$$(2.6) \quad \sup_{i \leq m} \|(\hat{x})_N(y_i)\| \leq \|\hat{x}\|_{cb} = \|x\| \leq (1 - \delta)^{-1} \sup_{i \leq m} \|(\hat{x})_N(y_i)\|.$$

Let $u : E \rightarrow \oplus_{i \leq m} M_N$ be the mapping defined by (here again $\hat{y}_i : E \rightarrow M_N$ is associated to y_i)

$$u(e) = \oplus_{i \leq m} \hat{y}_i(e)$$

for any $e \in E$. Let $F \subset \oplus_{i \leq m} M_N$ be the range of u . Then (2.6) says that $\|u_N\| \leq 1$ and $\|u_N^{-1}\| \leq 1 + \delta$, and hence $d_N(E, F) \leq (1 - \delta)^{-1}$. Thus $k_E(N, (1 - \delta)^{-1}) \leq m$. \square

Definition 2.12. An operator space X will be called *matricially C -subGaussian* if

$$\limsup_{N \rightarrow \infty} \frac{\log k_E(N, C)}{N^2} = 0.$$

for any finite dimensional subspace $E \subset X$. We say that X is *matricially subGaussian* if it is matricially C -subGaussian for some $C \geq 1$. (See Remark 3.2 for the reason behind “matricially”).

Note: If X itself is finite dimensional, it suffices to consider $E = X$.

We will denote by $C_g(X)$ the smallest C such that X is matricially C -subGaussian.

The preceding result (resp. Remark 2.10) shows that when $C < C_1$, then OH (resp. ℓ_1 or $R + C$) is not matricially C -subGaussian. In sharp contrast, any C -exact operator space (we recall the definition below) E is clearly matricially C -subGaussian since, for any $c > C$, it satisfies $k_E(N, c) = 1$ for all N large enough. We do not know whether conversely the latter property implies that E is C -exact (but we doubt it).

Remark 2.13. Given an operator space X , it is natural to introduce the following parameter:

$$k_X(N, C; d) = \sup\{k_E(N, C) \mid E \subset X, \dim(E) = d\}.$$

We will say that X is uniformly matricially subGaussian if there is C such that

$$\forall d \geq 1 \quad \limsup_{N \rightarrow \infty} \frac{\log k_X(N, C; d)}{N} = 0.$$

It is easy to check that if X is uniformly exact (resp. uniformly subexponential, resp. uniformly matricially subGaussian) then all ultrapowers of X are exact (resp. subexponential, resp. matricially subGaussian). Note however (I am indebted to Yanqi Qiu for this remark) that the converse is unclear.

For example, R or C (or $R \oplus C$), any commutative C^* algebra A , or any space of the form $A \otimes_{\min} M_N$ is uniformly exact. It would be interesting to characterize uniformly exact operator spaces.

We now turn to a different application of quantum expanders to operator spaces, that requires a refinement of our main result.

For any $n \times n$ matrix w and any $v \in M_N^n$, we denote by $w.v \in M_N^n$ the n -tuple defined by

$$(w.v)_i = \sum_j w_{ij} v_j.$$

Note that if w is unitary, i.e. $w \in U(n)$ then

$$(2.7) \quad \sum_i (w.v)_i \otimes \overline{(w.v)_i} = \sum_j v_j \otimes \bar{v}_j.$$

Also note that, if $w \in U(n)$, for any $v, v' \in M_N^n$ we have

$$(2.8) \quad d(w.v, w.v') \leq d(v, v').$$

Moreover, it is easy to check (e.g. using (1.1)) that for all $w \in M_n$ with operator norm $\|w\|$ and for all $v \in M_N^n$ we have

$$(2.9) \quad \left\| \sum_i (w.v)_i \otimes \overline{(w.v)_i} \right\| \leq \|w\|^2 \left\| \sum_j v_j \otimes \bar{v}_j \right\|,$$

and hence by (2.1) for any $u, v \in U(N)^n$

$$(2.10) \quad \left\| \sum_i u_i \otimes \overline{(w.v)_i} \right\| \leq \|w\| n.$$

Also

$$d(w.v, w.v') \leq \|w\| d(v, v').$$

We will say that $u, v \in U(N)^n$ are strongly δ -separated if v and $w.u$ are δ -separated for any $w \in U(n)$. Equivalently, for any pair $w, w' \in U(n)$ the pair $(w.u, w'.v)$ is δ -separated.

Explicitly, this can be written like this:

$$(2.11) \quad \forall w \in U(n) \quad \left\| \sum_{ij} w_{ij} u_j \otimes \bar{v}_i \right\| \leq n(1 - \delta).$$

We will use again (see Lemma 1.10) the following elementary fact : There is a positive constant D such that for each $0 < \xi < 1$ and each n there is an ξ -net $\mathcal{N}_\xi \subset U(n)$ with respect to the operator norm, of cardinality

$$|\mathcal{N}_\xi| \leq (D/\xi)^{2n^2}.$$

We will need the following refinement of Theorem 1.3.

Lemma 2.14. *For each $0 < \delta < 1$ there is a constant $\beta'_\delta > 0$ such that for any $0 < \varepsilon < 1$ and for all $n \geq n_0$ and all N such that $N^2/n \geq N_0$ (with n_0 depending on ε and δ , and N_0 depending on δ), there is a strongly δ -separated subset $T_1 \subset S_\varepsilon$ such that $|T_1| \geq \exp \beta'_\delta n N^2$. More generally, for each $\alpha > 0$, there are $\beta'_{\delta, \alpha} > 0$ and $n_0 = n_0(\varepsilon, \delta, \alpha)$ such that, if $n \geq n_0$ and $N^2/n \geq N_0$, any subset $A_N \subset U(N)^n$ with $\mathbb{P}(A_N) > \alpha$ contains a strongly δ -separated subset of S_ε with cardinal $\geq \exp \beta'_{\delta, \alpha} n N^2$.*

Proof. Fix $0 < \delta < 1$ and let $\xi = (1 - \delta)/2$ so that $\delta_1 = \delta + \xi = (1 + \delta)/2$. Note that $0 < \delta < \delta_1 < 1$. We define ε_0 so that $2\varepsilon_0^{1/2} = (1 - \delta_1)/2$ and ε' so that $\varepsilon'^{1/5}(2^{-4/5} + 2^{6/5}) = (1 - \delta_1)/2$. Note that $0 < \varepsilon_0, \varepsilon' < 1$ and

$$1 - \delta_1 = \varepsilon'^{1/5}(2^{-4/5} + 2^{6/5}) + 2\varepsilon_0^{1/2}.$$

Now assume $0 < \varepsilon \leq \varepsilon_0$. By Lemma 1.11 we know that for any $u \in S_\varepsilon$ and any $v \in U(N)^n$ such that $\|\sum v_j \otimes \bar{v}_j\| \leq n$ we have

$$(2.12) \quad \left\| \sum u_j \otimes \bar{v}_j \right\| > n(1 - \delta_1) \Rightarrow d'(u, v) < \sqrt{2n(1 - \varepsilon')}.$$

By Lemmas 1.9 and 1.10 and using (4.6) as in the proof of Theorem 1.3 we know that for $n \geq n_0(\varepsilon, \varepsilon')$

$$N(S_\varepsilon, d', \sqrt{2n(1 - \varepsilon')}) \geq \exp b' n N^2$$

for some b' depending only on δ (more precisely we set again $r = \varepsilon'^2$, $b = K\varepsilon'^2/2$ and $b' = b/2$).

Let $T_1 \subset S_\varepsilon$ be a maximal subset such that any two points in T_1 are strongly δ -separated. By maximality of T_1 for any $u \in S_\varepsilon$ there is $x \in T_1$ such that u, x are not strongly δ -separated. This means that there is $w \in U(n)$ such that

$$\left\| \sum u_j \otimes \overline{(w.x)_j} \right\| > n(1 - \delta).$$

Choose $w' \in \mathcal{N}_\xi$ such that $\|w - w'\| \leq \xi$. Then by (2.10) and the triangle inequality we have

$$\left\| \sum u_j \otimes \overline{(w'.x)_j} \right\| \geq \left\| \sum u_j \otimes \overline{(w.x)_j} \right\| - n\xi > n(1 - \delta - \xi) = 1 - \delta_1.$$

By (2.12) it follows that $d'(u, w'.x) < \sqrt{2n(1 - \varepsilon')}$. In other words, we find that the set $T_2 = \{w'.x \mid w' \in \mathcal{N}_\xi, x \in T_1\}$ is a $\sqrt{2n(1 - \varepsilon')}$ -net for S_ε , and hence

$$\exp b' n N^2 \leq N(S_\varepsilon, d', \sqrt{2n(1 - \varepsilon')}) \leq |T_2| \leq |\mathcal{N}_\xi| |T_1| \leq (D/\xi)^{2n^2} |T_1|$$

This yields

$$|T_1| \geq (2D/(1 - \delta))^{-2n^2} \exp b' n N^2.$$

Assuming $\varepsilon \leq \varepsilon_0$, this completes the proof, since for $N^2/n \geq N_0(\delta)$ the first factor can be absorbed, say, by choosing $\beta'_\delta = b'/2$. The case $\varepsilon_0 < \varepsilon < 1$ follows a fortiori since $S_{\varepsilon_0} \subset S_\varepsilon$.

The last assertion follows (for suitably adjusted values of β'_δ and n_0) as in Remark 1.4. Indeed, choosing n_0 large enough (depending on α) we can make sure that $\mathbb{P}(S_\varepsilon) > 1 - \alpha/2$ so that $\mathbb{P}(A_N \cap S_\varepsilon) > \alpha/2$. We can then run the preceding proof using the set $A_N \cap S_\varepsilon$ in place of S_ε . \square

Theorem 2.15. *For any $R > 1$, there are numbers $\beta_1 > 0$, $n_0 > 1$ and a function $n \mapsto N_0(n)$ from \mathbb{N} to itself such that for any $n \geq n_0$ and $N \geq N_0(n)$, there is a family $\{E_t \mid t \in T_1\}$ of n -dimensional subspaces of M_N , with cardinality $|T_1| \geq \exp \beta_1 n N^2$, such that for any $s \neq t \in T_1$ we have*

$$d_{cb}(E_s, E_t) > R.$$

Proof. Fix $0 < \delta < 1$. We will prove this for $R = (1 - \delta)^{-1}$. We will use the set T_1 from the preceding Lemma and we let $E_t = \text{span}\{t_1, \dots, t_n\}$. We may clearly assume (say by perturbation) that $\{t_1, \dots, t_n\}$ are linearly independent for all $t \in T_1$ so that $\dim(E_t) = n$ (but this will be automatic, see below). Consider $s \neq t \in T_1$. Let $W \in M_n$, and let $W : E_s \rightarrow E_t$ denote the associated linear map so that $Ws_j = \sum_i W_{ij}t_i$.

We claim that we can “make sure” that for all N large enough

$$\text{tr}|W| \leq n(1 - \delta)^{1/2}.$$

We first clarify what we mean here by “ N large enough”. Let $0 < \gamma_1 < 1$ be such that

$$(2.13) \quad (1 - \gamma_1)^{-1}(1 - \delta) = (1 - \delta)^{1/2},$$

let

$$\Delta_{N,n}(t) = n^2 \sup_{i \neq j} |\tau_N(t_i t_j^*)|.$$

Then we require that N is large enough (depending on a fixed n) so that with respect to the uniform probability on $U(N)^n$ we have

$$(2.14) \quad \mathbb{P}\{t \in U(N)^n \mid \Delta_{N,n}(t) < \gamma_1\} > 1/2.$$

Clearly this is possible because, by the almost sure weak convergence, we know that $\tau_N(t_i t_j^*) \rightarrow 0$ when $N \rightarrow \infty$ for any $1 \leq i \neq j \leq n$.

Using the last assertion in the preceding Lemma, we see that we may assume

$$\forall t \in T_1 \quad \Delta_{N,n}(t) < \gamma_1.$$

To verify the above claim, we will use an idea from [20] (refining one in [13]). First we note that for any matrix $a = [a_{ij}]$ and for any $t \in U(N)^n$, if we assume $\tau_N(t_i t_j^*) = 0$ for all $i \neq j$, then we have by (1.1)

$$|\text{tr}(a)| \leq \left\| \sum a_{ij} t_i \otimes \bar{t}_j \right\|.$$

More generally, with the notation from (1.1), we have $\langle \sum a_{ij} t_i \otimes \bar{t}_j(I), I \rangle = \sum a_{ii} + \sum_{i \neq j} a_{ij} \tau_N(t_i t_j^*)$ and $|\sum_{i \neq j} a_{ij} \tau_N(t_i t_j^*)| \leq \Delta_{N,n} \sup_{i \neq j} |a_{ij}|$. Therefore, without this assumption, we still have

$$(2.15) \quad |\text{tr}(a)| \leq \left\| \sum a_{ij} t_i \otimes \bar{t}_j \right\| + \gamma_1 \|a\|_1.$$

By (2.11) and an extreme point argument (since the unitaries are the extreme points of the unit ball of M_n) we have for any $s \neq t \in T_1$ and any $w \in M_n$

$$(2.16) \quad \left\| \sum \bar{w}_{ij} s_i \otimes \bar{t}_j \right\| \leq \|w\| n(1 - \delta).$$

Now we can write for any $W : E_s \rightarrow E_t$ by (2.16)

$$\left\| \sum W s_j \otimes \overline{(w \cdot t)_j} \right\| \leq \|W\|_{cb} \left\| \sum s_j \otimes \overline{(w \cdot t)_j} \right\| \leq \|W\|_{cb} \|w\| n(1 - \delta)$$

Therefore

$$\left\| \sum_{ijk} W_{ij} \overline{w_{jk}} t_i \otimes \bar{t}_k \right\| \leq \|W\|_{cb} \|w\| n(1 - \delta)$$

hence (replacing w by its transpose) by (2.15) we have

$$|\text{tr}(W w^*)| \leq \|W\|_{cb} \|w\| n(1 - \delta) + \gamma_1 \|W w^*\|_1,$$

and hence taking the sup over all $w \in U(n)$

$$\|W\|_1 = \text{tr}|W| \leq \|W\|_{cb}n(1 - \delta) + \gamma_1\|W\|_1.$$

Thus, we conclude by (2.13)

$$(2.17) \quad \text{tr}|W| \leq \|W\|_{cb}n(1 - \gamma_1)^{-1}(1 - \delta) = n\|W\|_{cb}(1 - \delta)^{1/2}.$$

Applying (2.17) with W^{-1} in place of W we find

$$\text{tr}|W^{-1}| \leq n\|W^{-1}\|_{cb}(1 - \delta)^{1/2},$$

and hence

$$\text{tr}|W|\text{tr}|W^{-1}| \leq n^2\|W\|_{cb}\|W^{-1}\|_{cb}(1 - \delta),$$

but we will immediately justify that any invertible matrix in M_n satisfies

$$(2.18) \quad n^2 \leq \text{tr}|W|\text{tr}|W^{-1}|,$$

so that we obtain

$$d_{cb}(E_s, E_t) \geq (1 - \delta)^{-1} = R.$$

To check (2.18) recall that for any pair $W_1, W_2 \in M_n$ the Schatten p -norms $\|\cdot\|_p$ satisfy whenever $0 < p, q, r$ and $1/r = 1/p + 1/q$

$$\|W_1 W_2\|_r \leq \|W_1\|_p \|W_2\|_q.$$

Moreover $\|I\|_r = n^{1/r}$. Therefore, (2.18) follows by taking $r = 1/2$ and $p = q = 1$. \square

3. Random matrices and subexponential operator spaces

In a forthcoming sequel to this paper [29], we introduce and study a generalization of the notion of exact operator space that we call subexponential. We briefly outline this here.

Our goal is to study a generalization of the notion of exact operator space for which the version of Grothendieck's theorem obtained in [31] is still valid.

Notation: Let E be a finite dimensional operator space. Fix $C > 0$. We denote by $K_E(N, C)$ the smallest integer K such that there is an operator subspace $F \subset M_K$ such that

$$d_N(E, F) \leq C.$$

Note that obviously

$$(3.1) \quad K_E(N, C) \leq Nk_E(N, C).$$

Definition 3.1. We say that an operator space X is C -subexponential if

$$\limsup_{N \rightarrow \infty} \frac{\log K_E(N, C)}{N} = 0,$$

for any finite dimensional subspace $E \subset X$. We say that X is subexponential if it is C -subexponential for some $C \geq 1$.

Note: If X itself is finite dimensional, it suffices to consider $E = X$.

We will denote by $C(X)$ the smallest C such that X is C -subexponential.

Recall that an operator space X is called C -exact if for any finite dimensional subspace $E \subset X$ and any $c > C$ there is a k and $F \subset M_k$ such that $d_{cb}(E, F) < c$. We denote by $ex(X)$ the smallest such C . We say that X is exact if it is C -exact for some $C \geq 1$.

We observe in [29] that a finite dimensional E is C -exact iff for any $c > C$ the sequence $N \mapsto K_E(N, c)$ is bounded. In this light “subexponential” seems considerably more general than “exact”.

As shown by Kirchberg, a C^* -algebra is exact iff it is 1-exact. We do not know whether the analogue of this for subexponential (or for matricially subGaussian) C^* -algebras is true. See [27, ch.17] or [5] for more background on exactness.

In [29] we show that for essentially all the results proved in either [13] or [31] we can replace exact by subexponential in the assumptions. Moreover, we show that there is a 1-subexponential C^* -algebra that is not exact.

Remark 3.2. In the same vein, it is natural to call an operator space X C -subGaussian if $\limsup_{N \rightarrow \infty} N^{-2} \log K_E(N, C) = 0$ for any finite dimensional subspace $E \subset X$. We do not have significant information about this class at this point, but to avoid confusion, we decided to call “matricially subGaussian” the spaces in Definition 2.12. Clearly by (3.1) “matricially subGaussian” implies “subGaussian” but the converse is unclear.

Problems:

- 1) Let $C > 1$. Assume that a finite dimensional space E satisfies $k_E(N, C) \leq 1$ for all N . What does that imply on E ? Is E exact with a control on its exactness constant?
- 2) Assume E subexponential for some constant C . What growth does that imply for $N \mapsto k_E(N, C)$ (here C could be a different constant)?
- 3) What is the order of growth (when $N \rightarrow \infty$) of $\log K_E(N, C)$ for $E = \ell_1^n$ or $E = OH_n$? In particular, when C is close to 1, is it $O(N)$? or to the contrary does it grow like N^2 ?

4. Appendix

In this appendix we give a quick proof of an inequality that can be substituted in §2 to Hastings’s result from [11], quoted above as Lemma 1.8. Our inequality is less sharp in some respect but stronger in some other. We only prove that (for some numerical constant C) $\mathbb{P}\{(u_j) \in U(N)^n \mid \|(\sum u_j \otimes \bar{u}_j)(1 - P)\| > 4C\sqrt{n} + \varepsilon n\} \rightarrow 1$ when $N \rightarrow \infty$ for any $\varepsilon > 0$, while Hastings proves this with $2\sqrt{n-1}$ in place of $4C\sqrt{n}$ which is best possible. However the inequality below remains valid with more general (and even matricial) coefficients, and it gives a bound valid uniformly for all sizes N (see (4.6)). It shows that up to a universal constant all moments of the norm of a linear combination of the form

$$S = \sum_j a_j U_j \otimes \bar{U}_j (1 - P)$$

are dominated by those of the corresponding Gaussian sum

$$S' = \sum_j a_j Y_j \otimes \bar{Y}_j'.$$

The advantage is that S' is now simply separately a Gaussian random variable with respect to the independent Gaussian random matrices (Y_j) and (Y_j') .

We recall that we denote by P the orthogonal projection onto the multiples of the identity. Also recall we denote by S_2^N the space M_N equipped with the Hilbert-Schmidt norm (recall $S_2^N \simeq \ell_2^N \otimes_2 \ell_2^N$). We will view elements of the form $\sum x_j \otimes \bar{y}_j$ with $x_j, y_j \in M_N$ as linear operators acting on S_2^N as follows

$$T(\xi) = \sum_j x_j \xi y_j^*,$$

so that

$$(4.1) \quad \left\| \sum x_j \otimes \bar{y}_j \right\| = \|T\|_{B(S_2^N)}.$$

We denote by (U_j) a sequence of i.i.d. random $N \times N$ -matrices uniformly distributed over the unitary group $U(N)$. We will denote by (Y_j) a sequence of i.i.d. Gaussian random $N \times N$ -matrices, more precisely each Y_j is distributed like the variable Y that is such that $\{Y(i, j)N^{1/2}\}$ is a standard family of N^2 independent complex Gaussian variables with mean zero and variance 1. In other words $Y(i, j) = (2N)^{-1/2}(g_{ij} + \sqrt{-1}g'_{ij})$ where g_{ij}, g'_{ij} are independent Gaussian normal $N(0, 1)$ random variables.

We denote by (Y'_j) an independent copy of (Y_j) .

We will denote by $\|\cdot\|_q$ the Schatten q -norm ($1 \leq q \leq \infty$), i.e. $\|x\|_q = (\text{tr}(|x|^q))^{1/q}$, with the usual convention that for $q = \infty$ this is the operator norm.

Lemma 4.1. *There is an absolute constant C such that for any $p \geq 1$ we have for any scalar sequence (a_j) and any $1 \leq q \leq \infty$*

$$\mathbb{E} \left\| \sum_1^n a_j U_j \otimes \bar{U}_j (I - P) \right\|_q^p \leq C^p \mathbb{E} \left\| \sum_1^n a_j Y_j \otimes \bar{Y}'_j \right\|_q^p,$$

(in fact this holds for all k and all matrices $a_j \in M_k$ with $a_j \otimes$ in place of a_j).

Proof. We assume that all three sequences (U_j) , (Y_j) and (Y'_j) are mutually independent. The proof is based on the well known fact that the sequence (Y_j) has the same distribution as $U_j|Y_j|$, or equivalently that the two factors in the polar decomposition $Y_j = U_j|Y_j|$ of Y_j are mutually independent. Let \mathcal{E} denote the conditional expectation operator with respect to the σ -algebra generated by (U_j) . Then we have $U_j \mathbb{E}|Y_j| = \mathcal{E}(U_j|Y_j|) = \mathcal{E}(Y_j)$, and moreover

$$(U_j \otimes \bar{U}_j) \mathbb{E}(|Y_j| \otimes \overline{|Y_j|}) = \mathcal{E}(U_j|Y_j| \otimes \overline{U_j|Y_j|}) = \mathcal{E}(Y_j \otimes \bar{Y}_j).$$

Let

$$T = \mathbb{E}(|Y_j| \otimes \overline{|Y_j|}) = \mathbb{E}(|Y| \otimes \overline{|Y|}).$$

Then we have

$$\sum a_j (U_j \otimes \bar{U}_j) T (I - P) = \mathcal{E}((\sum a_j Y_j \otimes \bar{Y}_j) (I - P)).$$

Note that by rotational invariance of the Gaussian measure we have $(U \otimes \bar{U}) T (U^* \otimes \bar{U}^*) = T$. Indeed since UYU^* and Y have the same distribution it follows that also $UYU^* \otimes \overline{UYU^*}$ and $Y \otimes \bar{Y}$ have the same distribution, and hence so do their modulus.

Viewing T as a linear map on $S_2^N = \ell_2^N \otimes \overline{\ell_2^N}$, this yields

$$\forall U \in U(N) \quad T(U \xi U^*) = U T(\xi) U^*.$$

Representation theory shows that T must be simply a linear combination of P and $I - P$. Indeed, the unitary representation $U \mapsto U \otimes \bar{U}$ on $U(N)$ decomposes into exactly two distinct irreducibles, by restricting either to the subspace $\mathbb{C}I$ or its orthogonal. Thus, by Schur's Lemma we know a priori that there are two scalars χ'_N, χ_N such that $T = \chi'_N P + \chi_N (I - P)$. We may also observe $\mathbb{E}(|Y|^2) = I$ so that $T(I) = I$ and hence $\chi'_N = 1$, therefore

$$T = P + \chi_N (I - P).$$

Moreover, since $T(I) = I$ and T is self-adjoint, T commutes with P and hence $T(I - P) = (I - P)T$, so that we have

$$(4.2) \quad \sum_1^n a_j(U_j \otimes \bar{U}_j)(1 - P)T = \mathcal{E} \sum_1^n a_j(Y_j \otimes \bar{Y}_j)(I - P).$$

We claim that T is invertible and that there is an absolute constant C_0 so that

$$\|T^{-1}\| = \chi_N^{-1} \leq C_0.$$

From this and (4.2) follows immediately that for any $p \geq 1$

$$(4.3) \quad \mathbb{E} \left\| \sum_1^n a_j(U_j \otimes \bar{U}_j)(1 - P) \right\|_q^p \leq C_0^p \mathbb{E} \left\| \sum_1^n a_j(Y_j \otimes \bar{Y}_j)(1 - P) \right\|_q^p.$$

To check the claim it suffices to compute χ_N . For $i \neq j$ we have a priori $T(e_{ij}) = e_{ij} \langle T(e_{ij}), e_{ij} \rangle$ but (since $\text{tr}(e_{ij}) = 0$) we know $T(e_{ij}) = \chi_N e_{ij}$. Therefore for any $i \neq j$ we have $\chi_N = \langle T(e_{ij}), e_{ij} \rangle$, and the latter we can compute:

$$\langle T(e_{ij}), e_{ij} \rangle = \mathbb{E} \text{tr}(|Y|e_{ij}|Y|^*e_{ij}^*) = \mathbb{E}(|Y|_{ii}|Y|_{jj}).$$

Therefore,

$$N(N - 1)\chi_N = \sum_{i \neq j} \mathbb{E}(|Y|_{ii}|Y|_{jj}) = \sum_{i,j} \mathbb{E}(|Y|_{ii}|Y|_{jj}) - \sum_j \mathbb{E}(|Y|_{jj}^2) = \mathbb{E}(\text{tr}|Y|)^2 - N\mathbb{E}(|Y|_{11}^2).$$

Note that $\mathbb{E}(|Y|_{11}^2) = \mathbb{E}(|Y|e_1, e_1)^2 \leq \mathbb{E}(|Y|^2 e_1, e_1) = \mathbb{E}\|Y(e_1)\|_2^2 = 1$, and hence

$$N(N - 1)\chi_N = \sum_{i \neq j} \mathbb{E}(|Y|_{ii}|Y|_{jj}) \geq \mathbb{E}(\text{tr}|Y|)^2 - N.$$

Now it is well known that $E|Y| = b_N I$ where b_N is determined by $b_N = N^{-1} \mathbb{E} \text{tr}|Y| = N^{-1} \|Y\|_1$ and $\inf_N b_N > 0$ (see e.g. [18, p. 80]). Actually, by a well known limit theorem originating in Wigner's work (see [36]), when $N \rightarrow \infty$, $N^{-1} \|Y\|_1$ tends almost surely to the L_1 -norm denoted by $\|c\|_1$ of a circular random variable c normalized in L_2 . Therefore, $N^{-2} \mathbb{E}(\text{tr}|Y|)^2$ tends to $\|c\|_1$. We have

$$\chi_N = (N(N - 1))^{-1} \sum_{i \neq j} \mathbb{E}(|Y|_{ii}|Y|_{jj}) \geq (N(N - 1))^{-1} \mathbb{E}(\text{tr}|Y|)^2 - (N - 1)^{-1},$$

and this implies

$$\liminf_{N \rightarrow \infty} \chi_N \geq (\|c\|_1)^2,$$

and actually $\chi_N \rightarrow (\|c\|_1)^2$. In any case, we have

$$\inf_N \chi_N > 0,$$

proving our claim with $C_0 = (\inf_N \chi_N)^{-1}$.

We will now deduce from (4.3) the desired estimate by a classical decoupling argument for multilinear expressions in Gaussian variables.

We first observe $\mathbb{E}((Y \otimes \bar{Y})(I - P)) = 0$. Indeed, by orthogonality, a simple calculation shows that $\mathbb{E}(Y \otimes \bar{Y}) = \sum_{ij} \mathbb{E}(Y_{ij} \bar{Y}_{ij}) e_{ij} \otimes \bar{e}_{ij} = \sum_{ij} N^{-1} e_{ij} \otimes \bar{e}_{ij} = P$, and hence $\mathbb{E}((Y \otimes \bar{Y})(I - P)) = 0$.

We will use

$$(Y_j, Y'_j) \stackrel{\text{dist}}{=} ((Y_j + Y'_j)/\sqrt{2}, (Y_j - Y'_j)/\sqrt{2})$$

and if \mathbb{E}_Y denotes the conditional expectation with respect to Y we have (recall $\mathbb{E}(Y_j \otimes \bar{Y}_j)(I - P) = 0$)

$$\sum_1^n a_j Y_j \otimes \bar{Y}_j (I - P) = \mathbb{E}_Y \left(\sum_1^n a_j Y_j \otimes \bar{Y}_j (I - P) - \sum_1^n a_j Y'_j \otimes \bar{Y}'_j (I - P) \right).$$

Therefore

$$\begin{aligned}
& \mathbb{E} \left\| \sum_1^n a_j Y_j \otimes \bar{Y}_j (1 - P) \right\|_q^p \leq \mathbb{E} \left\| \sum_1^n a_j Y_j \otimes \bar{Y}_j (1 - P) - \sum_1^n a_j Y'_j \otimes \bar{Y}'_j (I - P) \right\|_q^p \\
& = \mathbb{E} \left\| \sum_1^n a_j (Y_j + Y'_j) / \sqrt{2} \otimes \overline{(Y_j + Y'_j) / \sqrt{2}} (1 - P) - \sum_1^n a_j (Y_j - Y'_j) / \sqrt{2} \otimes \overline{(Y_j - Y'_j) / \sqrt{2}} (I - P) \right\|_q^p \\
& = \mathbb{E} \left\| \sum_1^n a_j (Y_j \otimes \bar{Y}'_j + Y'_j \otimes \bar{Y}_j) (1 - P) \right\|_q^p
\end{aligned}$$

and hence by the triangle inequality

$$\leq 2^p \mathbb{E} \left\| \sum_1^n a_j (Y_j \otimes \bar{Y}'_j) (1 - P) \right\|_q^p.$$

Thus we conclude a fortiori

$$\mathbb{E} \left\| \sum_1^n a_j U_j \otimes \bar{U}_j (1 - P) \right\|_q^p \leq (2C_0)^p \mathbb{E} \left\| \sum_1^n a_j (Y_j \otimes \bar{Y}'_j) \right\|_q^p,$$

so that we can take $C = 2C_0$. □

Theorem 4.2. *Let C be as in the preceding Lemma. Let*

$$\hat{S}^{(N)} = \sum_1^n a_j U_j \otimes \bar{U}_j (1 - P).$$

Then

$$(4.4) \quad \limsup_{N \rightarrow \infty} \mathbb{E} \|\hat{S}^{(N)}\| \leq 4C \left(\sum |a_j|^2 \right)^{1/2}.$$

Moreover we have almost surely

$$(4.5) \quad \limsup_{N \rightarrow \infty} \|\hat{S}^{(N)}\| \leq 4C \left(\sum |a_j|^2 \right)^{1/2}.$$

In addition, there is a constant $C' > 0$ such that for any scalars (a_j)

$$(4.6) \quad \forall N \geq 1 \quad \mathbb{E} \|\hat{S}^{(N)}\| \leq C' \left(\sum |a_j|^2 \right)^{1/2}.$$

Proof. A very direct argument is indicated in Remark 4.5 below, but we prefer to base the proof on [9] in the style of [29] in order to make clear that it remains valid with matrix coefficients. By [29, (1.1)] applied twice (for $k = 1$) (see also Remark 1.5 in [29]) one finds for any even integer p

$$(4.7) \quad \text{Etr} \left| \sum_1^n a_j (Y_j \otimes \bar{Y}'_j) \right|^p \leq (\text{Etr} |Y|^p)^2 \left(\sum |a_j|^2 \right)^{p/2}$$

Therefore by the preceding Lemma

$$\text{Etr} |\hat{S}^{(N)}|^p \leq C^p (\text{Etr} |Y|^p)^2 \left(\sum |a_j|^2 \right)^{p/2},$$

and hence a fortiori

$$E \|\hat{S}^{(N)}\|^p \leq N^2 C^p (\mathbb{E} \|Y\|^p)^2 \left(\sum |a_j|^2 \right)^{p/2}.$$

We then complete the proof, as in [29], using only the concentration of the variable $\|Y\|$. We have an absolute constant β' and $\varepsilon(N) > 0$ tending to zero when $N \rightarrow \infty$, such that

$$(\mathbb{E} \|Y\|^p)^{1/p} \leq 2 + \varepsilon(N) + \beta' \sqrt{p/N},$$

and hence

$$(E\|\hat{S}^{(N)}\|^p)^{1/p} \leq N^{2/p}C(2 + \varepsilon(N) + \beta'\sqrt{p/N})^2(\sum |a_j|^2)^{1/2}.$$

Fix $0 < \varepsilon < 1$. If we choose p minimal even integer so that $N^{2/p} \leq \exp \varepsilon$, i.e. if we set $p = 2([\varepsilon^{-1} \log N] + 1)$ (note that $p > 2\varepsilon^{-1} \log N$ and also $p \geq 2$) we obtain

$$E\|\hat{S}^{(N)}\| \leq (E\|\hat{S}^{(N)}\|^p)^{1/p} \leq 4e^\varepsilon C(1 + \varepsilon^{-1}\varepsilon'(N))(\sum |a_j|^2)^{1/2}$$

where $\varepsilon'(N)$ is independent of ε and satisfies $\varepsilon'(N) \rightarrow 0$ when $N \rightarrow \infty$. Clearly (4.4) and (4.6) follow.

Let $R_N = 4C(1 + \varepsilon^{-1}\varepsilon'(N))(\sum |a_j|^2)^{1/2}$. By Tchebyshev's inequality $(E\|\hat{S}^{(N)}\|^p)^{1/p} \leq e^\varepsilon R_N$ implies

$$\mathbb{P}\{\|\hat{S}^{(N)}\| > e^{2\varepsilon} R_N\} \leq \exp -\varepsilon p = N^{-2}.$$

From this it is immediate that almost surely

$$\limsup_{N \rightarrow \infty} \|\hat{S}^{(N)}\| \leq e^{2\varepsilon} 4C(\sum |a_j|^2)^{1/2}$$

and hence (4.5) follows. \square

Remark 4.3. The same argument can be applied when $a_j \in M_k$ for any integer $k > 1$. Then we find

$$\limsup_{N \rightarrow \infty} \mathbb{E} \left\| \sum_1^n a_j \otimes U_j \otimes \bar{U}_j (1 - P) \right\| \leq 4C \max\{\left\| \sum a_j^* a_j \right\|^{1/2}, \left\| \sum a_j a_j^* \right\|^{1/2}\}.$$

Moreover we have almost surely

$$\limsup_{N \rightarrow \infty} \left\| \sum_1^n a_j \otimes U_j \otimes \bar{U}_j (1 - P) \right\| \leq 4C \max\{\left\| \sum a_j^* a_j \right\|^{1/2}, \left\| \sum a_j a_j^* \right\|^{1/2}\}.$$

Remark 4.4. The preceding also allows us to majorize double sums of the form

$$\sum_{i \neq j} a_{ij} \otimes U_i \otimes \bar{U}_j.$$

Indeed, we have $\mathcal{E}(Y_i \otimes \bar{Y}_j) = (U_i \otimes \bar{U}_j)(\mathbb{E}|Y| \otimes \mathbb{E}|Y|)$ for any $i \neq j$, and there is a constant $b > 0$ (independent of N) such that $\mathbb{E}|Y| \geq bI$. Therefore, for any $p \geq 1$, any k , any sequence (a_{ij}) in M_k , and any $1 \leq q \leq \infty$, we have

$$\mathbb{E} \left\| \sum_{i \neq j} a_{ij} \otimes U_i \otimes \bar{U}_j \right\|_q^p \leq b^{-2p} \mathbb{E} \left\| \sum_{i \neq j} a_{ij} \otimes Y_i \otimes \bar{Y}_j \right\|_q^p \leq 2^p b^{-2p} \mathbb{E} \left\| \sum_{i \neq j} a_{ij} \otimes Y_i \otimes \bar{Y}_j' \right\|_q^p.$$

Remark 4.5. We refer the reader to [28, Theorem 16.6] for a self-contained proof of (4.7) for double sums of the form $\sum_{i,j} a_{ij} Y_i \otimes \bar{Y}_j'$ for scalar coefficients a_{ij} .

Acknowledgment. Thanks to Kate Juschenko for useful conversations at an early stage of this investigation. I am grateful to Mikael de la Salle for simplifying the proof of Lemma 1.11, and to him and Yanqi Qiu for numerous corrections. Thanks also to S. Szarek for a useful suggestion on the appendix.

References

- [1] B. Bekka, P. de la Harpe and A. Valette, Kazhdan's property (T). Cambridge University Press, Cambridge, 2008.
- [2] A. Ben-Aroya and A. Ta-Shma, Quantum expanders and the quantum entropy difference problem, 2007, arXiv:quant-ph/0702129.
- [3] A. Ben-Aroya, O. Schwartz, and A. Ta-Shma, Quantum expanders: motivation and constructions. *Theory Comput.* 6 (2010), 47-79.
- [4] Y. Benyamini and J. Lindenstrauss, Geometric nonlinear functional analysis. Vol. 1. American Mathematical Society, Providence, RI, 2000.
- [5] N.P. Brown and N. Ozawa, *C*-algebras and finite-dimensional approximations*, Graduate Studies in Mathematics, 88, American Mathematical Society, Providence, RI, 2008.
- [6] K. Davidson and S. Szarek, Local operator theory, random matrices and Banach spaces. *Handbook of the geometry of Banach spaces*, Vol. I, 317-366, North-Holland, Amsterdam, 2001.
- [7] J. Elton, Sign-embeddings of ℓ_1^n , *Trans. Amer. Math. Soc.* 279 (1983), 113-124.
- [8] A. Figà-Talamanca and D. Rider, A theorem of Littlewood and lacunary series for compact groups. *Pacific J. Math.* 16 (1966), 505-514.
- [9] U. Haagerup and S. Thorbjørnsen, Random matrices and K -theory for exact C^* -algebras, *Doc. Math.* 4 (1999), 341-450 (electronic).
- [10] P. de la Harpe, A.G. Robertson and A. Valette, On the spectrum of the sum of generators for a finitely generated group. *Israel J. Math.* 81 (1993), 6596.
- [11] M. Hastings, Random unitaries give quantum expanders, *Phys. Rev. A* (3) 76 (2007), no. 3, 032315, 11 pp.
- [12] S. Hoory, N. Linial, and A. Wigderson, Expander graphs and their applications *Bull. Amer. Math. Soc.* 43 (2006), 439-561.
- [13] M. Junge and G. Pisier, Bilinear forms on exact operator spaces and $B(H) \otimes B(H)$, *Geom. Funct. Anal.* 5 (1995), no. 2, 329-363.
- [14] M. Ledoux and M. Talagrand, *Probability in Banach Spaces. Isoperimetry and Processes*, Springer-Verlag, Berlin, 1991.
- [15] M. Ledoux, *The Concentration of Measure Phenomenon*, American Mathematical Society, Providence, RI, 2001.
- [16] F. Lehner, M_n -espaces, sommes d'unitaires et analyse harmonique sur le groupe libre. PhD thesis, Université Paris VI, 1997.
- [17] A. Lubotzky. *Discrete groups, expanding graphs and invariant measures*. Progress in Math. 125. Birkhäuser, 1994.

- [18] M.B. Marcus and G. Pisier, *Random Fourier series with Applications to Harmonic Analysis*. Annals of Math. Studies n°101, Princeton University Press. (1981).
- [19] R. Meshulam and A. Wigderson, Expanders in group algebras. *Combinatorica* 24 (2004), 659-680.
- [20] T. Oikhberg and É. Ricard, Operator spaces with few completely bounded maps. *Math. Ann.*, 328 (2004) 229-259.
- [21] A. Pajor, Sous-espaces ℓ_1^n des espaces de Banach. *Travaux en Cours*, 16. Hermann, Paris, 1985.
- [22] G. Pisier, Remarques sur un résultat non publié de B. Maurey. *Seminar on Functional Analysis*, 1980-1981, Exp. No. V, 13 pp., École Polytech., Palaiseau, 1981. [Available at <http://www.numdam.org/numdam-bin/feuilleter?j=SAF>]
- [23] G. Pisier, De nouvelles caractérisations des ensembles de Sidon. *Advances in Maths. Supplementary studies*, vol 7B (1981) 685-726.
- [24] G. Pisier, Condition d'entropie et caractérisations arithmétiques des ensembles de Sidon. *Modern Topics in Harmonic Analysis - Torino/Milano - June/July 1982*, *Inst. di Alta Math - Rome* (1983) vol. II., 911-944.
- [25] G. Pisier, *The volume of Convex Bodies and Banach Space Geometry*. Cambridge University Press. 1989.
- [26] G. Pisier, The operator Hilbert space OH , complex interpolation and tensor norms, *Mem. Amer. Math. Soc.* **122** (1996), no. 585.
- [27] G. Pisier, *Introduction to operator space theory*, Cambridge University Press, Cambridge, 2003.
- [28] G. Pisier, Grothendieck's Theorem, past and present. *Bull. Amer. Math. Soc.* 49 (2012), 237-323.
- [29] G. Pisier, Random matrices and subexponential operator spaces. *Israel J. Math.* To appear.
- [30] G. Pisier, On the metric entropy of the Banach-Mazur compactum, Preprint, arxiv 2013.
- [31] G. Pisier and D. Shlyakhtenko, Grothendieck's theorem for operator spaces, *Invent. Math.* **150** (2002), no. 1, 185-217.
- [32] O. Regev and T. Vidick, A simple proof of Grothendieck's theorem for completely bounded norms, Preprint 2012, to appear in *J. Op. Theory*.
- [33] S. Szarek, The finite-dimensional basis problem with an appendix on nets of Grassmann manifolds. *Acta Math.* 151 (1983), 153-179.
- [34] S. Szarek Nets of Grassmann manifold and orthogonal group, *Proceedings Research Workshop in Banach space Theory*, Univ. of Iowa, 1981.
- [35] D. Voiculescu, Property T and approximation of operators, *Bull. London Math. Soc.* 22 (1990), 25-30.

- [36] D. Voiculescu, K. Dykema and A. Nica, *Free random variables*, American Mathematical Society, Providence, RI, 1992.
- [37] S. Wassermann, C^* -algebras associated with groups with Kazhdan's property T. Ann. of Math. 134 (1991), 423-431.