

1. Solve the following parabolic initial-boundary-value problem:

$$\begin{cases} \frac{\partial u}{\partial t} - \frac{\partial^2 u}{\partial x^2} = \sin(2\pi x) \sin(4\pi t), & (x, t) \in (0, 1) \times (0, 1), \\ u(0, t) = u(1, t) = 0, & t \in (0, 1], \\ u(x, 0) = 0, & x \in (0, 1), \end{cases}$$

by two-level finite difference schemes with the spatial step-size  $h = 0.025$  and the time steps  $k = 0.05$ . Calculate the solution for three time levels.

2. Solve the following parabolic PDE by using two-level and three-level finite difference schemes:

$$\begin{cases} \frac{\partial u}{\partial t} - \frac{\partial^2 u}{\partial x^2} = 0, & (x, t) \in (0, 1) \times (0, T), \quad T > 0 \\ u(0, t) = u(1, t) = 0, & t \in (0, T], \\ u(x, 0) = f(x), & x \in (0, 1), \end{cases}$$

for the following values of  $f$ :

- (a)  $f(x) = \sin(\pi x)$   
 (b)  $f(x) = x(1 - x)$ .

Here, the values of the step size in the  $x$ -direction is  $\delta x = 0.025$ , and in the  $t$ -direction is  $\delta t = 0.0025$ . Determine the solution for three time levels.

3. Solve the following parabolic problem:

$$\begin{cases} \frac{\partial u}{\partial t} - \frac{\partial^2 u}{\partial x^2} = 0, & (x, t) \in (0, 1) \times (0, t), \quad t > 0, \\ u(x, 0) = \cos(\pi x/2), & x \in (0, 1), \\ u_x(0, t) = 0, & t \geq 0, \\ u(1, t) = 0, & t \geq 0, \end{cases}$$

by using the Neumann boundary condition approximation as given below:

$$u_0^{n+1} = u_0^n + 2\frac{k}{h^2}(u_1^n - u_0^n).$$

Use the spatial step-size  $h = 0.10$ , the time steps  $k = 0.005$  and find the solution at  $t = 0.1$ .

4. Determine the numerical solution of the following one-dimensional parabolic IBVP:

$$\begin{cases} u_t - u_{xx} = 0, & (x, t) \in (0, 1/2) \times (0, T), \quad T > 0 \\ u(0, t) = 0, & u_x(1/2, t) = t^2, \quad \forall t \in (0, T] \\ u(x, 0) = x(1 - x), & x \in (0, 1/2), \end{cases}$$

by using the Crank-Nicolson scheme with  $h = 0.3$  and  $k = 0.05$  for three time levels.

---