

AMSC 714
NUMERICAL METHODS FOR STATIONARY PDE
 HOMEWORK # 2 (Pbs 1-3 due Mar 4, Pbs 4-6 due Mar 13)

1 (15 pts). *L²-interpolation estimate.* Let $u \in H^2(0,1)$, that is u admits two weak derivatives in $L^2(0,1)$. Let u_I be the piecewise linear interpolant of u over a partition $\mathcal{T} = \{T\}$ of $(0,1)$ of size h , i.e. $h = \max_{T \in \mathcal{T}} h_T$. Prove that

$$\|u - u_I\|_{L^2(0,1)} \leq C \left(\sum_{T \in \mathcal{T}} h_T^4 \|u''\|_{L^2(T)}^2 \right)^{\frac{1}{2}} \leq Ch^2 \|u''\|_{L^2(0,1)}.$$

Hint: Show the validity of Poincaré inequality for $v \in H^1(0,1)$ with $v(0) = 0$:

$$\|v\|_{L^2(0,1)} \leq \|v'\|_{L^2(0,1)}.$$

Then use a scaling argument as explained in class and take $v = u - u_I$.

2 (20 pts). *Robin Problem.* Let $\Omega := (0,1)$ and $u \in H^1(\Omega)$ be the solution of HW#1-Pb2 with $h = 1, g = 0$.

- (a) Write a finite element discretization using piecewise linear elements.
- (b) Show that the bilinear form is continuous and coercive in $H^1(\Omega)$. Hint: since the function c can vanish, this entails finding a norm equivalent to the full H^1 -norm that contains a boundary term.
- (c) Prove an error estimate for the error in $H^1(\Omega)$. Specify clearly the required regularity of u and that of the coefficients and forcing function.

3 (20 pts). *Computer implementation.* Write a MATLAB program 2-POINT that implements piecewise linear finite elements over a general mesh $\mathcal{T} = \{x_i\}_{i=0}^N$ for the 2-point boundary value problem of HW#1-Pb6.

- (a) Organize the program in such a way that computations are done elementwise and then assemble to give rise to the stiffness and mass matrices \mathbf{K}, \mathbf{M} and right-hand side \mathbf{F} . Compute the integrals using the midpoint rule.
- (b) Check the rate of convergence in the H^1 -norm and L^∞ -norm over a uniform mesh with meshsize $h = \frac{1}{5}2^{-k}$ for $0 \leq k \leq 5$ and $b = 0, 100$. Plot the H^1 -error and L^∞ -error vs h in a log-log plot. Explain the results.
- (c) Plot the exact solution and the finite element solution for $h = 1/20, 1/80$ and $b = 0, 100$. How do these plots compare with those in HW#1. Draw conclusions.
- (d) Consider a graded mesh \mathcal{T} of the form $x_i = 1 - \left(\frac{N-i}{N}\right)^\beta$ for some $\beta > 1$. Experiment with different values of β and see the consequences for $b = 100$. Try to find a suitable value of β using the principle of error equilibration.

4 (20 pts). *Graded meshes.* Consider $u(x) = \sqrt{x}$ over $I = [0,1]$, and $\mathcal{T} = \{x_j\}_{j=0}^J$. The behavior $u \approx \sqrt{x}$ corresponds to a crack in a two dimensional problem (reentrant corner with internal angle $\omega = 2\pi$).

- (a) Show that

$$\|u - I_{\mathcal{T}}u\|_{L^\infty(x_i, x_{i+1})} = \frac{(\sqrt{x_{i+1}} - \sqrt{x_i})^2}{4(\sqrt{x_{i+1}} + \sqrt{x_i})}.$$

Conclude that $\|u - I_{\mathcal{T}}u\|_{L^\infty(I)} \geq \frac{1}{4\sqrt{N}}$ provided \mathcal{T} is uniform, i.e. $h = h_i$ for all i .

(b) Suppose \mathcal{T} is graded so that $x_i = (\frac{i}{N})^4$. Show that

$$\|u - I_{\mathcal{T}}u\|_{L^\infty(x_i, x_{i+1})} = \frac{1}{4N^2} \left(2 - \frac{1}{i^2 + (i-1)^2} \right).$$

Conclude that the global maximum error is $\leq \frac{1}{2N^2}$, which is the best approximation possible with $N + 1$ points. Compare with (a) and draw conclusions.

5 (15 pts). *Quadrature.* Let $\mathcal{T} = \{x_i\}_{i=0}^N$ be a partition of $\Omega = (0, 1)$. Let $Q(w) = \sum_{i=1}^N Q_i(w)$ be the trapezoidal quadrature rule, where

$$Q_i(w) := \frac{h_i}{2} (w(x_{i-1}) + w(x_i))$$

and $h_i = x_i - x_{i-1}$ is the local meshsize. Show that for all $w \in W_1^2(\Omega)$ the following error estimate holds

$$\left| Q(w) - \int_{\Omega} w \right| \leq C \sum_{i=1}^N h_i^2 \int_{x_{i-1}}^{x_i} |w''|.$$

Hint: Use the fact that Q is exact for piecewise linear functions and apply the Bramble-Hilbert lemma.

6 (10 pts). *Advection-diffusion equation.* Consider the HW#1-Pb7, namely

$$-\epsilon u'' + u' = 0, \quad u(0) = u(1) = 0.$$

(a) Write the finite element discretization with piecewise linear elements over a uniform partition with meshsize h . Compare with the centered finite difference method of HW#1-Pb7.

(b) Consider the *modified* equation

$$-\left(\epsilon + \frac{h}{2}\right)u'' + u' = 0.$$

Write the corresponding finite element discretization. Compare with the upwind finite difference method of HW#1-Pb7.

(c) The *streamline diffusion method* is a Petrov-Galerkin method that utilizes test functions different from the trial ones. In 1D it reads

$$U \in \mathbb{V}_h : \quad B[U, V] = \int_0^1 \epsilon U' V' + U' \left(V + \frac{h}{2} V' \right) = 0 \quad \forall V \in V_h.$$

Show that this is equivalent to approximating the modified equation with a Galerkin method.