

# Multivariable Calculus (CS+AI)

Aron Hardeman

June 14, 2023

## 1 Triple integrals

# Triple integrals

## Observation:

Triple integrals have appeared in the homework, but not in past exams (at least not in the ones found on Cover).

The coming slides discuss triple integrals.

(I'm not saying you won't get a triple integral on your exam...)


# Triple integrals

## Observation:

Triple integrals have appeared in the homework, but not in past exams (at least not in the ones found on Cover).

The coming slides discuss triple integrals.

(I'm not saying you won't get a triple integral on your exam...)



# Triple integrals

One can also have a triple integral:  $\iiint_E f(x, y, z) dV$

# Triple integrals

One can also have a triple integral:  $\iiint_E f(x, y, z) dV$

When the region of integration is a box  $E = [a, b] \times [c, d] \times [r, s]$ , then:

$$\iiint_E f(x, y, z) dV = \int_a^b \int_c^d \int_r^s f(x, y, z) dz dy dx$$

(all 6 orders of integration are possible, in case of a box, since the bounds of the variables do not depend on each other)

This is an integral that can be solved with methods similar to the ones from double integrals.

# Triple integrals

One can also have a triple integral:  $\iiint_E f(x, y, z) dV$

When the region of integration is a box  $E = [a, b] \times [c, d] \times [r, s]$ , then:

$$\iiint_E f(x, y, z) dV = \int_a^b \int_c^d \int_r^s f(x, y, z) dz dy dx$$

(all 6 orders of integration are possible, in case of a box, since the bounds of the variables do not depend on each other)

This is an integral that can be solved with methods similar to the ones from double integrals.

We can also take triple integrals over general regions. For example:

$$E = \{(x, y, z) \mid 0 \leq y \leq 3, 0 \leq x \leq y^2, 0 \leq z \leq xy + 1\}$$

$$\Rightarrow \iiint_E f(x, y, z) dV = \int_0^3 \int_0^{y^2} \int_0^{xy+1} f(x, y, z) dz dx dy$$

# Example triple integral

**Question:** evaluate  $\int_0^3 \int_0^{z^2} \int_0^{y-z} (3x - 2y) \, dx \, dy \, dz$ .

**Solution:**



# Example triple integral

**Question:** evaluate  $\int_0^3 \int_0^{z^2} \int_0^{y-z} (3x - 2y) \, dx \, dy \, dz$ .

**Solution:**

$$\int_0^3 \int_0^{z^2} \int_0^{y-z} (3x - 2y) \, dx \, dy \, dz = \int_0^3 \int_0^{z^2} \left[ \frac{3}{2}x^2 - 2xy \right]_{x=0}^{x=y-z} dy \, dz$$

# Example triple integral

**Question:** evaluate  $\int_0^3 \int_0^{z^2} \int_0^{y-z} (3x - 2y) dx dy dz$ .

**Solution:**

$$\begin{aligned} \int_0^3 \int_0^{z^2} \int_0^{y-z} (3x - 2y) dx dy dz &= \int_0^3 \int_0^{z^2} \left[ \frac{3}{2}x^2 - 2xy \right]_{x=0}^{x=y-z} dy dz \\ &= \int_0^3 \int_0^{z^2} \left( \frac{3}{2}(y-z)^2 - 2(y-z)y \right) dy dz \end{aligned}$$

# Example triple integral

**Question:** evaluate  $\int_0^3 \int_0^{z^2} \int_0^{y-z} (3x - 2y) dx dy dz$ .

**Solution:**

$$\begin{aligned} \int_0^3 \int_0^{z^2} \int_0^{y-z} (3x - 2y) dx dy dz &= \int_0^3 \int_0^{z^2} \left[ \frac{3}{2}x^2 - 2xy \right]_{x=0}^{x=y-z} dy dz \\ &= \int_0^3 \int_0^{z^2} \left( \frac{3}{2}(y-z)^2 - 2(y-z)y \right) dy dz \\ &= \int_0^3 \int_0^{z^2} \left( -\frac{1}{2}y^2 - zy + \frac{3}{2}z^2 \right) dy dz \end{aligned}$$

# Example triple integral

**Question:** evaluate  $\int_0^3 \int_0^{z^2} \int_0^{y-z} (3x - 2y) dx dy dz$ .

**Solution:**

$$\begin{aligned} \int_0^3 \int_0^{z^2} \int_0^{y-z} (3x - 2y) dx dy dz &= \int_0^3 \int_0^{z^2} \left[ \frac{3}{2}x^2 - 2xy \right]_{x=0}^{x=y-z} dy dz \\ &= \int_0^3 \int_0^{z^2} \left( \frac{3}{2}(y-z)^2 - 2(y-z)y \right) dy dz \\ &= \int_0^3 \int_0^{z^2} \left( -\frac{1}{2}y^2 - zy + \frac{3}{2}z^2 \right) dy dz \\ &= \int_0^3 \left[ -\frac{1}{6}y^3 - \frac{1}{2}zy^2 + \frac{3}{2}z^2y \right]_{y=0}^{y=z^2} dz \end{aligned}$$

# Example triple integral

**Question:** evaluate  $\int_0^3 \int_0^{z^2} \int_0^{y-z} (3x - 2y) dx dy dz$ .

**Solution:**

$$\begin{aligned} \int_0^3 \int_0^{z^2} \int_0^{y-z} (3x - 2y) dx dy dz &= \int_0^3 \int_0^{z^2} \left[ \frac{3}{2}x^2 - 2xy \right]_{x=0}^{x=y-z} dy dz \\ &= \int_0^3 \int_0^{z^2} \left( \frac{3}{2}(y-z)^2 - 2(y-z)y \right) dy dz \\ &= \int_0^3 \int_0^{z^2} \left( -\frac{1}{2}y^2 - zy + \frac{3}{2}z^2 \right) dy dz \\ &= \int_0^3 \left[ -\frac{1}{6}y^3 - \frac{1}{2}zy^2 + \frac{3}{2}z^2y \right]_{y=0}^{y=z^2} dz \\ &= \int_0^3 \left( -\frac{1}{6}z^6 - \frac{1}{2}z^5 + \frac{3}{2}z^4 \right) dz \end{aligned}$$

# Example triple integral

**Question:** evaluate  $\int_0^3 \int_0^{z^2} \int_0^{y-z} (3x - 2y) dx dy dz$ .

**Solution:**


$$\begin{aligned} \int_0^3 \int_0^{z^2} \int_0^{y-z} (3x - 2y) dx dy dz &= \int_0^3 \int_0^{z^2} \left[ \frac{3}{2}x^2 - 2xy \right]_{x=0}^{x=y-z} dy dz \\ &= \int_0^3 \int_0^{z^2} \left( \frac{3}{2}(y-z)^2 - 2(y-z)y \right) dy dz \\ &= \int_0^3 \int_0^{z^2} \left( -\frac{1}{2}y^2 - zy + \frac{3}{2}z^2 \right) dy dz \\ &= \int_0^3 \left[ -\frac{1}{6}y^3 - \frac{1}{2}zy^2 + \frac{3}{2}z^2y \right]_{y=0}^{y=z^2} dz \\ &= \int_0^3 \left( -\frac{1}{6}z^6 - \frac{1}{2}z^5 + \frac{3}{2}z^4 \right) dz \\ &= \left[ -\frac{1}{42}z^7 - \frac{1}{12}z^6 + \frac{3}{10}z^5 \right]_0^3 = \boxed{-\frac{5589}{140}} \end{aligned}$$

# Spherical coordinates

In the 2D world, we have polar coordinates.

# Spherical coordinates

In the 2D world, we have polar coordinates. In 3D, we have **spherical coordinates**  $(\rho, \theta, \phi)$ . They look like this:



$\rho$  (rho) is the radial distance,  $\theta$  (theta) is the *azimuthal angle*, and  $\phi$  (phi) is the *polar angle*.



# Integration in spherical coordinates

In the 2D world, we have polar coordinates, where  $dA = r \cdot dr d\theta$ .

# Integration in spherical coordinates

In the 2D world, we have polar coordinates, where  $dA = r \cdot dr d\theta$ .

In 3D's spherical coordinates, we have  $dV = \rho^2 \sin \phi \cdot dr d\theta d\phi$ .

(The blue factors are Jacobians, if you want to know more about them)

# Integration in spherical coordinates

In the 2D world, we have polar coordinates, where  $dA = r \cdot dr d\theta$ .

In 3D's spherical coordinates, we have  $dV = \rho^2 \sin \phi \cdot dr d\theta d\phi$ .

(The blue factors are Jacobians, if you want to know more about them)

For spherical coordinates, we have:

$$x = \rho \sin \phi \cos \theta \quad y = \rho \sin \phi \sin \theta \quad z = \rho \cos \phi$$

$$x^2 + y^2 + z^2 = \rho^2$$

# Integration in spherical coordinates

In the 2D world, we have polar coordinates, where  $dA = r \cdot dr d\theta$ .

In 3D's spherical coordinates, we have  $dV = \rho^2 \sin \phi \cdot dr d\theta d\phi$ .

(The blue factors are Jacobians, if you want to know more about them)

For spherical coordinates, we have:

$$x = \rho \sin \phi \cos \theta \quad y = \rho \sin \phi \sin \theta \quad z = \rho \cos \phi$$

$$x^2 + y^2 + z^2 = \rho^2$$

**Note:** the slides use the convention of the book, where  $\rho$  is the radial distance,  $\theta$  is the azimuthal angle and  $\phi$  is the polar angle. However, some sources swap the meanings of  $\theta$  and  $\phi$  and/or write  $r$  instead of  $\rho$ , so be aware of that.

# Example integral in spherical coordinates (1/2)

**Question:** evaluate  $\iiint_E x e^{x^2+y^2+z^2} dV$ , where  $E$  is the region with  $x^2 + y^2 + z^2 \leq 4$  and  $0 \leq y \leq x$ .

# Example integral in spherical coordinates (1/2)

**Question:** evaluate  $\iiint_E x e^{x^2+y^2+z^2} dV$ , where  $E$  is the region with  $x^2 + y^2 + z^2 \leq 4$  and  $0 \leq y \leq x$ .

**Step 1:** do geometry; write  $E$  in spherical coordinates:

$$E = \left\{ (\rho, \theta, \phi) \mid 0 \leq \rho \leq 2, 0 \leq \theta \leq \frac{\pi}{4}, -\frac{\pi}{2} \leq \phi \leq \frac{\pi}{2} \right\}$$

# Example integral in spherical coordinates (1/2)

**Question:** evaluate  $\iiint_E x e^{x^2+y^2+z^2} dV$ , where  $E$  is the region with  $x^2 + y^2 + z^2 \leq 4$  and  $0 \leq y \leq x$ .

**Step 1:** do geometry; write  $E$  in spherical coordinates:

$$E = \left\{ (\rho, \theta, \phi) \mid 0 \leq \rho \leq 2, 0 \leq \theta \leq \frac{\pi}{4}, -\frac{\pi}{2} \leq \phi \leq \frac{\pi}{2} \right\}$$

**Step 2:** since  $x = \rho \sin \phi \cos \theta$  and  $x^2 + y^2 + z^2 = \rho^2$  in spherical coordinates, we can rewrite the integrand as  $\rho \sin \phi \cos \theta e^{\rho^2}$ .

# Example integral in spherical coordinates (1/2)

**Question:** evaluate  $\iiint_E x e^{x^2+y^2+z^2} dV$ , where  $E$  is the region with  $x^2 + y^2 + z^2 \leq 4$  and  $0 \leq y \leq x$ .

**Step 1:** do geometry; write  $E$  in spherical coordinates:

$$E = \left\{ (\rho, \theta, \phi) \mid 0 \leq \rho \leq 2, 0 \leq \theta \leq \frac{\pi}{4}, -\frac{\pi}{2} \leq \phi \leq \frac{\pi}{2} \right\}$$

**Step 2:** since  $x = \rho \sin \phi \cos \theta$  and  $x^2 + y^2 + z^2 = \rho^2$  in spherical coordinates, we can rewrite the integrand as  $\rho \sin \phi \cos \theta e^{\rho^2}$ .

**Step 3:** set up the integral. Do not forget the Jacobian  $\rho^2 \sin \phi$  for spherical coordinates!

$$\begin{aligned} \iiint_E x e^{x^2+y^2+z^2} dV &= \int_{-\pi/2}^{\pi/2} \int_0^{\pi/4} \int_0^2 \rho \sin \phi \cos \theta e^{\rho^2} \rho^2 \sin \phi d\rho d\theta d\phi \\ &= \left( \int_{-\pi/2}^{\pi/2} \sin^2 \phi d\phi \right) \left( \int_0^{\pi/4} \cos \theta d\theta \right) \left( \int_0^2 \rho^3 e^{\rho^2} d\rho \right) \end{aligned}$$

We were able to write the long integral as a product of three single-variable integrals by using the idea from slide ??.



## Example integral in spherical coordinates (2/2)

**Step 4:** solve the integral.

$$\iiint_E x e^{x^2+y^2+z^2} dV = \left( \int_{-\pi/2}^{\pi/2} \sin^2 \phi d\phi \right) \left( \int_0^{\pi/4} \cos \theta d\theta \right) \left( \int_0^2 \rho^3 e^{\rho^2} d\rho \right)$$

## Example integral in spherical coordinates (2/2)

**Step 4:** solve the integral.

$$\iiint_E x e^{x^2+y^2+z^2} dV = \left( \int_{-\pi/2}^{\pi/2} \sin^2 \phi d\phi \right) \left( \int_0^{\pi/4} \cos \theta d\theta \right) \left( \int_0^2 \rho^3 e^{\rho^2} d\rho \right)$$

The red one can be solved by subbing  $u = \rho^2$  (such that  $du = 2\rho d\rho$ ), followed by integration by parts:

$$\int_0^2 \rho^3 e^{\rho^2} d\rho = \frac{1}{2} \int_0^2 u e^u du = \frac{1}{2} \left( [u e^u]_0^2 - \int_0^2 e^u du \right) = \frac{1}{2} (4e^4 - (e^4 - 1)) = \frac{3e^4 + 1}{2}$$

## Example integral in spherical coordinates (2/2)

**Step 4:** solve the integral.

$$\iiint_E x e^{x^2+y^2+z^2} dV = \left( \int_{-\pi/2}^{\pi/2} \sin^2 \phi d\phi \right) \left( \int_0^{\pi/4} \cos \theta d\theta \right) \left( \int_0^2 \rho^3 e^{\rho^2} d\rho \right)$$

The red one can be solved by subbing  $u = \rho^2$  (such that  $du = 2\rho d\rho$ ), followed by integration by parts:

$$\int_0^2 \rho^3 e^{\rho^2} d\rho = \frac{1}{2} \int_0^2 u e^u du = \frac{1}{2} \left( [u e^u]_0^2 - \int_0^2 e^u du \right) = \frac{1}{2} (4e^4 - (e^4 - 1)) = \frac{3e^4 + 1}{2}$$

The green one can be solved by using  $\sin^2 \phi = \frac{1}{2} (1 - \cos 2\phi)$ :

$$\int_{-\pi/2}^{\pi/2} \sin^2 \phi d\phi = \frac{1}{2} \int_{-\pi/2}^{\pi/2} (1 - \cos 2\phi) d\phi = \frac{\pi}{2} - \frac{1}{2} \left[ \frac{1}{2} \sin 2\phi \right]_{-\pi/2}^{\pi/2} = \frac{\pi}{2}$$

## Example integral in spherical coordinates (2/2)

**Step 4:** solve the integral.

$$\iiint_E x e^{x^2+y^2+z^2} dV = \left( \int_{-\pi/2}^{\pi/2} \sin^2 \phi d\phi \right) \left( \int_0^{\pi/4} \cos \theta d\theta \right) \left( \int_0^2 \rho^3 e^{\rho^2} d\rho \right)$$

The red one can be solved by subbing  $u = \rho^2$  (such that  $du = 2\rho d\rho$ ), followed by integration by parts:

$$\int_0^2 \rho^3 e^{\rho^2} d\rho = \frac{1}{2} \int_{0^2}^{2^2} u e^u du = \frac{1}{2} \left( [u e^u]_0^4 - \int_0^4 e^u du \right) = \frac{1}{2} (4e^4 - (e^4 - 1)) = \frac{3e^4 + 1}{2}$$

The green one can be solved by using  $\sin^2 \phi = \frac{1}{2} (1 - \cos 2\phi)$ :

$$\int_{-\pi/2}^{\pi/2} \sin^2 \phi d\phi = \frac{1}{2} \int_{-\pi/2}^{\pi/2} (1 - \cos 2\phi) d\phi = \frac{\pi}{2} - \frac{1}{2} \left[ \frac{1}{2} \sin 2\phi \right]_{-\pi/2}^{\pi/2} = \frac{\pi}{2}$$

The orange one is relatively straightforward, so the answer is:

$$\iiint_E x e^{x^2+y^2+z^2} dV = \left( \frac{\pi}{2} \right) \left( \frac{1}{2} \sqrt{2} \right) \left( \frac{3e^4 + 1}{2} \right) = \boxed{\frac{\pi\sqrt{2}}{8} (3e^4 + 1)}$$

P.S. The need to use substitution, integration by parts and a trigonometric identity makes this question harder than exam-level (no warranty).