

Genome-scale analysis of small molecule regulatory networks reveals principles of metabolic regulation - Supplementary Text

March 15, 2017

Contents

1	Futile Cycles and Small Molecule Regulation	2
2	Metabolic Control Analysis	2
2.1	What is elasticity?	2
2.2	Single-substrate Control in Michaelis-Menten Kinetics	3
2.3	Regulatory Effectors	4
2.3.1	Activation	5
2.3.2	Pure non-Competitive Inhibition	5
2.3.3	Competitive Inhibition	6
2.3.4	Uncompetitive Inhibition	7
2.3.5	Generalized Inhibition Model	8
2.3.6	The Monod-Changeux-Jacob model	9
2.4	A Trade-off between Enzyme Cost and Flux Control	10
3	Supplementary Figures	11
	References	11

1 Futile Cycles and Small Molecule Regulation

We gathered information regarding reactions that are involved in futile cycling (dissipation of ATP), using the *E. coli* genome scale model **iJO1366**. We mapped these reactions to enzymes, using EC number information (where available) and ended up with 75 unique reactions that take part in futile cycles (`FutileCycleReactionsECnumber.csv`). From these, using our SMRN, we find that 37 are regulated by at least one small molecule, whereas 38 are not regulated (`FutileCycleRegulatoryInfo.csv`).

	In Futile Cycle	Not in Futile Cycle
Regulated by small molecule	37	327
Not regulated	38	266

Using a right-tailed Fischer's exact test (`fishertest` function, MATLAB) to detect if there are significantly more small molecule regulation in futile cycle participating reactions, we verify that there is no significant overrepresentation of regulation in reactions that take part in futile cycling ($p = 0.86$).

2 Metabolic Control Analysis

2.1 What is elasticity?

Metabolic Control Analysis [1] is a mathematical framework to understanding the dynamics and control of metabolic networks. In particular, it defines local properties called elasticities that can be used to quantify how the control of metabolic fluxes depends on other quantities in the system (*i.e.* the topology of the metabolic network, and the levels of metabolites). *Scaled elasticity* is defined as the infinitesimal response of a single flux (v) to one of the parameters (a), using a partial derivative of the log-scaled functions:

$$\epsilon_a^v \equiv \frac{\partial \ln(v)}{\partial \ln(a)} = \frac{\partial v}{\partial a} \frac{a}{v} \quad (1)$$


The elasticity has several properties which make it desirable for modeling metabolic systems. In particular, for a variety of kinetic rate laws (*e.g.* Michaelis-Menten kinetics), the elasticity (for Michaelis-Menten kinetics, the substrate elasticity is bounded between 0 and 1).

2.2 Single-substrate Control in Michaelis-Menten Kinetics

To illustrate the concept of elasticity, let us consider an enzyme-catalyzed reaction, described by a one-substrate irreversible Michaelis-Menten kinetic rate law:


$$v = V^+ \frac{s}{K_M + s} \quad (2)$$

where s is the substrate concentrations (in units of molar), and K_M is the Michaelis-Menten coefficient (also in molar).



The scaled elasticity for s is given by:

$$\begin{aligned} \epsilon_s^v &= \frac{\partial v}{\partial s} \frac{s}{v} = V^+ \frac{K_M + s - s}{(K_M + s)^2} \frac{s}{v} \\ &= \frac{K_M}{(K_M + s)^2} (K_M + s) = \frac{K_M}{K_M + s} = 1 - \frac{s}{K_M + s} \end{aligned} \quad (3)$$



Note that the scaled elasticity is bounded between 0 and 1. More specifically, ϵ_s^v is maximized (*i.e.* the substrate has the highest control potential) at low substrate concentrations, and minimized at high substrate concentrations.

As shown in [2], this formula for substrate elasticity can be generalized to reversible Michaelis-Menten reactions, where the saturation term is separated from the thermodynamic term:

$$v = V^+ \cdot \kappa \cdot \gamma \quad (4)$$

$$\kappa \equiv \frac{s/K_S}{1 + s/K_S + p/K_P} \quad (5)$$

$$\gamma \equiv 1 - \frac{p/s}{K'_{eq}} \quad (6)$$

where p is the product concentration, K_S and K_P are the Michaelis-Menten constants for the substrate and product, and K'_{eq} is the apparent equilibrium constant. In this case, the elasticity of the substrate is:

$$\epsilon_s^v = \gamma^{-1} - \kappa \quad (7)$$

which converges to equation 3 as $p \rightarrow 0$.

2.3 Regulatory Effectors

Although previous publications have derived elasticities associated with small-molecule effectors for different types of rate laws [3, 4], the relationship between the elasticity and the relative activity of the enzyme has not been discussed. Here, we will demonstrate that in almost all cases, *there is a direct trade-off between the two, namely that a regulator must decrease the activity of the enzyme in order to have a non-zero elasticity.*

Without loss of generality, we will keep the separable form of the rate law

$$v = V^+ \cdot \kappa \cdot \gamma \cdot \theta(x)$$

where we add a multiplicative term $\theta(x)$ that will represent the decrease of activity due to the small-molecule regulation (x). As long as $\theta(x)$ is the only term affected by x , *i.e.* $\frac{\partial \kappa}{\partial x} = \frac{\partial \gamma}{\partial x} = 0$, the exact forms of κ and γ are irrelevant.

2.3.1 Activation

First, consider a cooperative [5, 6] activator with Hill coefficient h and activation coefficient K_A :

$$\theta = \frac{a^h}{K_A^h + a^h}. \quad (8)$$

The elasticity with respect to the activator concentration x will thus be:

$$\begin{aligned} \epsilon_a^v &= \frac{\partial v}{\partial a} \frac{a}{v} = V^+ \kappa \gamma \frac{h a^{h-1}(K_A^h + a^h) - h a^{h-1} a^h}{(K_A^h + a^h)^2} \frac{a}{v} \\ &= h \frac{K_A^h}{K_A^h + a^h} = h(1 - \theta) \end{aligned} \quad (9)$$


[to make this even more general, we can also mention the case of an activator that increases an already existing activity: instead of θ , we get $\alpha + (1 - \alpha)\theta$ with a constant relative baseline activity α . In this case, the elasticity formula remains the same, except for a constant prefactor; the same logic can also be applied to inhibitors.]

2.3.2 Pure non-Competitive Inhibition

Next, we consider a pure non-competitive inhibitor with Hill coefficient h and inhibition coefficient K_I :

$$\theta = 1 - \frac{x^h}{K_I^h + x^h} = \frac{K_I^h}{K_I^h + x^h} \quad (10)$$


The following plot shows the response of θ to the concentration x in log-log scale.



In this case, the elasticity with regards to the inhibitor concentration would be:

$$\begin{aligned}\epsilon_x^v &= \frac{\partial v}{\partial x} \cdot \frac{x}{v} = V^+ \kappa \gamma \frac{-h x^{h-1}}{(K_I^h + x^h)^2} \frac{x}{v} \\ &= -\frac{h x^h (K_I^h + x^h)}{(K_I^h + x^h)^2} = -h \frac{x^h}{K_I^h + x^h} = -h(1 - \theta).\end{aligned}\quad (11)$$

Plotting the elasticity as a function of x , we see that it is a monotonically decreasing negative function:



This means that substrates have the most control ($|\epsilon_s^v| \rightarrow 1$) when they are much below saturation ($s \ll K_M$) while inhibitors have the most control ($|\epsilon_x^v| \rightarrow h$) when they are saturated ($x \gg K_I$).

2.3.3 Competitive Inhibition

Competitive inhibition is one case, where a separable kinetic rate law is not sufficient since $\frac{\partial \kappa}{\partial x} \neq 0$. To analyze this case, we pick a simple one-substrate irreversible reaction, where the inhibitor affects the K_M according to the following formula:

$$v = V^+ \frac{s}{K_M \left(1 + \frac{x^h}{K_I^h}\right) + s} \quad (12)$$

In this case, we can define an *effective* inhibition constants K_{IC} , that will allow us to rewrite this rate law in a form identical to non-competitive

inhibition:

$$\begin{aligned}
K_{IC} &\equiv K_I \sqrt[h]{\frac{K_M + s}{K_M}} \\
v &= V^+ \frac{s}{K_M \left(1 + \frac{x^h}{K_I^h}\right) + s} = V^+ \frac{s}{K_M \left(1 + \frac{x^h (K_M + s)}{K_{IC}^h K_M}\right) + s} = \\
&= V^+ \frac{s}{K_M + s + \frac{x^h (K_M + s)}{K_{IC}^h}} = V^+ \frac{s}{K_M + s} \cdot \frac{1}{1 + \frac{x^h}{K_{IC}^h}}
\end{aligned} \tag{13}$$

so we can see here that in this case $\theta = \frac{K_{IC}^h}{K_{IC}^h + x^h}$, exactly like in the case of non-competitive inhibition. Of course, the difference here is that K_{IC} is not a binding constant but rather a function of s , K_M , and K_I . Nevertheless, we can use the same formula for the elasticity (since K_{IC} is a constant with regards to x):

$$\epsilon_x^v = \frac{\partial v}{\partial x} \frac{x}{v} = -h(1 - \theta) \tag{14}$$

2.3.4 Uncompetitive Inhibition

Uncompetitive inhibition is another case where substrate and inhibitor saturations are entangled:

$$v = V^+ \frac{s}{K_M + s \left(1 + \frac{x^h}{K_I^h}\right)} \tag{15}$$

However, the same procedure as in competitive inhibition can also be used here, by defining an effective constant K_{IU} , that will allow us to rewrite this rate law in a form identical to non-competitive inhibition:

$$\begin{aligned}
K_{IU} &\equiv K_I \sqrt[h]{\frac{K_M + s}{s}} \\
v &= V^+ \frac{s}{K_M + s \left(1 + \frac{x^h}{K_I^h}\right)} = V^+ \frac{s}{K_M + s \left(1 + \frac{x^h (K_M + s)}{K_{IU}^h s}\right)} = \\
&= V^+ \frac{s}{K_M + s + \frac{x^h (K_M + s)}{K_{IU}^h}} = V^+ \frac{s}{K_M + s} \cdot \frac{1}{1 + \frac{x^h}{K_{IU}^h}}
\end{aligned} \tag{16}$$

which shows that just like in the case of competitive inhibition, $\theta = \frac{K_{IU}^h}{K_{IU}^h + x^h}$, and the formula for elasticity remains

$$\epsilon_x^v = -h(1 - \theta). \tag{17}$$

2.3.5 Generalized Inhibition Model

A general formula for reversible reactions with non-cooperative competitive, uncompetitive, mixed and pure non-competitive inhibition is given in Chapter 5 of Sauro [7]:

$$v = \frac{\frac{V_m}{K_S} \left(s - \frac{p}{K_{eq}} \right) \left(1 + b \frac{x}{aK_I} \right)}{\left(1 + \frac{s}{K_S} + \frac{p}{K_P} \right) + \frac{x}{K_I} \left(1 + \frac{s}{aK_S} + \frac{p}{aK_P} \right)}. \quad (18)$$

The parameter b represents the amount of inhibition, i.e. the activity of the enzyme when the inhibitor is bound to it. If we assume there is no activity in the bound state, also known as **complete inhibition**, then we can set $b = 0$. Then, depending on the value of a , the above formula converges to the different cases discussed earlier. For example, if $a \rightarrow \infty$, then this rate law is equivalent to competitive inhibition. When $a = 1$ it is equivalent to pure non-competitive inhibition. Finally, if $a \rightarrow 0$ (more precisely, when $K_I \gg aK_I$), it becomes uncompetitive inhibition.

The elasticity we get for this general form is:

$$\epsilon_x^v = \frac{b \frac{x}{aK_I}}{1 + b \frac{x}{aK_I}} - \frac{\frac{x}{K_I} \left(1 + \frac{s}{aK_S} + \frac{p}{aK_P} \right)}{\left(1 + \frac{s}{K_S} + \frac{p}{K_P} \right) + \frac{x}{K_I} \left(1 + \frac{s}{aK_S} + \frac{p}{aK_P} \right)} \quad (19)$$

as with previous cases, we can define an effective K_E :

$$K_{E1} \equiv K_I \frac{a}{b} \quad (20)$$

$$K_{E2} \equiv K_I \frac{1 + \frac{s}{K_S} + \frac{p}{K_P}}{1 + \frac{s}{aK_S} + \frac{p}{aK_P}} \quad (21)$$

and therefore that we get:

$$\begin{aligned} v &= \frac{\frac{V_m}{K_S} \left(s - \frac{p}{K_{eq}} \right)}{\left(1 + \frac{s}{K_S} + \frac{p}{K_P} \right)} \cdot \frac{1 + \frac{x}{K_{E1}}}{1 + \frac{x}{K_{E2}}} \\ \epsilon_x^v &= \frac{x}{K_{E1} + x} - \frac{x}{K_{E2} + x} \end{aligned} \quad (22)$$

As expected, in all cases of complete inhibition ($b \rightarrow 0$ which means that $K_{E1} \rightarrow \infty$), we reach exactly the same conclusion as in the extreme cases, namely:

$$\begin{aligned} \theta &= \frac{1 + \frac{x}{K_{E1}}}{1 + \frac{x}{K_{E2}}} \rightarrow \frac{K_{E2}}{K_{E2} + x} \\ \epsilon_x^v &\rightarrow -(1 - \theta) \end{aligned} \quad (23)$$

For incomplete inhibition, however, there is a more complex relationship between relative activity and inhibition.

2.3.6 The Monod-Changeux-Jacob model

Probably the most widespread model for cooperativity of substrate and effector binding was proposed by Monod et al. [6]. In this model the rate law is given by the following formula:

$$v = V^+ \cdot \frac{s}{K_M + s} \cdot \frac{\left(1 + \frac{s}{K_M}\right)^h \left(1 + \frac{a}{K_A}\right)^h \left(1 + \frac{x}{K_I}\right)^{-h}}{L + \left(1 + \frac{s}{K_M}\right)^h \left(1 + \frac{a}{K_A}\right)^h \left(1 + \frac{x}{K_I}\right)^{-h}} \quad (24)$$

where a is the concentration of an inhibitor, and K_M , K_A and K_I are the dissociation constants for the active form of the enzyme (for s , a and x respectively), n is the number of subunits, and L is the allosteric constant. As before, we can define the saturation term κ and the relative activity term θ (i.e. the decrease from maximal activity):

$$\begin{aligned} \kappa &\equiv \frac{s}{K_M + s} \\ \theta &\equiv \frac{\left(1 + \frac{s}{K_M}\right)^h \left(1 + \frac{a}{K_A}\right)^h \left(1 + \frac{x}{K_I}\right)^{-h}}{L + \left(1 + \frac{s}{K_M}\right)^h \left(1 + \frac{a}{K_A}\right)^h \left(1 + \frac{x}{K_I}\right)^{-h}} \\ v &= V^+ \cdot \kappa \cdot \theta \end{aligned} \quad (25)$$

To simplify the form of the equations, let

$$g(x) \equiv \left(1 + \frac{s}{K_M}\right)^h \left(1 + \frac{a}{K_A}\right)^h \left(1 + \frac{x}{K_I}\right)^{-h} \quad (26)$$

so that $\theta = \frac{g(x)}{L+g(x)}$, which we write in shorthand as $\theta = \frac{g}{L+g}$. Then,

$$\epsilon_x^v = V^+ \cdot \kappa \cdot \frac{\partial \theta}{\partial x} \cdot \frac{x}{v} = V^+ \cdot \kappa \cdot \frac{L}{(L+g)^2} \cdot \frac{\partial g}{\partial x} \cdot \frac{x}{v} \quad (27)$$

Now, we can calculate that

$$\frac{\partial g}{\partial x} = \frac{-h}{K_I + x} \cdot g \quad (28)$$

Substituting this form of $\frac{\partial g}{\partial x}$ into the prior equation, we arrive at

$$\epsilon_x^v = V^+ \cdot \kappa \cdot \frac{g}{L+g} \cdot \frac{-hx}{K_I + x} \cdot \frac{L}{L+g} \cdot \frac{1}{v} \quad (29)$$

Note that the first three terms are precisely equal to v , so they cancel with the last term. Also note that $\frac{L}{L+g} = 1 - \frac{g}{L+g} = 1 - \theta$. Thus, we arrive at

$$\epsilon_x^v = \frac{\partial \ln v}{\partial \ln x} = \frac{\partial \ln \theta}{\partial \ln x} = -h \frac{x}{K_I + x} (1 - \theta) . \quad (30)$$


In [3], the elasticity of the inhibitor was calculated for this rate law (denoted as *effector strength* or X_I in the original text) and matches the form above. Although the final formulae for the elasticities in the MWC case are not identical to the other cases dealt with before, we can still reach a similar conclusion, where the absolute value of the elasticity is larger when the relative activity is smaller.

2.4 A Trade-off between Enzyme Cost and Flux Control

By taking the absolute value of the elasticity, we can compress the results for activators (Equation 9) and inhibitors (Equations 11, 13, 17, and 23, excluding the MWC model) into one formula:

$$|\epsilon_x^v| = h(1 - \theta) . \quad (31)$$

This relationship can be visualized in the following plot (for different values of h):



Since θ represents the fraction of active enzyme, we see here that there is direct trade-off between the activity of the enzyme and the elasticity. It should be noted, that evolution can easily adjust θ for an individual enzyme by changing the K_A or K_I values, even without changing the concentration of the small-molecule effector (assuming it has other crucial functions in

the cell). Therefore, evolution needs to weigh between how much of the enzyme is "wasted" by inhibition (or by inactivation), versus how much control it has on the flux. This can be viewed as a trade-off between the short-term goal of being able to adjust things quickly and the long-term goal of allocating resources efficiently in order to grow as fast as possible.

Another corollary of Equation 31 is that the control can be increased by changing the Hill coefficient (h). This could be a reason why mechanisms with very high cooperativity evolve for allosteric regulation [6], as it increases elasticity without the added cost of losing enzyme activity.

3 Supplementary Figures

References

- [1] D Fell. *Understanding the Control of Metabolism*. Portland Press, 1 November 1996.
- [2] E Noor, A Flamholz, W Liebermeister, A Bar-Even, and R Milo. A note on the kinetics of enzyme action: a decomposition that highlights thermodynamic effects. *FEBS Lett.*, 587:2772–2777, 2 September 2013.
- [3] R Heinrich and A T Rapoport. A linear Steady-State treatment of enzymatic chains. general properties, control and effector strength. *Eur. J. Biochem.*, 42:89–95, 1 February 1974.
- [4] Wolfram Liebermeister, Jannis Uhlendorf, and Edda Klipp. Modular rate laws for enzymatic reactions: thermodynamics, elasticities and implementation. *Bioinformatics*, 26(12):1528–1534, 15 June 2010.
- [5] J Barcroft and A V Hill. The nature of oxyhaemoglobin, with a note on its molecular weight. *J. Physiol.*, 39(6):411–428, 8 March 1910.
- [6] J Monod, J Wyman, and J P Changeux. On the nature of allosteric transitions: A plausible model. *J. Mol. Biol.*, 12:88–118, May 1965.
- [7] H M Sauro. *Enzyme Kinetics for Systems Biology*. Future Skill Software, 2011.




Figure 1: Distance between metabolites and enzymes in the (a) genome-scale metabolite network iJO1366 and (b)-(d) the SMRN of *E. coli*. In general, interactions in the SMRN traverse a shorter distance than the typical distance between a randomly chosen metabolite and enzyme in metabolism. Several highly-connected metabolites (i.e. co-factors) that were removed from the network before preparing the bipartite graph: h, h2o, co2, o2, pi, atp, adp, amp, nad, nadh, nadp, nadph, coa, thf, 5mthf, 5fthf, methf, mlthf, nh4, cmp, q8, q8h2, udp, updg, fad, fadh2, ade, ctp, gtp, h2o2, mql8, mqn8, na1, ppi, acp




Figure 2: Histogram of the number of interactions for reactions and metabolite in the (a)-(b) entire E. coli SMRN, and (c)-(d) in the region of the SMRN restricted to the central carbon metabolism (CCM) of E. coli.




Figure 3: Number of interactions in the SMRN targeting each pathway in *E. coli* metabolism.




Figure 4: Schematic of the relationship between kinetic rate laws and elasticity for (a)-(b) substrates obeying Michaelis-Menten kinetics and (c)-(d) inhibitors obeying non-competitive inhibition. Importantly, substrate elasticity is maximized at low substrate concentrations, and inhibitor elasticity is maximized at high inhibitor concentration.




Figure 5: A heatmap of the median values of each metabolite's elasticities across a variety of nutrient conditions. Left-hand-side corresponds to substrate interactions for that metabolite, and right-hand-side corresponds to inhibitory interactions.




Figure 6: For 45 metabolites, we were able to obtain 2 or more estimates each of K_M and K_I values. This enabled us to directly compare, on a single-metabolite basis, whether the distributions of its K_M and K_I values differed substantially. Doing so, we found that 5 metabolites (PEP, ATP, methionine, succinate, and dGMP) each showed significantly higher values of K_I values compared to K_M values (FDR-adjusted Mann-Whitney p-value < 0.1) and only one metabolite (phosphate) had significantly lower K_I values.




Figure 7: A map of the reactions branching from central carbon metabolism for amino acid biosynthesis. Amino acids in red color feedback inhibit the first enzyme of their biosynthesis pathway, based on the SMRN. For the remaining four amino acids (Glycine, Alanine, Aspartate and Glutamate) there is no evidence of feedback inhibition.




Figure 8: A map of the CCM reactions that are regulated by small molecule(s) in the SMRN. Orange reactions only show evidence of inhibitory interactions, blue reactions only show evidence of activating interactions, and purple reactions show evidence of both activating and inhibiting interactions.