


Statistical Methods for Data Mining

Kuangnan Fang

Xiamen University *Email:* xmufkn@xmu.edu.cn



Linear Model Selection and Regularization

- Recall the linear model

$$Y = \beta_0 + \beta_1 X_1 + \cdots + \beta_p X_p + \epsilon.$$

- In the lectures that follow, we consider some approaches for extending the linear model framework. In the lectures covering Chapter 7 of the text, we generalize the linear model in order to accommodate *non-linear*, but still *additive*, relationships.
- In the lectures covering Chapter 8 we consider even more general *non-linear* models.

In praise of linear models!

- Despite its simplicity, the linear model has distinct advantages in terms of its *interpretability* and often shows good *predictive performance*.
- Hence we discuss in this lecture some ways in which the simple linear model can be improved, by replacing ordinary least squares fitting with some alternative fitting procedures.

线性模型的优势·可解释性与
预测能力的平衡

Why consider alternatives to least squares?

引进原因：①预测精度， p 很大时，预测方差大
与预测 $p > n$ 时，OLS 无法估计
②模型可解形性：(易得零点、
变量选择)

- *Prediction Accuracy*: especially when $p > n$, to control the variance.
- *Model Interpretability*: By removing irrelevant features — that is, by setting the corresponding coefficient estimates to zero — we can obtain a model that is more easily interpreted. We will present some approaches for automatically performing *feature selection*.

目的：降方差，提解释力
子集选择 \rightarrow OLS
Three classes of methods 收缩估计 \rightarrow 正则化。
降维 \rightarrow 投影

- *Subset Selection*. We identify a subset of the p predictors that we believe to be related to the response. We then fit a model using least squares on the reduced set of variables.
- *Shrinkage*. We fit a model involving all p predictors, but the estimated coefficients are shrunk towards zero relative to the least squares estimates. This shrinkage (also known as *regularization*) has the effect of reducing variance and can also perform variable selection.
- *Dimension Reduction*. We project the p predictors into a M -dimensional subspace, where $M < p$. This is achieved by computing M different *linear combinations*, or *projections*, of the variables. Then these M projections are used as predictors to fit a linear regression model by least squares.

Subset Selection


最优子集选择 (穷举)
逐步回归

Best subset and stepwise model selection procedures

Best Subset Selection

1. Let \mathcal{M}_0 denote the *null model*, which contains no predictors. This model simply predicts the sample mean for each observation.
2. For $k = 1, 2, \dots, p$:
 - (a) Fit all $\binom{p}{k}$ models that contain exactly k predictors.
 - (b) Pick the best among these $\binom{p}{k}$ models, and call it \mathcal{M}_k . Here best is defined as having the smallest RSS, or equivalently
largest R^2 .
k个变量的模型中, Best 由 RSS_{min} 或 $R^2_{largest}$.
3. Select a single best model from among $\mathcal{M}_0, \dots, \mathcal{M}_p$ using cross-validated prediction error, C_p (AIC), BIC, or adjusted R^2 .
各 k 之间用 C_p (AIC), BIC, Adj R^2 选 Best

Example- Credit data set



For each possible model containing a subset of the ten predictors in the **Credit** data set, the RSS and R^2 are displayed. The red frontier tracks the **best** model for a given number of predictors, according to RSS and R^2 . Though the data set contains only ten predictors, the x -axis ranges from 1 to 11, since one of the variables is categorical and takes on three values, leading to the creation of two dummy variables

Extensions to other models

Best subset 也可以用于 logistic reg

近似元 RSS. 用 $-2 \log(L(x)) = \text{deviance}$

似然模型(同 k 中)

- Although we have presented best subset selection here for least squares regression, the same ideas apply to other types of models, such as logistic regression.
- The *deviance*— negative two times the maximized log-likelihood— plays the role of RSS for a broader class of models.

Stepwise Selection

- For computational reasons, best subset selection cannot be applied with very large p . *Why not?* ① Best subset 计算量随 p 增大而增大
- Best subset selection may also suffer from statistical problems when p is large: larger the search space, the higher the chance of finding models that look good on the training data, even though they might not have any predictive power on future data. ② 可能选出样本最优 → overfitting & 高方差
- Thus an enormous search space can lead to *overfitting* and high variance of the coefficient estimates.
- For both of these reasons, *stepwise* methods, which explore a far more restricted set of models, are attractive alternatives to best subset selection.

Forward Stepwise Selection

- Forward stepwise selection begins with a model containing no predictors, and then adds predictors to the model, one-at-a-time, until all of the predictors are in the model.
- In particular, at each step the variable that gives the greatest *additional* improvement to the fit is added to the model.

In Detail

Forward Stepwise Selection

每一步都用
 $R^2 \downarrow$ 表示
每轮保存一个
最好的 M_k 集合
 \downarrow

1. Let \mathcal{M}_0 denote the *null* model, which contains no predictors.
2. For $k = 0, \dots, p - 1$:
 - 2.1 Consider all $p - k$ models that augment the predictors in \mathcal{M}_k with one additional predictor. *添加一个*
 - 2.2 Choose the *best* among these $p - k$ models, and call it \mathcal{M}_{k+1} . Here *best* is defined as having smallest RSS or highest R^2 .
3. Select a single best model from among $\mathcal{M}_0, \dots, \mathcal{M}_p$ using *CV*, *BIC*, *Adj R²*, *CV 等价*, *(by) MSE*, *平均数*.

$$\text{7. } \frac{1}{n} [p + (p+1) + (p+2) + \dots + 1] = \frac{p(p+1)}{2} + 1 \rightarrow p^2 \text{ 次 (仍为平方项)}$$

More on Forward Stepwise Selection

不保证选出样本内最优

- Computational advantage over best subset selection is clear.
- It is not guaranteed to find the best possible model out of all 2^p models containing subsets of the p predictors. *Why not? Give an example.*

Credit data example

# Variables	Best subset	Forward stepwise
One	rating	rating
Two	rating, income	rating, income
Three	rating, income, student	rating, income, student
Four	cards, income student, limit	rating, income, student, limit

The first four selected models for best subset selection and forward stepwise selection on the Credit data set. The first three models are identical but the fourth models differ.

Backward Stepwise Selection

- Like forward stepwise selection, *backward stepwise selection* provides an efficient alternative to best subset selection.
- However, unlike forward stepwise selection, it begins with the full least squares model containing all p predictors, and then iteratively removes the least useful predictor, one-at-a-time.

Backward Stepwise Selection: details

Backward Stepwise Selection

1. Let \mathcal{M}_p denote the *full* model, which contains all p predictors.
2. For $k = p, p-1, \dots, 1$:
 - 2.1 Consider all k models that contain all but one of the predictors in \mathcal{M}_k , for a total of $k-1$ predictors.
 - 2.2 Choose the *best* among these k models, and call it \mathcal{M}_{k-1} . Here *best* is defined as having smallest RSS or highest R^2 .
3. Select a single best model from among $\mathcal{M}_0, \dots, \mathcal{M}_p$ using cross-validated prediction error, C_p (AIC), BIC, or adjusted R^2 .

More on Backward Stepwise Selection

- Like forward stepwise selection, the backward selection approach searches through only $1 + p(p + 1)/2$ models, and so can be applied in settings where p is too large to apply best subset selection
缺点① 不保证选出样本内最优
- Like forward stepwise selection, backward stepwise selection is not guaranteed to yield the *best* model containing a subset of the p predictors.
- Backward selection requires that the *number of samples n is larger than the number of variables p* (so that the full model can be fit). In contrast, forward stepwise can be used even when $n < p$, and so is the only viable subset method when p is very large.
② p>n时无法执行

Choosing the Optimal Model

- The model containing all of the predictors will always have the smallest RSS and the largest R^2 , since these quantities are related to the training error.
- We wish to choose a model with low test error, not a model with low training error. Recall that training error is usually a poor estimate of test error.
- Therefore, RSS and R^2 are not suitable for selecting the best model among a collection of models with different numbers of predictors.

rss, R^2 险度量数↑，越优
(模型复杂度↑) (过拟合)

Estimating test error: two approaches

不同參量數量下，估計預測误差

(R^2 , less R^2 , etc.)




{ ① 修正 (BIC , AIC , C_p)
② CV.

- We can indirectly estimate test error by making an *adjustment* to the training error to account for the bias due to overfitting.
- We can *directly* estimate the test error, using either a validation set approach or a cross-validation approach, as discussed in previous lectures.
- We illustrate both approaches next.

C_p , AIC, BIC, and Adjusted R^2

- These techniques adjust the training error for the model size, and can be used to select among a set of models with different numbers of variables.
- The next figure displays C_p , BIC, and adjusted R^2 for the best model of each size produced by best subset selection on the **Credit** data set.

Credit data example



min

- Mallow's C_p :

Now for some details

$$C_p^* = \frac{\text{RSS}}{\hat{\sigma}^2} + 2d - n$$

$$C_p = \frac{1}{n} \hat{\sigma}^2 (C_p^* + n)$$

$$C_p = \frac{1}{n} (\text{RSS} + 2d\hat{\sigma}^2),$$

$\left\{ \begin{array}{l} \text{RSS} \downarrow \\ d \uparrow \end{array} \right.$

where d is the total # of parameters used and $\hat{\sigma}^2$ is an estimate of the variance of the error ϵ associated with each response measurement.

min

- The **AIC** criterion is defined for a large class of models fit by maximum likelihood:

$$\text{AIC} = -2 \log L + 2 \cdot d$$

where L is the maximized value of the likelihood function for the estimated model. 当线性模型误差项为高斯误差时, $ML = LS$

- In the case of the linear model with Gaussian errors, C_p & AIC maximum likelihood and least squares are the same thing, and C_p and AIC are equivalent. Prove this. $AIC = \frac{1}{n\hat{\sigma}^2} (\text{RSS} + 2d\hat{\sigma}^2)$

$$\text{AIC} = 1/(n\hat{\sigma}^2)(\text{RSS} + 2d\hat{\sigma}^2)$$

Details on BIC

min

$$\text{BIC} = \frac{1}{n} (\text{RSS} + \log(n)d\hat{\sigma}^2).$$

training error ↑ 練習误差↑
penalty ↑ 2(C_p)

- Like C_p , the BIC will tend to take on a small value for a model with a low test error, and so generally we select the model that has the lowest BIC value.
- Notice that BIC replaces the $2d\hat{\sigma}^2$ used by C_p with a $\log(n)d\hat{\sigma}^2$ term, where n is the number of observations.
- Since $\log n > 2$ for any $n > 7$, the BIC statistic generally places a heavier penalty on models with many variables, and hence results in the selection of smaller models than C_p . See Figure on slide 19.

$n > 7, \log n > 2 \Rightarrow \text{BIC has penalty更重}$
 倾向于更少的变量

$$\begin{aligned} p \uparrow \Rightarrow & TSS = \sum (y_i - \bar{y})^2 \text{ 不变} \\ \text{Adjusted } R^2 & \downarrow \quad \text{RSS} = \sum (f_i - y_i)^2 \end{aligned} \quad \Rightarrow R^2 \uparrow$$

- For a least squares model with d variables, the adjusted R^2 statistic is calculated as

$$\text{Adjusted } R^2 = 1 - \frac{\text{RSS}/(n-d-1)}{\text{TSS}/(n-1)}.$$

↗ $\text{RSS} \downarrow$
 ↗ $n-d-1 \downarrow$
 ↗ 调整常数
 ↗ 看引入变量对 RSS 改善效果

where TSS is the total sum of squares.

- Unlike C_p , AIC, and BIC, for which a *small* value indicates a model with a low test error, a *large* value of adjusted R^2 indicates a model with a small test error.
- Maximizing the adjusted R^2 is equivalent to minimizing $\frac{\text{RSS}}{n-d-1}$. While RSS always decreases as the number of variables in the model increases, $\frac{\text{RSS}}{n-d-1}$ may increase or decrease, due to the presence of d in the denominator.
- Unlike the R^2 statistic, the adjusted R^2 statistic pays a price for the inclusion of unnecessary variables in the model. See Figure on slide 19.

Validation and Cross-Validation

- Each of the procedures returns a sequence of models \mathcal{M}_k indexed by model size $k = 0, 1, 2, \dots$. Our job here is to select \hat{k} . Once selected, we will return model $\mathcal{M}_{\hat{k}}$

调整法的缺陷 · 合理性不足
↓
自适应不好 . → CV更实用

用途:
从 M_0, M_1, \dots, M_K 中选出 M_k






Validation and Cross-Validation

- Each of the procedures returns a sequence of models \mathcal{M}_k indexed by model size $k = 0, 1, 2, \dots$. Our job here is to select \hat{k} . Once selected, we will return model $\mathcal{M}_{\hat{k}}$
- We compute the validation set error or the cross-validation error for each model \mathcal{M}_k under consideration, and then select the k for which the resulting estimated test error is smallest.
- This procedure has an advantage relative to AIC, BIC, C_p , and adjusted R^2 , in that it provides a direct estimate of the test error, and doesn't require an estimate of the error variance σ^2 .
- It can also be used in a wider range of model selection tasks, even in cases where it is hard to pinpoint the model degrees of freedom (e.g. the number of predictors in the model) or hard to estimate the error variance σ^2 .

优点

Credit data example



one standard error rule
往左取一个标准差，更向左

Details of Previous Figure

$k=4$

- The validation errors were calculated by randomly selecting three-quarters of the observations as the training set, and the remainder as the validation set.
- The cross-validation errors were computed using $k = 10$ folds. In this case, the validation and cross-validation methods both result in a six-variable model.
- However, all three approaches suggest that the four-, five-, and six-variable models are roughly equivalent in terms of their test errors.
- In this setting, we can select a model using the one-standard-error rule. We first calculate the standard error of the estimated test MSE for each model size, and then select the smallest model for which the estimated test error is within one standard error of the lowest point on the curve. *What is the rationale for this?* 一个标准误差范围内

bootstrap

Shrinkage Methods

Ridge regression and *Lasso* 亂世幫·皇者降臨之言。

- The subset selection methods use least squares to fit a linear model that contains a subset of the predictors.
- As an alternative, we can fit a model containing all p predictors using a technique that *constrains* or *regularizes* the coefficient estimates, or equivalently, that *shrinks* the coefficient estimates towards zero.
- It may not be immediately obvious why such a constraint should improve the fit, but it turns out that shrinking the coefficient estimates can significantly reduce their variance.

Ridge regression

- Recall that the least squares fitting procedure estimates $\beta_0, \beta_1, \dots, \beta_p$ using the values that minimize

$$\text{RSS} = \sum_{i=1}^n \left(y_i - \beta_0 - \sum_{j=1}^p \beta_j x_{ij} \right)^2.$$

- In contrast, the ridge regression coefficient estimates $\hat{\beta}^R$ are the values that minimize

$$\sum_{i=1}^n \left(y_i - \beta_0 - \sum_{j=1}^p \beta_j x_{ij} \right)^2 + \lambda \underbrace{\sum_{j=1}^p \beta_j^2}_{\text{penalty}} = \text{RSS} + \lambda \sum_{j=1}^p \beta_j^2,$$

where $\lambda \geq 0$ is a *tuning parameter*, to be determined separately.

Ridge regression (主要用于解决多重共线性)

- Writing the criterion in matrix form

$$RSS(\lambda) = (y - X\beta)^T(y - X\beta) + \lambda\beta^T\beta$$

- the ridge regression solutions are easily seen to be

$$\beta^{ridge} = (X^T X + \lambda I)^{-1} X^T y$$

where I is the $p \times p$ identity matrix.


仍存解折解 $(\underbrace{X'X + \lambda I}_{\Downarrow})^{-1} X' Y$.

使后 $X'X$ 在其设计下不至为奇异
(non singular)


Ridge regression: continued (CV & λ)

- As with least squares, ridge regression seeks coefficient estimates that fit the data well, by making the RSS small.
- However, the second term, $\lambda \sum_j \beta_j^2$, called a *shrinkage penalty*, is small when β_1, \dots, β_p are close to zero, and so it has the effect of *shrinking* the estimates of β_j towards zero.
- The tuning parameter λ serves to control the relative impact of these two terms on the regression coefficient estimates.
- Selecting a good value for λ is critical; cross-validation is used for this.

Credit data example



$$\begin{aligned} L_1 &:= \sum |\beta_j| \\ L_2 &:= \sqrt{\sum \beta_j^2} \\ L_\infty &:= \sum |\beta_j| \quad (\beta_j \neq 0) \end{aligned}$$



$$\begin{aligned} \|\cdot\|_2 &: L_2\text{-norm} = \text{范数} \\ \|\hat{\beta}_\lambda^R\|_2 / \|\hat{\beta}\|_2 & \downarrow \quad \downarrow \\ \text{ridge} & \quad \text{OLS} \end{aligned}$$

Details of Previous Figure

- In the left-hand panel, each curve corresponds to the ridge regression coefficient estimate for one of the ten variables, plotted as a function of λ .
- The right-hand panel displays the same ridge coefficient estimates as the left-hand panel, but instead of displaying λ on the x -axis, we now display $\|\hat{\beta}_\lambda^R\|_2/\|\hat{\beta}\|_2$, where $\hat{\beta}$ denotes the vector of least squares coefficient estimates.
- The notation $\|\beta\|_2$ denotes the ℓ_2 norm (pronounced “ell 2”) of a vector, and is defined as $\|\beta\|_2 = \sqrt{\sum_{j=1}^p \beta_j^2}$.

Ridge regression: scaling of predictors

- The standard least squares coefficient estimates are *scale equivariant*: multiplying X_j by a constant c simply leads to a scaling of the least squares coefficient estimates by a factor of $1/c$. In other words, regardless of how the j th predictor is scaled, $X_j \hat{\beta}_j$ will remain the same.
- In contrast, the ridge regression coefficient estimates can change *substantially* when multiplying a given predictor by a constant, due to the sum of squared coefficients term in the penalty part of the ridge regression objective function.
- Therefore, it is best to apply ridge regression after *standardizing the predictors*, using the formula


标准化
公式
除以标准差
即可

$$\tilde{x}_{ij} = \frac{x_{ij}}{\sqrt{\frac{1}{n} \sum_{i=1}^n (x_{ij} - \bar{x}_j)^2}}$$

在 LS 中，变量成倍
增加时， β 也会成倍
增加
在 Ridge 中则不会
因为 Ridge 前面有变量
标准化

Why Does Ridge Regression Improve Over Least Squares?

The Bias-Variance tradeoff



Simulated data with $n = 50$ observations, $p = 45$ predictors, all having nonzero coefficients. Squared bias (black), variance (green), and test mean squared error (purple) for the ridge regression predictions on a simulated data set, as a function of λ and $\|\hat{\beta}_\lambda^R\|_2/\|\hat{\beta}\|_2$. The horizontal dashed lines indicate the minimum possible MSE. The purple crosses indicate the ridge regression models for which the MSE is smallest.

The Lasso

Ridge 缺点：无法筛选变量

- Ridge regression does have one obvious disadvantage: unlike subset selection, which will generally select models that involve just a subset of the variables, ridge regression will include all p predictors in the final model
- The *Lasso* is a relatively recent alternative to ridge regression that overcomes this disadvantage. The lasso coefficients, $\hat{\beta}_\lambda^L$, minimize the quantity

$$\sum_{i=1}^n \left(y_i - \beta_0 - \sum_{j=1}^p \beta_j x_{ij} \right)^2 + \boxed{\lambda \sum_{j=1}^p |\beta_j|} = \text{RSS} + \lambda \sum_{j=1}^p |\beta_j|.$$

- In statistical parlance, the lasso uses an ℓ_1 (pronounced “ell 1”) penalty instead of an ℓ_2 penalty. The ℓ_1 norm of a coefficient vector β is given by $\|\beta\|_1 = \sum |\beta_j|$.

$$\|\beta\|_2 = \sqrt{\sum \beta_j^2} \quad \|\beta\|_0 = \sum |\beta_j|$$



$$\|\beta\|_2 = \sqrt{\sum \beta_j^2} \quad \|\beta\|_0 = \sum |\beta_j|$$

The Lasso: continued

- As with ridge regression, the lasso shrinks the coefficient estimates towards zero.
- However, in the case of the lasso, the ℓ_1 penalty has the effect of forcing some of the coefficient estimates to be exactly equal to zero when the tuning parameter λ is sufficiently large.
- Hence, much like best subset selection, the lasso performs *variable selection*.
- We say that the lasso yields *sparse models* — that is, models that involve only a subset of the variables.
- As in ridge regression, selecting a good value of λ for the lasso is critical; cross-validation is again the method of choice.

好
處
縮減
係
數
變
量
選
擇

Example: Credit dataset



The Variable Selection Property of the Lasso

Why is it that the lasso, unlike ridge regression, results in coefficient estimates that are exactly equal to zero?

The Variable Selection Property of the Lasso

Why is it that the lasso, unlike ridge regression, results in coefficient estimates that are exactly equal to zero?

One can show that the lasso and ridge regression coefficient estimates solve the problems

$$\underset{\beta}{\text{minimize}} \sum_{i=1}^n \left(y_i - \beta_0 - \sum_{j=1}^p \beta_j x_{ij} \right)^2$$

推导出， β 的最小化问题

$$\text{subject to } \sum_{j=1}^p |\beta_j| \leq s$$

gg penalty
- 一范数惩罚
趋向归零

Lasso.


$$\underset{\beta}{\text{minimize}} \sum_{i=1}^n \left(y_i - \beta_0 - \sum_{j=1}^p \beta_j x_{ij} \right)^2$$

$$\text{subject to } \sum_{j=1}^p \beta_j^2 \leq s,$$

ridge

and respectively.

The Lasso Picture



soft threshold.




shrinkage methods

TABLE 3.4. Estimators of β_j in the case of orthonormal columns of \mathbf{X} . M and λ are constants chosen by the corresponding techniques; sign denotes the sign of its argument (± 1), and x_+ denotes “positive part” of x . Below the table, estimators are shown by broken red lines. The 45° line in gray shows the unrestricted estimate for reference.

Estimator	Formula
Best subset (size M)	$\hat{\beta}_j \cdot I(\hat{\beta}_j \geq \hat{\beta}_{(M)})$
Ridge	$\hat{\beta}_j / (1 + \lambda)$
Lasso	$\text{sign}(\hat{\beta}_j)(\hat{\beta}_j - \lambda)_+$

$$\hat{\beta} = \begin{cases} 0 & \\ \hat{\beta}_1 > 0 & \\ \hat{\beta}_2 > 0 & \\ \vdots & \\ \hat{\beta}_M > 0 & \end{cases}$$

Best Subset



Coordinate descent algorithm for Lasso

Algorithm 1 Coordinate descent minimization

1: Let $\beta^{[0]} \in \mathbb{R}^p$ be an initial parameter vector. Set $m = 0$.

2: **repeat**

3: Increase m by one: $m \leftarrow m + 1$.

Denote by $\mathcal{S}^{[m]}$ the index cycling through the coordinates $\{1, \dots, p\}$:

$\mathcal{S}^{[m]} = \mathcal{S}^{[m-1]} + 1 \bmod p$. Abbreviate by $j = \mathcal{S}^{[m]}$ the value of $\mathcal{S}^{[m]}$.

4: if $|G_j(\beta_{-j}^{[m-1]})| \leq \lambda$: set $\beta_j^{[m]} = 0$,

otherwise: $\beta_j^{[m]} = \arg \min_{\beta_j} Q_\lambda(\beta_{+j}^{[m-1]})$,

where $\beta_{-j}^{[m-1]}$ is the parameter vector where the j th component is set to zero and $\beta_{+j}^{[m-1]}$ is the parameter vector which equals $\beta^{[m-1]}$ except for the j th component where it is equal to β_j (i.e. the argument we minimize over).

5: **until** numerical convergence

$$\begin{array}{l} \beta^{[0]} = (\dots) \\ m=0 \end{array} \Rightarrow \begin{array}{l} m=1 \text{ (迭代次数)} \\ \text{一次只变一个系数,用上一步的值} \end{array}$$

Coordinate descent algorithm for Lasso

The coordinatewise optimization above can easily incorporate the more general case where some parameters are unpenalized, i.e.,

$$\hat{\beta} = \arg \min_{\beta} Q_{\lambda}(\beta),$$

$$Q_{\lambda}(\beta) = \|\mathbf{Y} - \mathbf{X}\beta\|_2^2/n + \lambda \sum_{j=r+1}^p |\beta_j|,$$

选择部分变量进行惩罚.

and thus, β_1, \dots, β_r are unpenalized. The up-dating step in the optimization algorithm then looks as follows:


$$\text{if } j \in \{1, \dots, r\} : \beta_j^{[m]} = \arg \min_{\beta_j} Q_{\lambda}(\beta_{-j}^{[m-1]}),$$

$$\text{if } j \in \{r+1, \dots, p\} :$$

$$\text{if } |G_j(\beta_{-j}^{[m-1]})| \leq \lambda : \text{set } \beta_j^{[m]} = 0,$$

$$\text{otherwise: } \beta_j^{[m]} = \arg \min_{\beta_j} Q_{\lambda}(\beta_{-j}^{[m-1]}).$$


Comparing the Lasso and Ridge Regression



Left: Plots of squared bias (black), variance (green), and test MSE (purple) for the lasso on simulated data set of Slide 32.

Right: Comparison of squared bias, variance and test MSE between lasso (solid) and ridge (dashed). Both are plotted against their R^2 on the training data, as a common form of indexing. The crosses in both plots indicate the lasso model for which the MSE is smallest.

Comparing the Lasso and Ridge Regression: continued



Left: Plots of squared bias (black), variance (green), and test MSE (purple) for the lasso. The simulated data is similar to that in Slide 38, except that now only two predictors are related to the response. *Right:* Comparison of squared bias, variance and test MSE between lasso (solid) and ridge (dashed). Both are plotted against their R^2 on the training data, as a common form of indexing. The crosses in both plots indicate the lasso model for which the MSE is smallest.

Lasso vs ridge

Conclusions

① 互有优势

② 当自变量仅有-部分与Y有关，
用Lasso，否则 Ridge更好(预测)
③ 实际数据中不知道到底多少变量有关

用CV
判断.

- These two examples illustrate that neither ridge regression nor the lasso will universally dominate the other.
- In general, one might expect the lasso to perform better when the response is a function of only a relatively small number of predictors.
- However, the number of predictors that is related to the response is never known *a priori* for real data sets.
- A technique such as cross-validation can be used in order to determine which approach is better on a particular data set.

Adaptive Lasso

真实非零集合包含在选出的集合中.

LASSO
估计的
集合
 \Rightarrow
真实 model 中非零集合
 $\mathbf{P}[\hat{S} \supseteq S_0] \rightarrow 1 (p \geq n \rightarrow \infty)$

LASSO 选出变量会完全
 \Downarrow

解决:

Adaptive
LASSO.

$$\hat{\beta}_{\text{adapt}}(\lambda) = \operatorname{argmin}_{\beta} \left(\|\mathbf{Y} - \mathbf{X}\beta\|_2^2 / n + \lambda \sum_{j=1}^p \frac{|\beta_j|}{|\hat{\beta}_{\text{init},j}|} \right),$$

不影响

where $\hat{\beta}_{\text{init}}$ is an initial estimator.

初试估计 (可调 LASSO)

The adaptive Lasso has the following obvious property: 相当于用初
步估计作为权

$$\hat{\beta}_{\text{init},j} = 0 \Rightarrow \hat{\beta}_{\text{adapt},j} = 0.$$

重. 原初值大
的惩罚小. 入

若 $\hat{\beta}_i \approx 0$
直接令 $\beta_i = 0$

原初值小的惩罚
大. 逐步缩小

Adaptive Lasso

Lasso (倾向多选) $\xrightarrow{\text{解决}}$ Adaptive Lasso





Fig. 2.4 Estimated regression coefficients in the linear model with $p = 1000$ and $n = 50$. Left: Lasso. Right: Adaptive Lasso with Lasso as initial estimator. The 3 true regression coefficients are indicated with triangles. Both methods used with tuning parameters selected from 10-fold cross-validation.

Generalization of shrinkage methods

- We can generalize ridge regression and the lasso,
Consider the criterion

$$\tilde{\beta} = \underset{\beta}{\operatorname{argmin}} \left\{ \sum_{i=1}^N (y_i - \beta_0 - \sum_{j=1}^p x_{ij} \beta_j)^2 + \lambda \sum_{j=1}^p |\beta_j|^q \right\}$$




FIGURE 3.12. Contours of constant value of $\sum_j |\beta_j|^q$ for given values of q .

- $q > 1$, $|\beta_j|^q$ is differentiable at 0, and so does not share the ability of lasso.

Generalization of shrinkage methods

- Zou and Hastie (2005) introduced the elastic- net penalty

$$\lambda \sum_{j=1}^p (\alpha \beta_j^2 + (1 - \alpha) |\beta_j|)$$

Between Lasso & Ridge

a different compromise between ridge and lasso.





FIGURE 3.13. Contours of constant value of $\sum_j |\beta_j|^q$ for $q = 1.2$ (left plot), and the elastic-net penalty $\sum_j (\alpha \beta_j^2 + (1 - \alpha) |\beta_j|)$ for $\alpha = 0.2$ (right plot). Although visually very similar, the elastic-net has sharp (non-differentiable) corners, while the $q = 1.2$ penalty does not.

MCP - Group Lasso . SCAD .

Selecting the Tuning Parameter for Ridge Regression and Lasso

- As for subset selection, for ridge regression and lasso we require a method to determine which of the models under consideration is best.
- That is, we require a method selecting a value for the tuning parameter λ or equivalently, the value of the constraint s .
(不需要知道矩阵,但计算慢点)
- *Cross-validation* provides a simple way to tackle this problem. We choose a grid of λ values, and compute the cross-validation error rate for each value of λ .
- We then select the tuning parameter value for which the cross-validation error is smallest.
- Finally, the model is re-fit using all of the available observations and the selected value of the tuning parameter.


Credit data example



Left: Cross-validation errors that result from applying ridge regression to the **Credit** data set with various values of λ .

Right: The coefficient estimates as a function of λ . The vertical dashed lines indicates the value of λ selected by cross-validation.

Simulated data example



Left: Ten-fold cross-validation MSE for the lasso, applied to the sparse simulated data set from Slide 39. *Right:* The corresponding lasso coefficient estimates are displayed. The vertical dashed lines indicate the lasso fit for which the cross-validation error is smallest.

Dimension Reduction Methods

- The methods that we have discussed so far in this chapter have involved fitting linear regression models, via least squares or a shrunken approach, using the original predictors, X_1, X_2, \dots, X_p .
- We now explore a class of approaches that *transform* the predictors and then fit a least squares model using the transformed variables. We will refer to these techniques as *dimension reduction* methods.

Dimension Reduction Methods: details

- Let Z_1, Z_2, \dots, Z_M represent $M < p$ linear combinations of our original p predictors. That is,

$$Z_m = \sum_{j=1}^p \phi_{mj} X_j \quad (1)$$

loadings.

$Z_{im} = \sum_{j=1}^p (\phi_{mj} X_{ij})$

score (第 i 个 case 的第 m 个主成分得分)

for some constants $\phi_{m1}, \dots, \phi_{mp}$.

- We can then fit the linear regression model,

$$X: \begin{pmatrix} Z_{11} & Z_{12} & \dots & Z_{1m} \\ Z_{21} & Z_{22} & \dots & \vdots \\ \vdots & \vdots & \ddots & \vdots \\ & & & Z_{n1} & Z_{n2} & \dots & Z_{nm} \end{pmatrix}_{n \times m} \quad y_i = \theta_0 + \sum_{m=1}^M \theta_m z_{im} + \epsilon_i, \quad i = 1, \dots, n, \quad (2)$$

各主成分得分的系数 (m < p)
且解决共线问题 \rightarrow var

优势: 预测效果好
缺点: 不易解释

using ordinary least squares.

- Note that in model (2), the regression coefficients are given by $\theta_0, \theta_1, \dots, \theta_M$. If the constants $\phi_{m1}, \dots, \phi_{mp}$ are chosen wisely, then such dimension reduction approaches can often outperform OLS regression.

- Notice that from definition (1),

$$\sum_{m=1}^M \theta_m z_{im} = \sum_{m=1}^M \theta_m \sum_{j=1}^p \phi_{mj} x_{ij} = \sum_{j=1}^p \sum_{m=1}^M \theta_m \phi_{mj} x_{ij} = \sum_{j=1}^p \beta_j x_{ij},$$

where

$$\boxed{\beta_j = \sum_{m=1}^M \theta_m \phi_{mj}.}$$

可将用主成分回归
后得到的参数还原为 β

(3)

- Hence model (2) can be thought of as a special case of the original linear regression model.
- Dimension reduction serves to constrain the estimated β_j coefficients, since now they must take the form (3).
- Can win in the bias-variance tradeoff. 相当于对 β 进行约束


Principal Components Regression



- Here we apply principal components analysis (PCA) (discussed in Chapter 10 of the text) to define the linear combinations of the predictors, for use in our regression.
- The first principal component is that (normalized) linear combination of the variables with the largest variance.
- The second principal component has largest variance, subject to being uncorrelated with the first.
- And so on.
- Hence with many correlated original variables, we replace them with a small set of principal components that capture their joint variation.



用力量主成分表示原数据的方差 .

Pictures of PCA





The population size (**pop**) and ad spending (**ad**) for 100 different cities are shown as purple circles. The green solid line indicates the first principal component, and the blue dashed line indicates the second principal component.

Pictures of PCA: continued





A subset of the advertising data. Left: The first principal component, chosen to minimize the sum of the squared perpendicular distances to each point, is shown in green. These distances are represented using the black dashed line segments. Right: The left-hand panel has been rotated so that the first principal component lies on the x-axis.

Pictures of PCA: continued





*Plots of the first principal component scores z_{i1} versus **pop** and **ad**. The relationships are strong.*

Pictures of PCA: continued




*Plots of the second principal component scores z_{i2} versus **pop** and **ad**. The relationships are weak.*

Application to Principal Components Regression



PCR was applied to two simulated data sets. The black, green, and purple lines correspond to squared bias, variance, and test mean squared error, respectively. **Left:** Simulated data from slide 32. **Right:** Simulated data from slide 39.

Choosing the number of directions M



Left: PCR standardized coefficient estimates on the Credit data set for different values of M . **Right:** The 10-fold cross validation MSE obtained using PCR, as a function of M .

Partial Least Squares

- PCR identifies linear combinations, or *directions*, that best represent the predictors X_1, \dots, X_p .
- These directions are identified in an *unsupervised* way, since the response Y is not used to help determine the principal component directions.
- That is, the response does not *supervise* the identification of the principal components.
- Consequently, PCR suffers from a potentially serious drawback: there is no guarantee that the directions that best explain the predictors will also be the best directions to use for predicting the response.

Partial Least Squares: continued

- Like PCR, PLS is a dimension reduction method, which first identifies a new set of features Z_1, \dots, Z_M that are linear combinations of the original features, and then fits a linear model via OLS using these M new features.
- But unlike PCR, PLS identifies these new features in a supervised way – that is, it makes use of the response Y in order to identify new features that not only approximate the old features well, but also that *are related to the response*.
- Roughly speaking, the PLS approach attempts to find directions that help explain both the response and the predictors.

Details of Partial Least Squares

- After standardizing the p predictors, PLS computes the first direction Z_1 by setting each ϕ_{1j} in (1) equal to the coefficient from the simple linear regression of Y onto X_j .
- One can show that this coefficient is proportional to the correlation between Y and X_j .
- Hence, in computing $Z_1 = \sum_{j=1}^p \phi_{1j} X_j$, PLS places the highest weight on the variables that are most strongly related to the response.
- Subsequent directions are found by taking residuals and then repeating the above prescription.

Summary

- Model selection methods are an essential tool for data analysis, especially for big datasets involving many predictors.
- Research into methods that give *sparsity*, such as the *lasso* is an especially hot area.
- Later, we will return to sparsity in more detail, and will describe related approaches such as the *elastic net*.