


Effect size: raw vs covariate-adjusted



M57: Covariate Analysis - Effect Survival Table

Covariate: final_answer_length

Measure	Raw d	Raw p	Adj d	Adj p	Atten	Survival
seq_typing_max_run	-0.41	0.043	-0.52	0.013	-27%	SURVIVED
seq_typing_entropy	+0.52	0.012	+0.69	0.001	-32%	SURVIVED
seq_typing_mean_run_explore	-0.32	0.122	-0.44	0.038	-38%	SURVIVED
seq_typing_mean_run_exploit	+0.03	0.884	+0.13	0.547	-327%	KILLED
pasted_chars	-0.42	0.043	+0.16	0.429	61%	KILLED
n_paste_events	-0.29	0.148	-0.01	0.977	98%	KILLED
seq_time_mean_run_explore	-0.35	0.091	-0.38	0.075	-9%	PARTIAL
seq_topic_mean_run_exploit	-0.32	0.110	-0.42	0.043	-31%	SURVIVED
first_writing_time_s	-0.32	0.115	-0.47	0.025	-48%	SURVIVED
prop_pages_with_writing	+0.05	0.790	+0.12	0.561	-125%	KILLED


Survival rules:

SURVIVED -> adjusted p < .05 AND |adj d| >= 0.3
PARTIAL -> |adj d| >= 50% of |raw d| AND |adj d| >= 0.2
KILLED -> effect mostly attributable to covariate

Adj d denominator = sqrt(MSE) from full model (different scale than raw pooled SD; absolute comparison is approximate).


seq_typing_entropy

raw $d=+0.52$ ($p=0.012$) -> adj $d=+0.69$ ($p=0.001$) [SURVIVED]




pasted_chars

raw $d=-0.42$ ($p=0.043$) -> adj $d=+0.16$ ($p=0.429$) [KILLED]



seq_typing_max_run

raw d=-0.41 (p=0.043) -> adj d=-0.52 (p=0.013) [SURVIVED]



seq_time_mean_run_explore

raw d=-0.35 (p=0.091) -> adj d=-0.38 (p=0.075) [PARTIAL]

