Looking Away

Building 1: Moves

Arrows from a floor to itself, and what counting their "hows" gives.

Contents

Part I. Moves on things 2

Chapter 1. Rearrangements a bag of n things → its moves 2

Chapter 2. Words letters and their undos → the free move system 5

Part II. Move systems as rows 8

Chapter II.1. The table of move systems move systems → classes 8

Chapter II.2. Blurs that respect one side a move system → its cosets 11

Chapter II.3. Blurs that respect both sides a move system → its quotients 13

Chapter II.4. Moves on matrices: the Smith form relation matrices → their normal form 15

Part III. A cubic and its moves 18

Chapter III.1. Points glued by lines sums of points → points 18

Chapter III.2. The torus carries the law $\mathbb{C}/\Lambda \to C(\mathbb{C})$ 21

Chapter III.3. Counting points over a clock cubics over $\mathbb{Z}/p$ → their sizes 23

A *move* is an arrow from a floor to itself that is an inside and a picture (Chapter 0, W2). Forgetting *how* one thing moves to another gives a blur: Building 0. This book keeps the how. It reads Building 0 and is never read by it (the rule F7, between buildings).

Chapter numbers restart in this book; Building 0 is cited as "Building 0, Chapter n". The rules are Chapter 0 of Building 0.

Planned, not yet written (`plans/ALGEBRA-PATH.md`). Part II: Actions; Counting and Sylow; Divisibility in any ring; Linear maps; Similarity; Forms; Field extensions; Finite fields; the Galois correspondence; Solvable ladders and radicals. The plane, the cubic and the torus are in Building 0 (Chapters 35–37); Part III here uses them.

Part I. Moves on things

Chapter 1. Rearrangements

a bag of n things $\to$ its moves · from: Building 0, Chapters 0, 1, 3 · build, select, blur

Select the moves of n things. Glue all swaps together and exactly two bags are left: the sign appears. Glue a move to its relabellings and the bags are the shapes of its cycles.

The first chapter that keeps the *how*. A move is an arrow from a bag to itself that is an inside and a picture (Chapter 0, W2). Building 0 forgot how one thing was carried to another; here the carrying is the thing.

Given

G1 (Building 0, Chapter 0.) Build M1–M7; blur L1–L9; arrows W1–W4; W2: inside and picture.

G2 (Building 0, Chapter 3.) $\mathbb{Z}/2$ with $+$.

G3 (Building 0, Chapter 1.) Whole numbers; counting a finite bag.

Definitions

D1 (Build, M7; Select, M4.) Fix $n \geq 1$ and $[n] := \{1, \ldots, n\}$. S_n: the maps $\sigma : [n] \to [n]$ that are one-to-one (then also onto). Each is a *move* of $[n]$. $\sigma\tau := \sigma \circ \tau$ (do τ first). e: the move that does nothing.

D2 A *swap* $(i\,j)$ exchanges i and j and keeps the rest. A *cycle* $(i_1\, i_2 \ldots i_k)$ sends each i_r to i_{r+1} and i_k to i_1.

D3 (Inversions.) $\mathrm{inv}(\sigma)$: the number of pairs $i < j$ with $\sigma(i) > \sigma(j)$.

D4 (Relabelling.) For π in S_n: $\sigma^\pi := \pi\sigma\pi^{-1}$, the move σ with the things renamed by π.

Propositions: the floor

P1 (Laws, carried by W2.) $\circ$ is associative; e does nothing on either side; every move has an undo σ^{-1}. Not commutative for $n \geq 3$: $(1\,2)(2\,3) = (1\,2\,3)$ while $(2\,3)(1\,2) = (1\,3\,2)$.

P2 (Count.) S_n has $n!$ members.

P3 (Normal form: cycles.) Every move is a product of cycles on disjoint sets of things, in exactly one way up to the order of the cycles (follow each thing until it comes back).

P4 (Swaps are enough.) Every move is a product of swaps, by P3 and

$$(i_1 \ldots i_k) = (i_1\, i_2)(i_2\, i_3) \cdots (i_{k-1}\, i_k) \quad \text{(do the right-hand swap first).}$$

Blur 1: the sign

On S_n, $n \geq 2$. *O*: $\circ$.

B1 (From below.) Gluings: $s \sim t$ for any two swaps s, t.

B2 (From above.) Question: the parity of $\mathrm{inv}(\sigma)$, in $\mathbb{Z}/2$.

P5 (Survives: parity respects $\circ$.) $\mathrm{inv}(\sigma\tau) \equiv \mathrm{inv}(\sigma) + \mathrm{inv}(\tau) \pmod 2$. (For each pair $\{i, j\}$, $\sigma\tau$ turns it over exactly when one of τ on $\{i, j\}$ and σ on $\{\tau i, \tau j\}$ does; count the pairs mod 2.) A swap has an odd number of inversions: $(i\,j)$ with $i < j$ has $2(j - i) - 1$.

P6 (Appears.) The two blurs agree. There are exactly two bags, A_n (even) and its other bag (odd). (B1 puts every product of an even number of swaps with e and every odd product with $(1\,2)$, by P4; B2 tells them apart, by P5.) Name: $\mathrm{sgn}(\sigma) := (-1)^{\mathrm{inv}(\sigma)}$.

P7 (Dies: how many swaps.) $(1\,3) = (1\,2)(2\,3)(1\,2)$: one swap, or three. The number of swaps is not a question on moves; only its parity is.

P8 (Survives into Building 0.) $\mathrm{sgn} : S_n \to \{1, -1\}$ respects $\circ$ and $\times$: an arrow onto Building 0's $\mathbb{Z}/2$ (G2), written multiplicatively. By W4, $S_n \to \mathbb{Z}/2$ is the blur B1 followed by a matching.

Blur 2: relabelling

On S_n. *O*: nothing (L7: this blur respects neither $\circ$ nor its undo in general; say so).

B3 (From below.) Gluings: $\sigma \sim \sigma^\pi$ for every π.

B4 (From above.) Question: the *cycle type*, the list of cycle lengths of P3 in falling order (a partition of n).

P9 (Appears.) The two blurs agree. $(\pi(i_1 \dots i_k)\pi^{-1} = (\pi i_1 \dots \pi i_k)$: relabelling renames the cycles and keeps their lengths; two moves of the same type are relabellings of each other.) Normal form: one move per partition of n; the bags number $p(n)$: $1, 2, 3, 5, 7, 11$ for $n = 1, \dots, 6$.

P10 (Dies: $\circ$.) $(1\,2) \sim (2\,3)$, but $(1\,2)(1\,2) = e$ while $(2\,3)(1\,2) = (1\,3\,2)$.

P11 (Survives: sign and order.) $\mathrm{sgn}(\sigma) = (-1)^{n-c}$ with c the number of cycles (counting fixed things); the order of σ (least $m \geq 1$ with $\sigma^m = e$) is the lcm of the cycle lengths. Both are read off the type.

P12 (Bag sizes.) The bag of type $1^{m_1} 2^{m_2} \cdots$ has $n! \,/\, \prod_k k^{m_k}\, m_k!$ members. (Count the relabellings that give the same move.)

Theorems

T1 (Survey: every blur of S_n that respects $\circ$.) A blur of S_n respecting $\circ$ is fixed by its bag of e, which is a union of bags of Blur 2 closed under $\circ$. For $n \leq 6$ the survey is:
$n = 2$: two blurs. $n = 3$: three (e only; A_3; everything). $n = 4$: four (e; the three double swaps with e; A_4; everything). $n = 5$ and $n = 6$: three (e; A_n; everything).

Machine-checked (all unions of types closed under $\circ$). For $n \geq 5$ the classical theorem is that A_n glues to nothing smaller: A_n is *simple*. That is a count, proved in Part II.

T2 (The sign is the only arrow to $\mathbb{Z}/2$.) For $n \geq 2$, the arrows $S_n \to \{1, -1\}$ respecting $\circ$ are $\sigma \mapsto 1$ and sgn. (Relabelling glues all swaps (P9), so an arrow takes one value on them; P4.)

T3 (Every move system lives here. Cayley.) For any finite bag G of moves of something, closed under $\circ$ and undo, $g \mapsto (h \mapsto gh)$ is an inside of G into the moves of the bag G, respecting $\circ$. So every finite move system is found inside some S_n, $n =$ its size.

Remarks

R1 (What the blur did.) The sign is an Appears line: "glue all swaps" from below meets "parity of inversions" from above. That the bags are two and not one is the kept question P5, and it is the whole content of the classical theorem.

R2 (Building 0 seen from here.) The blur of Building 0, Chapter 3 ($n \sim 0$ on $\mathbb{Z}$) is "forget how far you moved" for the moves $x \mapsto x + k$. Every blur in Building 0 is such a forgetting; this book keeps the moves.

R3 (Old names.) S_n: the symmetric group. A_n: the alternating group. P3: disjoint cycle decomposition. P6: the sign is well defined. B3: conjugacy; its bags are conjugacy classes. P12: class sizes. T1: the normal subgroups of S_n. T3: Cayley's theorem.

R4 (Machine check, `checks/b1-ch1-check.py`.) P1, P2, P4 on all of S_n, $n \leq 6$; P5 for $n \leq 5$; P6 (closure of the swap gluing); P7; P9 and P12 (types and bag sizes); P11; T1 (survey); T2.

Exercises

E1 Write $\sigma = (1\,4\,2)(3\,5)$ in S_5 as swaps, and find $\operatorname{sgn}\sigma$ and the order of σ in two ways.

E2 Count the moves of type $2\,2\,1$ in S_5 by P12, and by listing.

E3 Show that in S_n the even moves are exactly the products of 3-cycles ($n \geq 3$).

E4 Which move of S_4 has the most inversions? How many does it have?

E5 Why is "σ and τ commute" not kept by Blur 2? Give a witness in S_3.

Chapter 2. Words

letters and their undos → the free move system · from: Building 0, Chapters 0, 1, 3; Chapter 1 · build, blur

Write words in letters and their undos, and glue a letter next to its undo to nothing. Each bag holds exactly one word with nothing left to cancel. Glue more, and every move system appears.

Chapter 1 started from moves already given. This chapter builds moves from nothing but names, then blurs. It is the build-then-blur shape of Building 0, Chapter 1, one level up.

Given

G1 (Building 0, Chapter 0, M3.) Words in a list of names; maps out of words by recursion.

G2 (Chapter 1.) Moves of a bag; $\circ$; P1.

G3 (Building 0, Chapters 1, 3.) $\mathbb{Z}$ and $\mathbb{Z}/n$ with $+$.

Definitions

D1 (Build, M3.) Fix a bag L of *letters*. For each letter s take a second name s^{-1}. $W(L)$: the finite strings in the names s, s^{-1}, with the empty string 1. Joining strings, $u \cdot v$, is the operation.

D2 A word is *reduced* if no s stands next to s^{-1} (in either order).

D3 (Each letter is a move of the reduced words.) For a reduced word r: $s * r := r$ with its first name removed if that name is s^{-1}, and sr otherwise; $s^{-1} * r$ in the same way. Each of these is a move (its undo is the other), so a word $u = x_1 \cdots x_m$ acts by $u * r := x_1 * (\cdots (x_m * r))$.

The blur

On $W(L)$. O: joining.

B1 (From below.) Gluings: $s\, s^{-1} \sim 1$ and $s^{-1} s \sim 1$, for each letter s.

B2 (From above.) Question: the move $r \mapsto u * r$ of the reduced words (D3).

Propositions

P1 (Appears.) The two blurs agree. (B1 keeps the question: $s * (s^{-1} * r) = r$. Every word is glued by B1 to a reduced one, by cancelling. Two reduced words $r \neq r'$ act differently: $r * 1 = r$.)

P2 (Normal form.) Each bag holds exactly one reduced word. (P1.) The bags, with joining, form a move system $F(L)$: the *free move system* on L. Undo: $(x_1 \cdots x_m)^{-1} = x_m^{-1} \cdots x_1^{-1}$.

P3 (Count.) With two letters, the reduced words of length $k \geq 1$ number $4 \cdot 3^{k-1}$: four choices for the first name, then three (anything but the undo of the one before).

P4 (Dies: length.) $s\,s^{-1}s \sim s$, of lengths 3 and 1.

P5 (Survives: length mod 2, and each letter's exponent sum.) Cancelling removes two names, one s and one s^{-1}. So $\ell(u) \bmod 2$ and $\varepsilon_s(u) :=$ (number of s) $-$ (number of s^{-1}) survive: arrows $F(L) \to \mathbb{Z}/2$ and $F(L) \to \mathbb{Z}$, respecting joining and $+$.

P6 (One letter.) $F(\{s\}) \to \mathbb{Z}$, $u \mapsto \varepsilon_s(u)$, is a matching. Building 0, Chapter 1 builds $\mathbb{Z}$ from pairs; here it is the free move system on one letter.

Theorems

T1 (Every move system is a picture of a free one.) Let G be a move system and choose g_s in G for each letter s. There is exactly one arrow $F(L) \to G$ respecting joining with $s \mapsto g_s$. (Recursion on words, G1; it respects B1 because $g_s g_s^{-1} = e$.) If the g_s generate G, the arrow is a picture, and by W4

$$G \text{ matches } F(L) \text{ blurred by the bag of words sent to } e.$$

T2 (Presentations: glue more.) For a list R of words, glue $r \sim 1$ for each r in R, respecting joining. Write the bags $\langle L \mid R \rangle$. Then the bag of 1 is the smallest bag that holds R, joins, undos, and relabellings uru^{-1} (as Chapter 1, T1: a zero bag must be kept by relabelling).

T3 (Old floors, as presentations.)
$\langle s \mid s^n \rangle$ matches $\mathbb{Z}/n$ (Building 0, Chapter 3).
$\langle s, t \mid sts^{-1}t^{-1} \rangle$ matches $\mathbb{Z}^2$: glue $st \sim ts$ and the normal form is $s^a t^b$ (the whole-number points of $\mathbb{R}^2$, Building 0, Chapter 6).
$\langle s, t \mid s^2, t^2, (st)^3 \rangle$ matches S_3 (Chapter 1), with $s = (1\,2)$, $t = (2\,3)$.
$\langle s, t \mid s^2, t^2, (st)^n \rangle$ has $2n$ members: the moves of a regular n-gon.
$\langle s, t \mid s^2, t^3, (st)^5 \rangle$ has 60 members.

T4 (Dies: deciding.) For some finite L and R, no procedure decides whether a word lies in the bag of 1 (classical, Novikov and Boone; reported). From below, a blur can be written down and still not be computable: Building 0 never met this, because its gluings always came with a normal form.

Remarks

R1 (Build, then blur, one level up.) Building 0, Chapter 1: pairs, glue $(a+1, b+1) \sim (a, b)$, normal form $a - b$. Here: words, glue $ss^{-1} \sim 1$, normal form the reduced word. The question that makes the blur from above (D3) is itself a move: in Building 1 even the questions are moves.

R2 (Where it goes.) T1 makes every move system a blur of a free one. Part II asks which blurs of a move system respect joining (cosets, normal bags), and counts bags.

R3 (Old names.) $F(L)$: the free group on L. D3 and P1: van der Waerden's proof of the normal form. P5: abelianisation to $\mathbb{Z}^L$ (exponent sums). T1: the universal property of free groups. T2: group presentations; the bag of 1 is the normal closure of R. T3: cyclic, free abelian, symmetric, dihedral groups; the last is A_5. T4: the word problem.

R4 (Machine check, `checks/b1-ch2-check.py`.) P1 on all words of length ≤ 8 in two letters; P3 to length 8; P5; T3 (orders by coset enumeration, and the arrow onto S_3).

Exercises

E1 Reduce $s\,t\,t^{-1}s^{-1}s\,t^{-1}\,t\,s$.

E2 Show that in $F(\{s,t\})$, $st \neq ts$, using the exponent sums or P2.

E3 In $\langle s,t \mid s^2, t^2, (st)^3\rangle$, show $sts = tst$.

E4 How many reduced words of length at most 3 are there in two letters?

E5 Why is "length is even" kept by B1 but "length is ≤ 5" not?

Part II. Move systems as rows

Chapter II.1. The table of move systems

move systems → classes · from: Part I; Building 0, Chapters 1, 3, 8, 9, 14, II.1 · table, blur, class

The rows are now move systems. Every row passes the three laws of moves; the table is about what else they pass.

Building 0, Chapter II.1 again, with move systems for rows. A *move system* is a bag G with $\circ$, a do-nothing e and undos, where $\circ$ is associative: exactly the laws Chapter 1, P1 found for the moves of a bag. By Chapter 1, T3 every finite one is a bag of moves.

Given

G1 (Rows.) $\mathbb{Z}$ with $+$ (Building 0, Ch. 1); $\mathbb{Z}/6$ (Ch. 3); the circle $\mathbb{T}$ (Ch. 8); $\mathbb{Q}^\times$, the non-zero rationals with $\times$ (Ch. 2); $(\mathbb{Z}/8)^\times = \{1,3,5,7\}$ with $\times$ (Ch. 3); S_3 (Chapter 1); D_4, the moves of S_4 that carry a square's corners $1,2,3,4$ (in order round the square) to corners of the square; $GL_2(\mathbb{R})$, the straight maps of $\mathbb{R}^2$ with $\det \neq 0$ (Building 0, Ch. 14); F_2 (Chapter 2); $\mathbb{Z}^2$ (Chapter 2, T3).

G2 (Facts.) The lines of those chapters, read as yes/no facts.

Tests

D1 comm: $gh = hg$ for all g, h.

D2 fin: finitely many members.

D3 cyc: some g has every member as a power g^k, k in $\mathbb{Z}$.

D4 tf: no $g \neq e$ has $g^k = e$ for any $k \geq 1$.

D5 div: for every g and every $n \geq 1$ there is h with $h^n = g$.

The table

	comm	fin	cyc	tf	div
$\mathbb{Z}$	✓	·	✓	✓	·
$\mathbb{Z}/6$	✓	✓	✓	·	·
$\mathbb{T}$	✓	·	·	·	✓
$\mathbb{Q}^\times$	✓	·	·	·	·
$(\mathbb{Z}/8)^\times$	✓	✓	·	·	·
S_3	·	✓	·	·	·
D_4	·	✓	·	·	·
$GL_2(\mathbb{R})$	·	·	·	·	·
F_2	·	·	·	✓	·
$\mathbb{Z}^2$	✓	·	·	✓	·

Reasons for the less obvious entries: $\mathbb{T}$ is not cyc (if a turns a rational amount its powers are finitely many; if not, $a/2$ is no power of a) and is div (Building 0, Chapter 8, T2). $\mathbb{Q}^\times$: -1 has $(-1)^2 = 1$; 2 has no square root. $(\mathbb{Z}/8)^\times$: every member squares to 1. $GL_2(\mathbb{R})$: $-I$ squares to I; $\mathrm{diag}(-1, 1)$ has no square root (det of a square is ≥ 0). F_2: s has no square root (exponent sums, Chapter 2, P5); no power of a reduced word $\neq 1$ is 1.

Blur, and what appears

B1 (Building 0, II.1, B1.) For a set U of tests, glue two rows with the same answers on U.

P1 (Not sharp.) With all five tests, S_3 and D_4 share a bag. A separating test is the size: 6 against 8. Every other pair of rows is told apart.

P2 (Classes.) As T runs over the sets of tests, class(T) takes 11 values. Old names: class(comm), *abelian*; class(cyc), *cyclic* ($\mathbb{Z}$, $\mathbb{Z}/6$); class(tf), *torsion-free* ($\mathbb{Z}$, F_2, $\mathbb{Z}^2$).

P3 (Read off the table.) cyc $\Rightarrow$ comm; div $\Rightarrow$ comm. Never together: fin and tf; fin and div; cyc and div; tf and div.

Theorems

For every move system with more than one member, not only the ten rows.

T1 cyc $\Rightarrow$ comm. ($g^j g^k = g^{j+k} = g^k g^j$.)

T2 fin and tf never together. (The powers of g repeat, $g^i = g^j$ with $i < j$, so $g^{j-i} = e$.)

T3 fin and div never together. (By T2's reason each g has $g^m = e$ for some $1 \leq m \leq |G|$; so with $n = |G|!$ every n-th power is e, and $g \neq e$ has no n-th root.)

T4 cyc and div never together. (Finite: T3. Infinite: it matches $\mathbb{Z}$, and $2h = 1$ has no answer there.)

T5 (Matching rows.) $\mathbb{Z}/6$ matches $\mathbb{Z}/2 \times \mathbb{Z}/3$, and $(\mathbb{Z}/8)^\times$ matches $\mathbb{Z}/2 \times \mathbb{Z}/2$; $\mathbb{Z}/6$ and $(\mathbb{Z}/8)^\times$ differ already by size, and $\mathbb{Z}/4$ and $(\mathbb{Z}/8)^\times$ by cyc.

Remarks

R1 (Accidents of a small table; Building 0, II.1, R1.) Two lines of P3 die when the right row is added. div $\Rightarrow$ comm dies on the rotations of space: every rotation has n-th roots (turn by a $1/n$ of the angle about the same axis), and two rotations about different axes need not commute. "tf and div never together" dies on $\mathbb{Q}$ with $+$. The other four survive every row: T1–T4.

R2 (A conjecture from the table, F6.) Every abelian row with fin in the table matches a product of clocks $\mathbb{Z}/n_1 \times \cdots \times \mathbb{Z}/n_k$. This is true for every finite abelian move system; it is proved in the chapter on modules, from a normal form of matrices.

R3 (Old names.) Move system: group. comm: abelian. tf: torsion-free. div: divisible. D_4: the dihedral group of order 8. T2: finite groups are torsion groups.

R4 (Machine check, `checks/b1-p2-ch1-check.py`.) Every finite entry by brute force; the 11 classes (`b1-p2-ch1-classes.py`); P1; T5 (the two matchings checked).

Exercises

E1 List the members of D_4 as moves of the four corners, and show D_4 is not comm.

E2 Add the row $\mathbb{Q}$ with $+$. How many classes now?

E3 Find a test that separates S_3 from D_4 without counting members.

E4 Show $\mathbb{Z}^2$ is not cyc.

Chapter II.2. Blurs that respect one side

a move system → its cosets · from: Chapter II.1; Building 0, Chapters 3, II.2 · blur, count

Ask a blur to respect moving on the left only. Its bags are copies of the bag of e, all of one size. Counting them gives Lagrange, and Fermat's little theorem falls out.

Building 0, II.2 asked which blurs of a ring respect $+$ and $\times$. Here the question is asked of a move system with less: respect $g\circ$ on the left, and nothing on the right. Rule L7 (say what the blur respects) is the whole story of this chapter and the next.

Given

G1 (Chapter II.1.) Move systems; the rows.

G2 (Building 0, Chapter 3.) $\mathbb{Z}/n$; $(\mathbb{Z}/n)^\times$, the bags with an inverse for $\times$; Chapter 3, T5 there.

The blur

On a move system G. O: for each g, the move $x \mapsto gx$ (left only).

B1 (From below.) For a bag $H \subseteq G$: glue $h \sim e$ for each h in H.

B2 (From above.) For a sub move system K (closed under $\circ$ and undo, holding e): the question $x \mapsto$ the bag $xK := \{xk : k \in K\}$.

Propositions

P1 (Survey: every such blur.) A blur of G respecting left moves is fixed by its bag of e, which is a sub move system K; and $x \sim y \iff x^{-1}y \in K$. Conversely every sub move system gives one. So these blurs and the sub move systems match.

P2 (Appears.) B1 for H equals B2 for $K =$ the smallest sub move system holding H. Its bags are the xK: the *left cosets*.

P3 (Survives: moving a bag.) $x \mapsto gx$ carries the bag yK onto the bag gyK, one-to-one. (Its undo is $x \mapsto g^{-1}x$.) So all bags have the same size as K.

P4 (Dies: the right side.) In S_3 with $K = \{e, (1\,2)\}$: $e \sim (1\,2)$, but $e(1\,3) = (1\,3)$ and $(1\,2)(1\,3) = (1\,3\,2)$ are in different bags, since $(1\,3)^{-1}(1\,3\,2) = (2\,3) \notin K$.

P5 (Normal form.) None in general: no rule picks one member of each xK. For $\mathbb{Z}$ and $K = n\mathbb{Z}$ the normal forms $0, \ldots, n-1$ of Building 0, Chapter 3 are such a rule.

Theorems

T1 (Counting bags: Lagrange. A count, not a Survives line.) If G is finite, $|G| = |K| \cdot$ (number of bags). (P3: the bags are disjoint and each has $|K|$ members.) So $|K|$ divides $|G|$.

T2 (The order of a move divides the size.) For g in a finite G, the least $m \geq 1$ with $g^m = e$ divides $|G|$. ($K = \{e, g, \ldots, g^{m-1}\}$ has m members: T1.)

T3 ($g^{|G|} = e$.) From T2.

T4 (Building 0, Chapter 3, T5, recovered.) For a with $\gcd(a, n) = 1$: $a^{\varphi(n)} \equiv 1 \pmod{n}$, where $\varphi(n) = |(\mathbb{Z}/n)^\times|$. For $n = p$ prime: $a^{p-1} \equiv 1$. (T3 in $(\mathbb{Z}/n)^\times$.)

T5 (Dies: "every divisor is a size".) A_4 has 12 members and no sub move system with 6. (Machine-checked; classical.) T1 is "at most these sizes", never "these sizes occur".

T6 (Prime size.) If $|G| = p$ is prime, G matches $\mathbb{Z}/p$. (Any $g \neq e$ has order dividing p, so p: its powers are all of G.)

Remarks

R1 (What the blur did.) The cosets are the bags of the one blur that respects left moves; their equal size is a Survives line (P3); Lagrange is a count of bags (T1). The method's own split: blur keeps (P3), counting gives existence and sizes.

R2 (Respecting both sides.) If the blur also respects right moves, K must be kept by relabelling, $gKg^{-1} = K$, and the bags themselves form a move system. That is Chapter II.3.

R3 (Old names.) Left cosets; index = number of bags; T1: Lagrange's theorem; T4: Euler's theorem, Fermat's little theorem; T5: the converse of Lagrange fails (A_4).

R4 (Machine check, `checks/b1-p2-ch2-check.py`.) P1 (for every bag $K \ni e$ of S_3 and D_4: "$x^{-1}y \in K$" is a blur exactly when K is a sub move system); P3; P4; T1–T3 on all sub move systems of S_4; T4 for $n \leq 200$; T5.

Exercises

E1 List the left cosets of $K = \{e, (1\,2)\}$ in S_3, and then the right cosets Kx. Are they the same bags?

E2 Find $3^{100} \bmod 7$ with T4.

E3 Which sizes of sub move systems occur in S_4? Compare with the divisors of 24.

E4 Show that a move system of size 4 is comm. (Each $g \neq e$ has order 2 or 4.)

Chapter II.3. Blurs that respect both sides

a move system → its quotients · from: Chapters 1, 2, II.2 · blur

Ask the blur to respect moving on both sides. The bag of e must now be kept by relabelling, and the bags form a move system of their own. Every arrow is such a blur followed by an inside.

The same gluing as Chapter II.2, with one more operation in O. Building 0, Chapter 3 is the case $G = \mathbb{Z}$, where left and right are the same and every sub move system will do.

Given

G1 (Chapter II.2.) Left-respecting blurs and cosets; T1 (Lagrange).

G2 (Chapter 1.) Relabelling σ^π; the bags of relabelling (types); T1 (survey for S_n, $n \leq 6$).

G3 (Chapter 2.) Presentations; P5 (exponent sums).

The blur

On a move system G. O: $x \mapsto gx$ and $x \mapsto xg$ for each g.

B1 (From below.) For a bag H: glue $h \sim e$ for each h in H.

B2 (From above.) For a sub move system N with $gNg^{-1} = N$ for every g (*kept by relabelling*): the question $x \mapsto xN$.

Propositions

P1 (Survey.) A blur respecting both sides is a left-respecting blur (Chapter II.2, P1) whose bag of e, N, is kept by relabelling: $n \sim e$ gives $gng^{-1} \sim gg^{-1} = e$. Conversely each such N gives one. Then $xN = Nx$: left and right bags agree.

P2 (Appears.) B1 for H equals B2 for the smallest N kept by relabelling that holds H (Chapter 2, T2).

P3 (Survives: $\circ$ on bags.) $xN \cdot yN := xyN$ does not depend on the members chosen. The bags form a move system G/N, and $x \mapsto xN$ is a picture respecting $\circ$.

P4 (Dies: "kept by relabelling" does not chain.) In D_4 (corners $1, 2, 3, 4$ round a square), let s be the flip $(2\,4)$ and $V = \{e, s, (1\,3), (1\,3)(2\,4)\}$. Then $\{e, s\}$ is kept by relabelling in V (V is comm), and V is kept in D_4 (it has half the members). But $\{e, s\}$ is not kept in D_4: the quarter turn $r = (1\,2\,3\,4)$ gives $rsr^{-1} = (1\,3)$.

P5 (Half is always kept.) If N has half the members of G, it is kept by relabelling: the two left bags and the two right bags are both N and the rest.

Theorems

T1 (Every arrow is a blur, then an inside. W4.) For an arrow $f : G \to G'$ respecting $\circ$, the blur "$f(x) = f(y)$" respects both sides; its bag of e is $N = \{x : f(x) = e\}$;

and $G/N \to G'$, $xN \mapsto f(x)$, is an inside. So G/N matches the picture of f.
Example: sgn : $S_n \to \{1, -1\}$ gives S_n/A_n matching $\mathbb{Z}/2$ (Chapter 1, P8).

T2 (Blurs of a blur.) The two-sided blurs of G/N match the two-sided blurs of G that are blurrier than N's, by $M/N \leftrightarrow M$; and $(G/N)/(M/N)$ matches G/M. Blurring twice is blurring once, with the coarser bag of e.

T3 (A sub and a blur.) For a sub move system K and N kept by relabelling: KN is a sub move system, $K \cap N$ is kept by relabelling in K, and KN/N matches $K/(K \cap N)$. (T1 for $K \to G/N$.)

T4 (Appears: the blur to comm.) Glue $xy \sim yx$ for all x, y, respecting both sides. The bags form a comm move system G^{ab}, and every arrow from G to a comm move system passes through it, in exactly one way. From above: the questions are all arrows to comm systems.
S_n^{ab} matches $\mathbb{Z}/2$, by sgn ($n \geq 2$). F_2^{ab} matches $\mathbb{Z}^2$, by the exponent sums (Chapter 2, P5). D_4^{ab} matches $\mathbb{Z}/2 \times \mathbb{Z}/2$.

T5 (Simple: only two such blurs. A count.) G is *simple* if its only two-sided blurs are "glue nothing" and "glue everything". $\mathbb{Z}/p$ is simple (Chapter II.2, T1). A_5 is simple: its relabelling bags (inside A_5) have sizes $1, 15, 20, 12, 12$; a bag of e kept by relabelling is a union of these holding the 1, and its size divides 60 (Lagrange); the only such unions have 1 or 60 members.

Remarks

R1 (L7 is the whole story.) The same gluing $h \sim e$, with left moves in O, gives cosets; with both sides, quotients. Nothing else changed.

R2 (Simple pieces.) A ladder of blurs $G \to G/N_1 \to \cdots \to \{e\}$ in which each rung's blur is simple lists the *atoms* of G (the atoms ladder of the project, rung 2: the list survives, the gluing does not). A_5 is the first atom that is not a clock.

R3 (Old names.) Kept by relabelling: normal subgroup. G/N: quotient group. T1: the first isomorphism theorem. T2: the correspondence and third isomorphism theorems. T3: the second isomorphism theorem. T4: abelianisation, $G/[G, G]$. T5: simple groups; A_5 simple. R2: Jordan–Hölder.

R4 (Machine check, `checks/b1-p2-ch3-check.py`.) P1 on all bags of S_3, D_4 (and kept-by-relabelling sub systems of S_4); P3; P4; P5 on S_4; T1 for sgn and det on $GL_2(\mathbb{Z}/3)$; T4 sizes for S_3, S_4, D_4; T5 (class sizes of A_5, all unions).

Exercises

E1 List the sub move systems of S_3 kept by relabelling.

E2 Show that every sub move system of a comm system is kept by relabelling. Which line of Building 0, Chapter 3 is this?

E3 Find S_3^{ab} by gluing $(1\,2)(2\,3) \sim (2\,3)(1\,2)$ and closing up.

E4 Why can A_5 have no arrow onto $\mathbb{Z}/2$?

Chapter II.4. Moves on matrices: the Smith form

relation matrices → their normal form · from: Chapters II.1, II.3; Building 0, Chapters 18, II.2, II.5 · blur, build

Glue a matrix to every matrix reached by moving its rows and columns. The kept questions are the gcds of the minors; the normal form is a diagonal in which each entry goes into the next. Every finite comm move system is then a product of clocks.

The first of five chapters with one template: a move system acts on matrices, the blur forgets the moves, kept questions are found, and a normal form shows they are all. Here the moves are on both sides, $A \mapsto PAQ$.

Given

G1 (Building 0, Chapter II.5.) $K[x]$; division with remainder by any $d \neq 0$ when K is a field, and in $\mathbb{Z}$; det over any ring; P10 (equations are carried); P11 (inverse iff det is a unit); T2: $\mathbb{Z}$ and $K[x]$ pass the same lines.

G2 (Building 0, Chapter 18.) gcd, and "goes into" as a one-way blur.

G3 (Building 0, Chapter II.2.) Blurs of a ring by ideals; quotients R/I.

G4 (Chapter II.3.) Quotients of move systems; T1.

Definitions

D1 R: $\mathbb{Z}$, or $K[x]$ with K a field. Its units: ± 1, or the non-zero constants.

D2 An *R-module*: a bag M with $+$ (a comm move system) and scaling by members of R, with the laws of Building 0, Chapter 6, P2. R^n is one; so is every comm move system, with $R = \mathbb{Z}$ ($k \cdot m := m + \cdots + m$).

D3 (Build, then blur.) For an $m \times n$ matrix A over R: $\operatorname{coker} A := R^n$ blurred by gluing each row of A to 0, respecting $+$ and scaling. Every finite comm move system M is a $\operatorname{coker} A$: one generator x_g for each member g, one relation $x_g + x_h - x_{g+h}$ for each pair (Chapter 2, T1 and Chapter II.3, T1, read with $+$).

D4 $d_k(A)$: the gcd of all $k \times k$ minors of A (made monic, or ≥ 0); $d_0 := 1$.

The blur

On $m \times n$ matrices over R. O: none beyond the moves themselves.

B1 (From below.) Gluings: $A \sim PAQ$ for P, Q square with det a unit (G1, P11). Equivalently, by row and column moves: swap two rows; add r times one row to another; scale a row by a unit; and the same for columns.

B2 (From above.) Questions: $d_1(A), d_2(A), \ldots$.

Propositions

P1 (Survives: the d_k.) Each $k \times k$ minor of PA is an R-combination of $k \times k$ minors of A (an equation in the entries, carried to every R: G1, P10). So $d_k(A)$ goes into $d_k(PA)$, and back again by P^{-1}: d_k is kept. The same for AQ.

P2 (Survives: the module.) $\operatorname{coker}(PAQ)$ matches $\operatorname{coker} A$. (P only remixes the relations; Q renames the generators.)

P3 (Dies: the entries.) $\begin{pmatrix} 2 & 0 \\ 0 & 3 \end{pmatrix} \sim \begin{pmatrix} 1 & 0 \\ 0 & 6 \end{pmatrix}$ over $\mathbb{Z}$: $d_1 = 1$, $d_2 = 6$ for both.

P4 (Normal form; at least one, by Euclid.) Every A is glued to a diagonal

$$\operatorname{diag}(e_1, e_2, \dots, e_r, 0, \dots)$$

with $e_1 \mid e_2 \mid \cdots \mid e_r$, all $\neq 0$. (Move a smallest non-zero entry to the corner; divide with remainder along its row and column; if a remainder is left, it is smaller, and repeat; sizes or degrees cannot fall forever (Building 0, Chapter 1, G8). Then clear the row and column, and make the corner go into every other entry by adding a row.)

P5 (At most one.) $e_1 e_2 \cdots e_k = d_k(A)$ up to a unit, so $e_k = d_k/d_{k-1}$: fixed by the kept questions.

P6 (Appears.) B1 and B2 agree: two matrices are glued exactly when all their d_k agree. (P1 one way; P4 and P5 the other.)

P7 (Dies over $\mathbb{Z}[x]$.) The 1×2 matrix $(2 \;\; x)$ over $\mathbb{Z}[x]$ has $d_1 = 1$, but is glued to no $(e \;\; 0)$: the multiples of 2 and x together are not the multiples of one member (the constant terms of $2a + xb$ are even). Division with remainder was the line that P4 used, and $\mathbb{Z}[x]$ fails it.

Theorems

T1 (Every module given by finitely many generators and relations.) It matches

$$R^s \times R/(e_1) \times \cdots \times R/(e_r), \qquad e_1 \mid e_2 \mid \cdots \mid e_r,$$

for exactly one s and one such list of e_i that are not units (up to units). (P2, P4, P5.)

T2 (Chapter II.1, R2, proved.) Every finite comm move system matches a product of clocks $\mathbb{Z}/e_1 \times \cdots \times \mathbb{Z}/e_r$ with $e_1 \mid \cdots \mid e_r$, in one way only. ($R = \mathbb{Z}$, $s = 0$.)

T3 (Counting them.) The comm move systems with n members, up to matching, number $\prod_p \mathrm{p}(a_p)$, where $n = \prod p^{a_p}$ and $\mathrm{p}(a)$ is the number of partitions of a. For $n = 8$: 3 ($\mathbb{Z}/8$, $\mathbb{Z}/2 \times \mathbb{Z}/4$, $\mathbb{Z}/2 \times \mathbb{Z}/2 \times \mathbb{Z}/2$). For $n = 72$: 6. (Split each e_i by primes: Building 0, Chapter 3, T6.)

T4 (Two clocks, again.) $\mathbb{Z}/m \times \mathbb{Z}/n$ matches $\mathbb{Z}/mn$ exactly when $\gcd(m, n) = 1$: the Smith form of $\operatorname{diag}(m, n)$ is $\operatorname{diag}(\gcd, \operatorname{lcm})$.

Remarks

R1 (The template.) A move system acting on matrices; the blur "reachable by a move"; kept questions; a normal form proving they are all. Next in Building 1: equivalence over a field (rank), similarity (Jordan), forms (Sylvester), and the cycle type of Chapter 1 was the first case.

R2 (Where existence came from.) Once: the Euclid steps of P4, a construction. Uniqueness came from kept questions (P5).

R3 (Old names.) R-module; coker A: a presented module. B1: equivalence of matrices. d_k: determinantal divisors. P4: the Smith normal form; e_i: invariant factors. T1: the structure theorem for finitely generated modules over a principal ideal domain (here Euclidean). T2: the fundamental theorem of finite abelian groups. P7: $\mathbb{Z}[x]$ is not a principal ideal domain.

R4 (Machine check, `checks/b1-p2-ch4-check.py`.) P1 (d_k kept under 200 random moves); P4–P6 against a Smith form routine on random integer matrices; P3; T3 for $n \leq 200$ by listing diagonal forms; T4.

Exercises

E1 Find the Smith form of $\begin{pmatrix} 2 & 4 & 4 \\ -6 & 6 & 12 \\ 10 & -4 & -16 \end{pmatrix}$ over $\mathbb{Z}$, and say which comm move system it presents.

E2 List the comm move systems with 36 members.

E3 Over $\mathbb{Q}[x]$, find the Smith form of $\begin{pmatrix} x & 1 \\ 0 & x \end{pmatrix}$.

E4 Show $\mathbb{Z}/4 \ncong \mathbb{Z}/2 \times \mathbb{Z}/2$ by P5, and by cyc.

Part III. A cubic and its moves

Building 0, Chapters 35–37 drew a cubic in the plane, found the functions on it, and met the same curve over $\mathbb{C}$ as a torus. This part glues its points by lines and finds a comm move system, then reads it over $\mathbb{C}$ and over $\mathbb{Z}/p$.

Chapter III.1. Points glued by lines

sums of points → points · from: Chapters 2, II.1, II.3; Building 0, Chapters 35, 36 · build, blur

Add points as letters, and glue the three points of each line to three copies of O. The bags form a move system for free, and each bag of size 0 holds exactly one point.

The usual road defines a sum of two points by a drawing and then spends pages proving it is associative. Here the sum is a blur of a free move system, so associativity is never in doubt; the work moves to "at most one point per bag".

Given

G1 (Building 0, Chapter 36.) k with $2, 3 \neq 0$; the cubic C: $y^2 = x^3 + ax + b$ with $\Delta \neq 0$, and O. P1 (the edge meets C only at O, three times); P2 (a line meets C in at most 3, counted); P3 (the mirror); T1 (Build: two points of $C(k)$ on a line give a third, and the counts add to exactly 3); P10 (no function has weight 1); P11.

G2 (Chapter 2, T1; Chapter II.1.) The free comm move system on a bag: finite sums $\sum n_P\, P$, n_P in $\mathbb{Z}$.

G3 (Chapter II.3.) The bags of a blur of a move system that respects both sides form a move system; for a comm one, respecting + is enough.

Definitions

D1 (Build.) $\mathbb{Z}[C]$: the free comm move system on the points of $C(k)$ (G2).
$\deg \sum n_P P := \sum n_P$.

D2 For a line L, $L \cdot C :=$ its meeting points with their counts, as a member of $\mathbb{Z}[C]$: $P + Q + R$, or $2P + R$ for a tangent, or $3O$ for the edge.

The blur

On $\mathbb{Z}[C]$. O: $+$.

B1 (From below.) Gluing: $L \cdot C \sim 3\, O$ for every line L that meets C at points of $C(k)$ (G1, T1: all three are then in $C(k)$). Respects $+$. Write $\mathrm{Pic}(C)$ for the bags.

B2 (From above.) Questions: the degree; and, for each D of degree 0, "is D glued to 0?", read through functions: $D \sim 0$ exactly when D is the list of zeros minus poles of a product of powers of lines (P2).

Propositions

P1 (Free: a move system.) $\mathrm{Pic}(C)$ is a comm move system (G3). Its law is associative because $+$ on $\mathbb{Z}[C]$ is. Nothing was drawn.

P2 (What a gluing is.) The affine line $\ell = \alpha x + \beta y + \gamma$ is a function on C (Building 0, Chapter 36) whose zeros, counted, are the affine points of $L \cdot C$, and whose only pole is at O, of order $w(\ell) = 3$ or 2 (P11 there). So $L \cdot C - 3O$ is "zeros minus poles" of ℓ, and $D \sim 0$ exactly when D is zeros minus poles of $\prod \ell_i^{n_i}$.

P3 (Survives: the degree.) Each gluing has degree $3 - 3 = 0$. So deg is kept, and $\mathrm{Pic}^0(C)$, the bags of degree 0, is a sub move system.

P4 (Two moves on points.) For P, Q in $C(k)$ let $P * Q$ be the third point on their line (G1, T1). Then

$$P + Q + (P * Q) \sim 3O, \qquad R + \bar{R} \sim 2O,$$

the second by the upright line through R and its mirror $\bar{R}$ (G1, P3). So $P + Q \sim O + \overline{P * Q}$: two points are glued to one point plus O.

Theorems

T1 (At least one: normal form.) Every D of degree 0 is glued to $P - O$ for some P in $C(k)$. (Write $D = \sum(P_i - O) - \sum(Q_j - O)$. $-(Q - O) \sim \bar{Q} - O$ by P4. Then $(P-O)+(Q-O) \sim \overline{P * Q} - O$ by P4, and repeat: each step is a Build of Building 0, Chapter 36, T1.)

T2 (At most one.) $P - O \sim Q - O$ only when $P = Q$. (Put $R := \overline{P * \bar{Q}}$; then $P - Q \sim R - O$, so $R - O$ is zeros minus poles of some $f = \prod \ell_i^{n_i}$ (P2). If $R \neq O$, f has no pole on the affine part, so f lies in $k[C]$ (classical, proof deferred: a function on C with no affine pole is a polynomial function); its only pole is a simple one at O, so $w(f) = 1$; Building 0, Chapter 36, P10 says no such f exists. So $R = O$, that is $P = Q$.)

T3 (Appears: the points are a move system.) $P \mapsto$ the bag of $P - O$ matches $C(k)$ one-to-one with $\mathrm{Pic}^0(C)$ (T1, T2). Carried back to the points, the law is

$$P \oplus Q = \overline{P * Q}, \qquad O \text{ does nothing}, \qquad \ominus P = \bar{P},$$

and $P \oplus Q \oplus R = O$ exactly when P, Q, R lie on one line. Associativity of $\oplus$ is P1, carried.

T4 (Formulas.) For $P = (x_1, y_1)$, $Q = (x_2, y_2)$ with $x_1 \neq x_2$: $P \oplus Q = (x_3, -y_3)$ with x_3, y_3 as in Building 0, Chapter 36, T1. $2P$: the same with the tangent's slope $(3x_1^2 + a)/(2y_1)$; $2P = O$ when $y_1 = 0$.

Propositions on examples

P5 (Over $\mathbb{Q}$, a clock.) $y^2 = x^3 + 1$: the points O, $(-1, 0)$, $(0, \pm 1)$, $(2, \pm 3)$, and $(2, 3)$ has order 6: $C(\mathbb{Q})$ holds a $\mathbb{Z}/6$.

P6 (Over $\mathbb{Q}$, no clock.) $y^2 = x^3 - 2$, $P = (3, 5)$: $2P = (\frac{129}{100}, -\frac{383}{1000})$. The denominators grow with each doubling, and P has no finite order.

P7 (Dies when $\Delta = 0$.) On $y^2 = x^3$ T2 fails at the bad point, which every construction through it returns. Leave that point out and the rest is again a move system, matching $(k, +)$ by $(x, y) \mapsto x/y$ (machine-checked over $\mathbb{Z}/p$; classical).

Remarks

R1 (What the method did.) Associativity, the hard line of the usual road, is free (P1): the blur is of a free move system. Existence of a normal form is Euclid-like steps (T1), each a Build. The one "at most one" line, T2, is Building 0, Chapter 36, P10, plus one classical step, deferred. A second road to T2 over $\mathbb{C}$ is Chapter III.2.

R2 (Kept questions.) The two questions of B2 are all: degree, and "zeros minus poles of a function". The same template as Chapter II.4: free thing, gluing, kept questions, normal form.

R3 (Old names.) $\mathbb{Z}[C]$: divisors; B1: linear equivalence (lines generate it on a cubic); $\mathrm{Pic}^0(C)$: the degree-zero Picard group; T3: the group law on an elliptic curve; T2: "no function with a single simple pole", a case of Riemann–Roch; P4: the chord-and-tangent construction; P6: a point of infinite order; T3 with $k = \mathbb{Q}$ and finitely many generators: Mordell–Weil (classical, reported, not used).

R4 (Machine check, `checks/b1-p3-check.py`.) Over $\mathbb{Z}/p$, $p = 5, 7, 11, 13$, every smooth (a, b): O does nothing, mirror undoes, $\oplus$ commutes, associativity on all triples (on 12 points per curve when there are more); P5; P6.

Exercises

E1 On $y^2 = x^3 + 1$ over $\mathbb{Q}$, compute $2(2, 3)$ and $3(2, 3)$.

E2 Show that $P \oplus P = O$ exactly when $P = O$ or $y(P) = 0$. How many such P are there over $\mathbb{C}$?

E3 Why does B1 include the edge line, and what does it glue?

E4 Over $\mathbb{Z}/5$, list $C(\mathbb{Z}/5)$ for $y^2 = x^3 + 1$ and find which clock it is.

Chapter III.2. The torus carries the law

$\mathbb{C}/\Lambda \to C(\mathbb{C})$ · from: Chapter III.1; Building 0, Chapters 8, 37 · arrow

Three points of the torus add to 0 exactly when their images lie on one line. So the arrow of Building 0, Chapter 37 respects the moves, and + on the torus is the chord law.

Given

G1 (Building 0, Chapter 37.) Λ, $\mathbb{C}/\Lambda$ with +; $\wp$; T4 (zeros and poles equal in number; their sums differ by a member of Λ); T8 (the arrow $\phi : z \mapsto (\wp(z) : \wp'(z) : 1)$ matches $\mathbb{C}/\Lambda$ with $C(\mathbb{C})$, C: $y^2 = 4x^3 - g_2x - g_3$).

G2 (Chapter III.1.) $\oplus$; T3: $P \oplus Q \oplus R = O$ exactly when P, Q, R lie on one line. (The 4 and the g_i change nothing: scale y by 2.)

Theorems

T1 (Lines are sums in Λ.) $\phi(z_1), \phi(z_2), \phi(z_3)$ lie on one line, with counts, exactly when $z_1 + z_2 + z_3 \in \Lambda$. (A line $\alpha x + \beta y + \gamma = 0$ with $\beta \neq 0$ gives the elliptic map $\alpha\wp + \beta\wp' + \gamma$, with one triple pole per bag, at 0. Its zeros are the z whose images lie on the line (G1, T8): three of them, and their sum is in Λ (G1, T4). For $\beta = 0$: $\wp - c$ has zeros z, $-z$, and the third meeting point is $O = \phi(0)$; again the sum is 0. Conversely, the line through $\phi(z_1)$ and $\phi(z_2)$ meets C a third time at some $\phi(z_3')$, and $z_3' \sim -z_1 - z_2$.)

T2 (Survives: the arrow respects the moves.) $\phi(z_1 + z_2) = \phi(z_1) \oplus \phi(z_2)$. (T1 and G2: both sides are the point R with $\phi(z_1) \oplus \phi(z_2) \oplus \ominus R = O$, and $\ominus\phi(z) = \phi(-z)$, the mirror, since $\wp$ is even and $\wp'$ odd.)

T3 (A second road to Chapter III.1, T2, over $\mathbb{C}$.) For a cubic reached by T8, $P \mapsto P - O$ is one-to-one: the torus side has no gluing to undo. And associativity of $\oplus$ is the associativity of + on $\mathbb{C}$, carried.

T4 (Points of finite order.) The P with $nP = O$ in $C(\mathbb{C})$ match $(\frac{1}{n}\Lambda)/\Lambda$, so there are n^2 of them and they form $\mathbb{Z}/n \times \mathbb{Z}/n$. For $n = 2$: O and the three points with $y = 0$. For $n = 3$: the nine points where the tangent meets C three times.

Remarks

R1 (What the method did.) One arrow. Its respecting + is one line of Building 0, Chapter 37 (T4, the sum of zeros) read through Chapter III.1, T3.

R2 (Reported.) Every smooth cubic over $\mathbb{C}$ is reached by some Λ (classical, uniformisation). With it, T3 covers every cubic over a k inside $\mathbb{C}$, and the deferred step of Chapter III.1, T2 is no longer needed there. Not used.

R3 (Old names.) T1: Abel's theorem on the torus; T2: the addition theorem of $\wp$; T4: $E[n] \cong (\mathbb{Z}/n)^2$; the $n = 3$ points: flexes.

R4 (Machine check, `checks/b1-p3-check.py`.) With $\omega_1 = 1$, $\omega_2 = 0.3 + 1.1i$: for random z_1, z_2 and $z_3 = -z_1 - z_2 + \omega_1 + \omega_2$ the three images have det below 10^{-6} (relative); moving z_3 by 0.13 gives det above $3 \cdot 10^{-3}$.

Exercises

E1 Use T2 to write $\wp(2z)$ in terms of $\wp(z)$ and $\wp'(z)$.

E2 Why are the points with $nP = O$ over $\mathbb{R}$ fewer than n^2?

E3 Show from T1 that $\phi(z)$ is a point where the tangent meets C three times exactly when $3z \in \Lambda$.

Chapter III.3. Counting points over a clock

cubics over $\mathbb{Z}/p \to$ their sizes · from: Chapters II.2, II.4, III.1; Building 0, Chapters 3, 17 · count

Over $\mathbb{Z}/p$ the move system of Chapter III.1 is finite. Its size is $p+1$ plus a sum of p signs, and the signs nearly cancel.

Given

G1 (Chapter III.1.) $C(\mathbb{Z}/p)$ with $\oplus$, for $p \neq 2, 3$ and $\Delta \neq 0$ in $\mathbb{Z}/p$.

G2 (Chapter II.2, T1, T2.) Lagrange; the order of a move divides the size.

G3 (Chapter II.4, T2.) A finite comm move system is $\mathbb{Z}/e_1 \times \cdots \times \mathbb{Z}/e_r$, $e_1 \mid \cdots \mid e_r$, in one way.

G4 (Building 0, Chapter 3, T4, T5.) $\mathbb{Z}/p$ is a field; $c^p = c$.

Definitions

D1 (The square sign.) For c in $\mathbb{Z}/p$: $\chi(c) := 0$ if $c = 0$; 1 if $c = y^2$ for some y; -1 otherwise. $N_p := |C(\mathbb{Z}/p)|$, O included.

Propositions

P1 (How many square roots.) $y^2 = c$ has $1 + \chi(c)$ answers in $\mathbb{Z}/p$. (y and $-y$ differ unless $y = 0$, as $2 \neq 0$; at most two by Building 0, Chapter 7, T4 over $\mathbb{Z}/p$.)

P2 (The count.)

$$N_p \;=\; p + 1 + \sum_{x \in \mathbb{Z}/p} \chi\big(x^3 + ax + b\big).$$

(P1 for each x, and O.)

P3 (Euler's test.) $\chi(c) = c^{(p-1)/2}$ in $\mathbb{Z}/p$, read as 0, 1, -1. (The squares $c \neq 0$ are $(p-1)/2$ in number (P1) and each has $c^{(p-1)/2} = y^{p-1} = 1$ (G4); $x^{(p-1)/2} - 1$ has no more roots than that.)

Theorems

T1 (Lagrange on the curve.) For every P in $C(\mathbb{Z}/p)$, $N_p\, P = O$. (G2.)

T2 (At most two clocks.) $C(\mathbb{Z}/p)$ matches $\mathbb{Z}/e_1 \times \mathbb{Z}/e_2$ with $e_1 \mid e_2$ (G3; at most two factors because the P with $\ell P = O$, ℓ prime, number at most ℓ^2: classical for $\mathbb{Z}/p$; over $\mathbb{C}$ it is Chapter III.2, T4). Also $e_1 \mid p - 1$ (classical, reported).

T3 (The signs nearly cancel. Reported; its road is Building 2.)

$$|N_p - p - 1| \;\leq\; 2\sqrt{p}.$$

The sum in P2 has p terms ± 1 and 0; if they were independent coin tosses its size would be about $\sqrt{p}$ (Building 0, Chapter 17's scale). T3 says the curve's signs never do worse than twice that.

The road: the counts N_{p^m} over the floors with p^m members form a ladder; one rough bound $|N_{p^m} - p^m - 1| \le c\, p^{m/2}$, the same c on every rung, is amplified along the ladder to the constant 2 (Amplify, Building 2). Two inputs of that road are classical and reported: the shape $N_{p^m} = p^m + 1 - \alpha^m - \beta^m$ with $\alpha\beta = p$, and the rough bound.

Remarks

R1 (What the method did.) Counting, and two lines from Part II. The real content, T3, is a cancellation line: the edge the survey of the book keeps meeting (Building 0, Chapter 17, P6).

R2 (Old names.) χ: the Legendre symbol; P3: Euler's criterion; T3: Hasse's theorem (the Riemann hypothesis for elliptic curves over finite fields); α, β: Frobenius eigenvalues; $p + 1 - N_p$: the trace of Frobenius.

R3 (Machine check, `checks/b1-p3-check.py`.) For every prime $5 \le p < 60$ and every smooth (a, b): P2 against direct listing; T1 (each point's order divides N_p); T2 ($e_1 e_2 = N_p$, $e_1 \mid e_2$, $e_1 \mid p - 1$); T3, with the worst case $|N_p - p - 1|/2\sqrt{p} = 0.991$.

Exercises

E1 Count $C(\mathbb{Z}/7)$ for $y^2 = x^3 + 2$ by P2, and check T3.

E2 Show that for $y^2 = x^3 + x$ and $p \equiv 3 \pmod 4$, $N_p = p + 1$. (Pair x with $-x$, and use $\chi(-1) = -1$.)

E3 Use T1 to find the order of $(0, 1)$ on $y^2 = x^3 + 1$ over $\mathbb{Z}/5$ without adding.