

subject	date	keywords
Foundations of Computing II	26 Feb 2024	Recursion Nonhomogenous Relations
topic	Nonhomogenous Recurrence Relations	

Announcements

Exam results released Wednesday

Read the textbook

Background

Define a recurrence relation  $\{a_n\}$  by

$$c_0 a_n + c_1 a_{n+1} + \dots + c_k a_{n+k} = f(n)$$

for constants  $c_j \in \mathbb{R}$ , some function  $f(n)$ 

(Book page 479)

Homogenous

When  $f(n)$  is identically (always) zero,  $\{a_n\}$  is homogenous

Nonhomogenous

For anything else,  $\{a_n\}$  non-homogenousOnly for some  $f(n)$  do "nice" closed forms exist.

Types of Solvable N-H Relations

We'll focus on (for  $c, r$  non-zero):

- $f(n) = c$  constant
- $f(n) = cn$  linear
- $f(n) = cn^2$  quadratic
- $f(n) = r^n$  geometric
- $f(n) = \text{combination of above}$  combination

We will not cover  
trigonometric equations

Steps

- 1) Write in the associated homogenous form,  $c_0 a_n + c_1 a_{n+1} + \dots + c_k a_{n+k} = 0$  !!!
- 2) Get form of the particular solution,  $a_n^{(p)}$ , based on  $f(n)$  (see pg 479)
- 3) Using non-homogenous relation, solve for  $a_n^{(p)}$  w/ initial conditions
- 4)  $a_n = a_n^{(h)} + a_n^{(p)}$

First-Order N.H.

Example

$$a_{n+1} + ca_n = f(n)$$

$$a_{n+1} = 2a_n + n \iff a_{n+1} - 2a_n = n$$

w/  $a_0 = 1$

→ Characteristic equation:  $r - 2 = 0 \iff r = 2$ 

$$\rightarrow a_n^{(h)} = c(2)^n$$

$$\rightarrow a_n^{(p)} = A_1 n + A_0 \text{ (from textbook)}$$

substitute into N.H. relation

$$\rightarrow a_{n+1}^{(p)} = 2a_n^{(p)} + n = 2(A_1 n + A_0) + n$$

$$= 2A_1 n + 2A_0 + n$$

I don't know what's going on ---

$$\leadsto a_n^{(p)} = -n - 1 \rightarrow a_n = a_n^{(h)} + a_n^{(p)} = c(2)^n - n - 1$$

$$1 = a_0 = c(2)^0 - 0 - 1 \Rightarrow c = 2$$

$$\rightarrow a_n = 2(2)^n - n - 1 = 2^{n+1} - n - 1$$

subject

date

keywords

topic

Example 2

$$a_{n+1} = a_n + n, \quad a_0 = 1$$

$$\rightarrow a_{n+1} - a_n = n \quad \leadsto \quad r-1=0 \rightarrow r=1$$

$$\rightarrow a_n^{(h)} = c \cdot 1^n = c$$

$$\rightarrow a_n^{(p)} = \cancel{A_1 n + A_0} \quad A_1 n^2 + A_0 n \quad (\text{exception to table})$$

$$\text{Solve } a_{n+1}^{(p)} = A_1(n+1)^2 + A_0(n+1) \quad \text{from def'n of relation}$$

$$= A_1(n^2 + 2n + 1) + A_0(n+1) = A_1 n^2 + A_0 n + n$$

$$\Rightarrow \begin{cases} A_1 = A_1 \\ 2A_1 + A_0 = A_0 + 1 \\ A_1 = A_0 \end{cases} \quad \text{match coefficients}$$


$$\Rightarrow \begin{cases} A_1 = \frac{1}{2} \\ A_0 = -\frac{1}{2} \end{cases}$$

$$\Rightarrow a_n = a_n^{(h)} + a_n^{(p)} = c + \frac{1}{2}n^2 + \frac{1}{2}n$$

$$\text{Let } a_0 = 1, \text{ then } 1 = c + \frac{1}{2}(0)^2 + \frac{1}{2}(0)$$

$$\Rightarrow a_n = \frac{1}{2}n^2 + \frac{1}{2}n + 1$$

See next page for better notes



subject

Foundations of Computing II

date

26 Feb 2024

keywords

Recurrence Relation  
Non homogeneous  
Linear  
General Solution

topic

10.3 Nonhomogeneous Linear Recurrence Relations

Notes outside of Class

 $a_n^{(p)}$  table

$f(n)$	$a_n^{(p)}$
c	$A$
$n$	$A_1 n + A_0$
$n^2$	$A_2 n^2 + A_1 n + A_0$
$n^t$	$A_t n^t + A_{t-1} n^{t-1} + \dots + A_1 n + A_0$
$r^n$	$A r^n$

$$\left. \begin{array}{l} a_n^{(h)} : \text{"homogenous" solution} \\ a_n^{(p)} : \text{"particular" solution} \end{array} \right\} a_n = a_n^{(h)} + a_n^{(p)} \quad \text{"general" solution}$$

Capital A's are unknown,  
not the value of  $a_i$

To Solve N-H:

Suppose we're given a linear recurrence relation...

- 1) Write it in standard form, i.e.  $c_0 a_n + c_1 a_{n-1} + \dots + c_k a_{n-k} = f(n)$   
 $\swarrow$  "set equal to"  $\searrow$   $a_n^{(h)}$
- 2) Solve  $a_n^{(h)} = 0$  using old techniques (e.g. characteristic eqn)
- 3) Use the above table to find the form of  $a_n^{(p)}$  (note exceptions)
- 4) Using the definition of the recurrence relation, solve for unknowns in  $a_n^{(p)}$  (match coefficients)
- 5) Solve  $a_n = a_n^{(h)} + a_n^{(p)}$  using boundary/initial conditions

Examples from Class

$$a_{n+1} = 2a_n + n, a_0 = 1 \quad *$$

- 1)  $a_{n+1} - 2a_n = n$
  - 2)  $a_n^{(h)} = a_{n+1} - 2a_n = 0 \Rightarrow r - 2 = 0 \Rightarrow r = 2$   
 $\Rightarrow a_n^{(h)} = A \cdot 2^n$
  - 3)  $f(n) = n \Rightarrow a_n^{(p)} = A_1 n + A_0$
  - 4)  $a_{n+1}^{(p)} = A_1(n+1) + A_0 = A_1 n + A_1 + A_0$  (from 3)  
 $a_{n+1}^{(p)} = 2a_n^{(p)} + n$  (from \*)  
 $= 2(A_1 n + A_0) + n = 2A_1 n + 2A_0 + n$
- $*$   $A_1 n + A_1 + A_0 = 2A_1 n + 2A_0 + n$   
 $\Rightarrow A_1 = 2A_1 + 1$  (everything w/  $n$ )  
 $\hookrightarrow A_1 = -1$   
 $\Rightarrow A_1 + A_0 = 2A_0$  (everything else)  
 $\hookrightarrow A_0 = -1$   
 $\Rightarrow a_n^{(p)} = A_1 n + A_0 = -n - 1$
- 5)  $a_n = a_n^{(h)} + a_n^{(p)} = C \cdot 2^n - n - 1$ , yet  $a_0 = 1$   
 $\Rightarrow 1 = C \cdot 2^0 - 0 - 1 \Rightarrow C = 2 \Rightarrow a_n = 2 \cdot 2^n - n - 1$

subject

date

keywords

topic

Example 2

$$a_{n+1} = a_n + n, a_0 = 1$$

$$1) a_{n+1} - a_n = n \quad (\text{standard form})$$

$$2) a_n^{(h)} = a_{n+1} - a_n = 0 \Rightarrow r - 1 = 0 \Rightarrow r = 1 \leadsto a_n^{(h)} = c \cdot (1)^n = c$$

$$3) f(n) = n \Rightarrow a_n^{(p)} = A_1 n + A_0$$

$$4) a_{n+1}^{(p)} = A_1(n+1) + A_0 = A_1 n + A_1 + A_0$$

$$a_{n+1}^{(p)} = a_n^{(p)} + n = A_1 n + A_0 + n$$

$$\Rightarrow A_1 = A_1 + 1 \quad ! \quad \text{no solution}$$

multiply by  $n$ , try again (this will happen from time to time...)

$$3) a_n^{(p)} = A_1 n^2 + A_0 n$$

$$4) a_{n+1}^{(p)} = A_1(n+1)^2 + A_0(n+1) = A_1(n^2 + 2n + 1) + A_0(n+1)$$

(replace  $n$  w/  $n+1$ )

$$a_{n+1}^{(p)} = a_n^{(p)} + n = A_1 n^2 + A_0 n + n$$

(from def'n of relation)

$$\Rightarrow \begin{cases} A_1 = A_1 & (n^2 \text{ coeff}) \\ 2A_1 + A_0 = A_0 + 1 & (n \text{ coeff}) \\ A_1 + A_0 = 0 & (\text{const coeff}) \end{cases}$$

$$\downarrow \quad A_0 = -A_1 \quad \rightarrow 2A_1 + (-A_1) = (-A_1) + 1 \Leftrightarrow A_1 = -A_1 + 1 \Rightarrow A_1 = \frac{1}{2} \Rightarrow A_0 = -\frac{1}{2} \quad \checkmark$$

$$\Rightarrow a_n^{(p)} = \frac{1}{2}n^2 - \frac{1}{2}n$$

$$5) a_n = a_n^{(h)} + a_n^{(p)} = c + \frac{1}{2}n^2 - \frac{1}{2}n, a_0 = 1$$

$$\Rightarrow \frac{1}{2}(0)^2 - \frac{1}{2}(0) + c = 1 \Rightarrow c = 1$$

$$\Rightarrow a_n = \frac{1}{2}n^2 - \frac{1}{2}n + 1$$



not sure if it's  $+\frac{1}{2}n$  or  $-\frac{1}{2}n$ ... may have missed a minus sign !!

to be continued...