

Project 1: Root Finding via Lagrange Polynomial Interpolation

Max Ortman Teammate 1 Teammate 2

February 15, 2026

The Algorithm

Each step we create a Lagrange Polynomial using four sampled points. We then evaluate the roots of that polynomial and add whatever point is closest to zero as a new sampled point. We then remove the point with the largest x value while making sure there is at least one point on each side of zero. This process is repeated until we have reached the desired number of iterations or the error is sufficiently small. Our method assumes that a range that has points on either side of zero is given, and the function only crosses zero once in that range.

The code is too long to fit in the doc so please check out the attached `main.py` file for the implementation as we handle edge cases. But the sudo code is as follows:

```

def rootfind(f_:Callable, min:float, max:float) -> float:
    # get four points
    points = [min, min+(max-min)/3, min+2*(max-min)/3, max]

    cubic_count = 0
    bisect_count = 0

    for i in range(max_iter_):
        a, b, c, d = make_lagrange_cubic(points)
        roots = cubic_roots(a, b, c, d)

        # Filter the roots to meet the following:
        # Root is within the range of points
        # Root of polynomial is within a tolerance of being a root
        # Root is not too close to the last best root (to prevent loops)
        candidate = get_candidate(roots, points)
        if candidate is not None:
            best = candidate
        else:
            # If no candidate roots are found we bisect the closest
            # positive and negative points to find a new candidate root
            best = (min_pos[0] + min_neg[0]) / 2

        # Sample new point and add to list
        best_y = f_(best)

        # Return if within tolerance
        if abs(best_y) < tol_:
            return best

        # Add point
        points += (best, best_y)

        # Remove the nearest point on both sides of the root

        # Get xs and ys without the closest pos and neg to keep bounds
        filtered_points = [points not in (min_pos, min_neg)]
        furthest = furthest_point(filtered_points, best)
        points.remove(furthest)

```

Explanation

You can use an interactive version in desmos with 3 points: <https://www.desmos.com/calculator/atqa25yh0u> to see how the method works. Blue is first step and red is the second.

Here is also our method taking on a particularly challenging function. You can see it samples really close then slowly moves the leftmost point inwards. This is a larger issue that leads to some cases not performing as well as we would like. But when it doesn't do this it can one shot some examples




Figure 1: Visualization of the method on a challenging function. It samples closely (blue) and moves the leftmost point inwards (red). Note the performance in later iterations.

To estimate the zero for this function, we use sampled points to construct a Lagrange Polynomial. The basis polynomials are defined as:

$$L_i(x) = \prod_{j \neq i} \frac{(x - x_j)}{(x_i - x_j)}$$

Using these basis functions, the unique polynomial of degree n that passes through all $n + 1$ points is:

$$P(x) = \sum_i y_i L_i(x)$$

To start our method we sample the beginning, end, and middle 2 points. This gives us a set of 4 points to construct our Lagrange Polynomial. We then evaluate $P(x) = 0$ to get our initial estimate of the root. We then add that as a middle point and remove the point with the largest x value while making sure there is at least one point on each side of zero. This process is repeated until we have reached the desired number of iterations or the error is sufficiently small.

Analysis & Results

Table 1: Rootfinding comparison averaged over 10 trials with randomized bounds

Function	Iterations		Function Calls		Step Type	
	Cubic	Newton	Cubic	Newton	Cubic	Bisect
$\sin(x) + 0.2$	3.0	3.0	7.0	10.0	3.0	0.0
$x^3 - x - 2$	1.0	3.7	5.0	12.1	1.0	0.0
$\cos(x) - x$	3.7	4.0	7.7	13.0	3.5	0.2
$e^x - 3$	4.0	3.8	8.0	12.4	4.0	0.0
$x^5 - x - 1$	33.9	5.8	36.7	18.4	3.5	30.4
$\ln(x) - 1$	5.0	4.0	9.0	13.0	5.0	0.0
$\tan(x) - 2x$	100.0	8.0	74.0	25.2	35.0	35.0
$x^2 - 2$	1.0	3.6	5.0	11.8	1.0	0.0
Average	18.9	4.5	19.1	14.5	7.0	8.2