

Ease: An R Package for Efficient Population Genetics Simulations

Ehouarn Le Faou ^{1*}

¹ Department of Ecology and Evolution, University of Lausanne, 1015 Lausanne, Switzerland * These authors contributed equally.

DOI: [10.xxxxxx/draft](https://doi.org/10.xxxxxx/draft)

Software

- [Review](#) 
- [Repository](#) 
- [Archive](#) 

Editor: [Open Journals](#) 

Reviewers:

- [@openjournals](#)

Submitted: 01 January 1970

Published: unpublished

License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License ([CC BY 4.0](#)).

Summary

Ease is an R package for efficient and reproducible population genetics simulations, designed to facilitate the implementation of theoretical models involving few to moderate numbers of loci with arbitrary epistasis and dominance patterns. The package implements efficient simulation of such models by tracking genotype frequencies using deterministic transition matrices for mutation, reproduction, and selection. It incorporates genetic drift via multinomial sampling. Ease provides a flexible simulation environment well suited to theory-driven research in population genetics.

Statement of need

Population genetics plays a crucial role in our understanding of the adaptation of species. Its methods are used, for example, to trace back speciation times, to understand the genetic diversity of species and the consequences of selection. The field has always been largely driven by theory, which, thanks to simplified models of reality, has made it possible to understand and predict how evolution shapes biodiversity. Along with mathematical treatment of those models, simulations are essential, particularly for exploring analytically intractable models or for assessing their behaviour when their assumptions are relaxed. A wide range of simulation frameworks are available, e.g. coalescent simulators optimised for neutral genome-scale models (e.g. *msprime*; Baumdicker et al. (2022)), or forward-in-time individual-based simulators designed for biologically realistic representations of selection, demography and drift (e.g. *SLiM*; Haller & Messer (2023)). These tools are well suited to many applications, but they are not optimised for models in which the genetic architecture is deliberately simple while selection depends on complex epistasis and dominance relations between finite set of alleles in arbitrarily large populations. Many canonical models (see, for example, models described in Crow & Kimura (1970); Bürger (2000); Ewens (2004)) fall into this category, which have not been the focus of efforts to produce an effective, flexible and easy-to-use simulator.

The R package Ease was developed to fill this gap. It is a lightweight and reproducible implementation of population genetics models that include only a few loci. Ease represents populations by genotype frequency vectors rather than by individual genomes, and updates these frequencies across generations under selection, mutation, recombination, migration, and genetic drift. This approach is computationally efficient when the number of loci and alleles is limited, and it closely matches the structure of many population genetic models that are formulated as recursions on genotype or haplotype frequencies.

38 Software design

39 Ease incorporate both nuclear and cytoplasmic (homoplasmic) loci. The loci of the first type
40 are diploid, with Mendelian transmission, while those of the second type are haploid with a
41 strict maternal transmission. Any number of alleles can be defined for each locus. Unlike
42 individual-based simulations, where each individual is explicitly defined (i.e. each with its own
43 combination of alleles), Ease tracks the frequencies of allele combinations themselves. Allele
44 combinations are defined as unique sets of alleles, and exist in two types: *haplotypes* and
45 *genotypes*. A *haplotype* is defined as the allele combination of a gamete, which is composed of
46 one *nuclear haplotype* and one *cytoplasmic haplotype*. Two *nuclear haplotypes* can be paired
47 together to form a *nuclear genotype* and combined with a *cytoplasmic haplotype* to form a
48 *genotype*. A *genotype* is therefore the allele combination of a zygote.

49 Once the user has defined how many loci (nuclear and cytoplasmic) and how many alleles per
50 locus they want to simulate, Ease defines automatically a set of transition matrices which are
51 used at each generation and constant throughout the simulation (Figure [Figure 1](#)). These
52 matrices allow, for example, the haplotype frequency vector of gametes to be calculated from
53 the genotype frequencies of the parents by a simple matrix multiplication that incorporates
54 recombination, random segregation of chromosomes, and mutation. Theoretically, no hard
55 limit is imposed onto the number of loci and the number of alleles per locus the user can
56 define, but the computational method Ease uses to track allele frequency changes imposes a
57 practical limit to them. This is because the size of the transition matrices becomes exceedingly
58 large when the number of haplotypes and the number of genotypes is large, and so when the
59 number of loci exceeds a handful. One advantage of this approach is that populations can
60 be of any size (up to millions or billions of individuals) without slowing down the simulation.
61 This is because individuals are not simulated explicitly. Genetic drift is instead simulated by
62 multinomial sampling, whose performance is only marginally affected by sample size.

63 The primary use of Ease is to track the trajectories of genotypes, haplotypes and allele
64 frequencies over generations. It does so through a canonical life cycle used in population
65 genetics, in which diploid zygotes are first produced by the fusion of one male and one female
66 haploid gametes resulting from independent meiosis (Figure [Figure 1](#)). These zygotes then
67 develop into adults, which then produce the gametes that will form the zygotes of the next
68 generation before dying (generations do not overlap). Along this life cycle, selection can
69 happen at three time points. Adult genotypes can be associated with absolute fecundity values
70 for female and male gamete production. Gametes, depending on their haplotype and their sex,
71 can be associated with survival probabilities. Finally, zygote genotypes can be associated with
72 different survival probabilities, which occur before they become adults.




Figure 1: Summaries of the various steps implemented in Ease to run simulations with a number G of genotypes and H of haplotypes. All the transition matrices (reproduction, selection, migration, ...) are available for consultation using the print method, and can be defined in a variety of ways, depending on user's requirements.

Ease assumes two types of gamete associated with two mating types, which are called male and female gametes for simplicity. For the sexual identity of adult individuals, there are two possible scenarios. In the default scenario, all individuals are hermaphrodites and therefore capable of producing both female and male gametes. A self-fertilisation rate can be define, as the probability for a zygote to be produced through the fusion of gametes produced by the same individual. In an alternative scenario, zygotes are assigned to develop into adults producing only male gametes or only female gametes, with a 50% chance for each. It is possible to simulate this alternative scenario from the default one by defining explicitly a sex-determining locus, which can be useful when simulating sex-linked loci or variable sex ratios. This can also help to simulate more complex sexual systems, such as gynodioecy or androdioecy, among others.

Finally, the package supports the simulation of multiple populations with any migration pattern between them, e.g. island models, stepping stone models, or models with different population sizes.

Example of use

Model implementation is standardised and involves defining the genome, allelic effects, population(s) and their interconnection within a metapopulation, before finally simulating and extracting the results of the simulation(s) (Figure 2). A model can be simulated multiple times (without parallelisation), their results are then combined in the result objects.



Figure 2: Steps for implementing a model with Ease and associated functions, from defining the genome and the selective effect of alleles to extracting the results of the model simulation.

In the example below, each individual is represented by a genotype at two diallelic loci. One of these is nuclear with wildtype allele "A" and alternative allele "a", and the other is cytoplasmic with wildtype allele "B" and alternative allele "b". By default, when defining a population, the first allele named (here the wildtypes "A" and "B", see below) is the one that is fixed in the population at the start of the simulation.

```
library(Ease)
# Definition of a genome with 2 locus, 2 alleles each:
DL <- list(dL = c("A", "a"))
HL <- list(hL = c("B", "b"))
```

Next, a list of possible mutation events from one allele to another is defined, each associated with a mutation rate, defined as the proportion of copies that will be mutated at each meiosis event.

```
mutations <- list(
  mutation(from = "A", to = "a", rate = 1e-3),
  mutation(from = "B", to = "b", rate = 1e-3)
)
genomeObj <- Ease::setGenome(listHapLoci = HL, listDipLoci = DL)
mutMatrixObj <- Ease::setMutationMatrix(genomeObj, mutations = mutations)
```

The population is then defined, specifying its name (here ExamplePop), size (here 1,000 individuals), sexual system (here dioecy, i.e. 500 females and 500 males), genetic system and

102 mutation pattern. In addition to this, it is necessary to define the type of selection. Here, for
103 simplicity's sake, the alleles are defined as neutral.

```
# Definition of a population of 1000 individuals, males and females
pop <- Ease::setPopulation(
  name = "ExamplePop",
  size = 1000,
  dioecy = TRUE,
  genomeObj = genomeObj,
  mutMatrixObj = mutMatrixObj,
  selectionObj = Ease::setSelectNeutral(genomeObj)
)
print(pop)
#> ----- Population OBJECT -----
#>           in details
#> Population 'ExamplePop' of 1000 dioecious individuals
#> There is no demography.
#> The initial genotypes frequency are:
#> A/A||B A/a||B a/a||B A/A||b A/a||b a/a||b
#>      1      0      0      0      0      0
#> Selection:
#> No selection defined.
#> -----
```

104 Once the population has been defined, a metapopulation must be defined (here with only a
105 single population). Below, its evolution is simulated over 2,000 generations with the genotype,
106 haplotype and allele frequencies of each generation being recorded (by default when setting
107 recording = TRUE).

```
metapop <- Ease::setMetapopulation(populations = list(pop))
metapop <- simulate(metapop, threshold = 2000, seed = 123, recording = TRUE)
rec = Ease::getRecords(metapop)
print(which(rec$s1$ExamplePop$a == 1)[1])
#> [1] 1072
print(which(rec$s1$ExamplePop$b == 1)[1])
#> [1] 1481
```

108 The recorded frequencies of alleles a and b can then be extracted using the function getRecords.
109 Here, having defined a (fairly high) directional mutation rate towards these alleles, each fixed
110 after 1072 and 1481 generations, respectively.

111 For more details on how the package works and a more detailed view of the options, see the
112 package vignette and documentation.

113 Acknowledgements

114 I would like to thank Sylvain Glémin and Lucas Marie-Orleach for inspiring me to take this
115 modelling approach, and Vitor Sudbrack for his valuable feedback.

116 References

117 Baumdicker, F., Bisschop, G., Goldstein, D., Gower, G., Ragsdale, A. P., Tsambos, G., Zhu, S.,
118 Eldon, B., Ellerman, E. C., Galloway, J. G., Gladstein, A. L., Gorjanc, G., Guo, B., Jeffery,
119 B., Kretschumar, W. W., Lohse, K., Matschiner, M., Nelson, D., Pope, N. S., ... Kelleher,
120 J. (2022). Efficient ancestry and mutation simulation with msprime 1.0. *Genetics*, 220(3),
121 iyab229.

- 122 Bürger, R. (2000). *The mathematical theory of selection, recombination, and mutation*. Wiley.
123 ISBN: 978-0-471-98653-9
- 124 Crow, J. F., & Kimura, M. (1970). *An introduction to population genetics theory*. Burgess
125 Publishing Company. ISBN: 978-0-8087-2901-3
- 126 Ewens, W. J. (2004). *Mathematical Population Genetics* (S. S. Antman, J. E. Marsden, L.
127 Sirovich, & S. Wiggins, Eds.; Vol. 27). Springer New York. <https://doi.org/10.1007/978-0-387-21822-9>
128
- 129 Haller, B. C., & Messer, P. W. (2023). SLiM 4: Multispecies eco-evolutionary modeling. *The*
130 *American Naturalist*, 201(5), E127–E139. <https://doi.org/10.1086/723601>

DRAFT