

Step current response of the HH Model

Eleftherios Ioannidis
elefthei@mit.edu

James Hobin
hobinjk@mit.edu

MIT EECS

November 20, 2014

Three possible responses to a step current




Figure : Phase diagram for stimulation with a step current.




Figure : Simulated single action potential.




Figure : Simulated ring response.




Figure : Simulated train of repeating potentials.

Finding train frequency; LSSA

To find the train frequency we used the Least-Squares spectral analysis method (LSSA). LSSA is a method of estimating a frequency spectrum, based on a least squares fit of sinusoids to data samples, similar to Fourier analysis.

It works *better* than Fourier Analysis on data with variable time intervals such as the ones we are studying.

Train frequency over increasing input step

HERE




Figure : HH Models step current response starting at 0 $\mu\text{A}/\text{cm}^2$




Figure : HH Models step current response starting at 0 $\mu\text{A}/\text{cm}^2$




Figure : HH Models step current response starting at 0 $\mu\text{A}/\text{cm}^2$




Figure : HH Models step current response starting at 0 $\mu\text{A}/\text{cm}^2$




Figure : HH Models step current response starting at 0 $\mu\text{A}/\text{cm}^2$




Figure : HH Models step current response starting at 0 $\mu\text{A}/\text{cm}^2$




Figure : HH Models step current response starting at 0 $\mu\text{A}/\text{cm}^2$




Figure : HH Models step current response starting at 0 $\mu\text{A}/\text{cm}^2$




Figure : HH Models step current response starting at 0 $\mu\text{A}/\text{cm}^2$




Figure : HH Models step current response starting at 0 $\mu\text{A}/\text{cm}^2$




Figure : HH Models step current response starting at 0 $\mu\text{A}/\text{cm}^2$




Figure : HH Models step current response starting at 0 $\mu\text{A}/\text{cm}^2$




Figure : HH Models step current response starting at 0 $\mu\text{A}/\text{cm}^2$




Figure : HH Models step current response starting at 0 $\mu\text{A}/\text{cm}^2$




Figure : HH Models step current response starting at 0 $\mu\text{A}/\text{cm}^2$




Figure : HH Models step current response starting at 0 $\mu\text{A}/\text{cm}^2$




Figure : HH Models step current response starting at 0 $\mu\text{A}/\text{cm}^2$





Figure : Incorrect behavior due to low precision

References

- 1 Weiss, T. F. (1995). Cellular Biophysics. Volume 1: Transport, MIT Press.
- 2 Weiss, T. F. (1995). Cellular Biophysics. Volume 2: Electrical Properties, MIT Press.
- 3 Blaustein, M.P., Kao, J.P.Y., Matteson, D.R. (2012). Cellular Physiology and Neurophysiology, 2nd edition, Elsevier-Mosby.
- 4 Gerstner, Wulfram, and Werner M. Kistler. Spiking neuron models: Single neurons, populations, plasticity. Cambridge university press, 2002.