

Hilbert Spaces

H.1

Solution:

☐

H.2

Solution:

☐

H.3

Solution:

☐

H.4

Solution:

☐

H.5a**Solution:**☐

H.11**Solution:**☐

H.12**Solution:**☐

H.13**Solution:**☐

H.15

$$(u \mid v) = \int_{-\infty}^{\infty} u(x)v(x) \frac{e^{-x^2}}{\sqrt{\pi}} dx$$

Determine the first three Hermite polynomials

Solution:

Using Gram-Schmidt with $u_n = [1, x, x^2]$

$$\begin{aligned}\varphi_0 &= 1 \\ (\varphi_0 | \varphi_0) &= \int_{-\infty}^{\infty} \frac{e^{-x^2}}{\sqrt{\pi}} dx = 1 \\ \psi_0 &= \frac{\varphi_0}{(\varphi_0 | \varphi_0)} = 1\end{aligned}$$

Next

$$\begin{aligned}\varphi_1 &= u_1 - P_{[\varphi_0]} u_1 = u_1 - \frac{(\varphi_0 | u_1)}{(\varphi_0 | \varphi_0)} \varphi_0 = x - \frac{(1 | x)}{(1 | 1)} 1 = \\ &\quad \underbrace{x - \int_{-\infty}^{\infty} x \frac{e^{-x^2}}{\sqrt{\pi}} dx}_{=0 \text{ (odd function)}} = x \\ \psi_1 &= \frac{\varphi_1}{(\varphi_1 | \varphi_1)} = \frac{x}{(x | x)} \\ (x | x) &= \int_{-\infty}^{\infty} x^2 \frac{e^{-x^2}}{\sqrt{\pi}} dx = -\frac{1}{2\sqrt{\pi}} \int_{-\infty}^{\infty} (x) (-2xe^{-x^2}) dx = \\ &= -\frac{1}{2\sqrt{\pi}} \left([xe^{-x^2}]_{-\infty}^{\infty} - \int_{-\infty}^{\infty} e^{-x^2} dx \right) = \frac{1}{2}\end{aligned}$$

Finally this gives

$$\psi_1 = \frac{x}{(x | x)^{\frac{1}{2}}} = \sqrt{2}x$$

Next

$$\begin{aligned}
\varphi_2 &= u_2 - P_{[\varphi_0, \varphi_1]} u_2 = x^2 - \frac{(1 \mid x^2)}{(1 \mid 1)} 1 - \frac{(x \mid x^2)}{(x \mid x)} x = \\
&= x^2 - \int_{-\infty}^{\infty} x^2 \frac{e^{-x^2}}{\sqrt{\pi}} dx - \underbrace{\int_{-\infty}^{\infty} x^3 \frac{e^{-x^2}}{\sqrt{\pi}} dx}_{=0 \text{ odd function}} = x^2 - \frac{1}{2} \\
\psi_2 &= \frac{\varphi_2}{(\varphi_2 \mid \varphi_2)^{\frac{1}{2}}} = \left(x^2 - \frac{1}{2}\right) \left(\int_{-\infty}^{\infty} \left(x^2 - \frac{1}{2}\right)^2 \frac{e^{-x^2}}{\sqrt{\pi}} dx \right)^{-\frac{1}{2}} = \\
&= \left(x^2 - \frac{1}{2}\right) \left(\int_{-\infty}^{\infty} \left(x^4 - x^2 + \frac{1}{4}\right) \frac{e^{-x^2}}{\sqrt{\pi}} dx \right)^{-\frac{1}{2}} \\
&\quad \text{(i) } \int_{-\infty}^{\infty} \frac{1}{4} \frac{e^{-x^2}}{\sqrt{\pi}} dx = \boxed{\frac{1}{4}} \\
&\quad \text{(ii) } \int_{-\infty}^{\infty} -x^2 \frac{e^{-x^2}}{\sqrt{\pi}} dx = \boxed{-\frac{1}{2}} \\
&\quad \text{(iii) } \int_{-\infty}^{\infty} x^4 \frac{e^{-x^2}}{\sqrt{\pi}} dx = -\frac{1}{2\sqrt{\pi}} \int_{-\infty}^{\infty} (x^3)(-2xe^{-x^2}) dx = \\
&\quad = \underbrace{\left[x^2 e^{-x^2}\right]_{-\infty}^{\infty}}_{=0} - \underbrace{\int_{-\infty}^{\infty} 3x^2 e^{-x^2} dx}_{=\frac{3\sqrt{\pi}}{2}} = \boxed{\frac{3}{4}}
\end{aligned}$$

So

$$\psi_2 = \left(x^2 - \frac{1}{2}\right) \left(\frac{1}{4} - \frac{1}{2} + \frac{3}{4}\right)^{-\frac{1}{2}} = \left(x^2 - \frac{1}{2}\right) \sqrt{2}$$

The three first ortonormal Hermite polynomials are

$$\psi_0 = \boxed{1}$$

$$\psi_1 = \boxed{\sqrt{2}x}$$

$$\psi_2 = \boxed{\sqrt{2}\left(x^2 - \frac{1}{2}\right)}$$

□

H.18

$$\int_0^\pi (x - a \sin(x) - b \sin(2x))^2 dx$$

is minimized.

Solution:

$$v \in \mathcal{M} = [\sin(x), \sin(2x)]$$

$\sin(x)$ and $\sin(2x)$ are orthogonal and we can use the projection formula which says that the minimum is obtained when

$$v = P_{[\mathcal{M}]}u$$

$$v_1 = a \sin(x) = P_{[\sin(x)]}u = \frac{(\sin(x) | x)}{(\sin(x) | \sin(x))} \sin(x)$$

$$v_2 = b \sin(2x)$$

From this, a is calculated as

$$a = \frac{(\sin(x) | x)}{(\sin(x) | \sin(x))}$$

$$(\sin(x) | x) = \int_0^\pi x \sin(x) dx = -[x \cos(x)]_0^\pi + \int_0^\pi \pi \cos(x) dx = \pi$$

$$(\sin(x) | \sin(x)) = \int_0^\pi \sin^2 x dx = \int_0^\pi \frac{1 - \cos(2x)}{2} dx = \frac{\pi}{2}$$

↓

$$a = \frac{\pi}{\frac{\pi}{2}} = 2$$

and b is calculated in the same way as

$$b = \frac{(\sin(2x) | x)}{(\sin(2x) | \sin(2x))}$$

$$(\sin(2x) | x) = \int_0^\pi x \sin(2x) dx = -\left[\frac{x \cos(2x)}{2}\right]_0^\pi + \int_0^\pi \frac{\cos(2x)}{2} dx = -\frac{\pi}{2}$$


$$(\sin(2x) | \sin(2x)) = \int_0^\pi \sin^2(2x) dx = \int_0^\pi \frac{1 - \cos(2x)}{2} dx = \frac{\pi}{2}$$

↓

$$b = -\frac{\frac{\pi}{2}}{\frac{\pi}{2}} = -1$$

In general, the functions $v \in \mathcal{M} = \{\sin(nx)\}_{n=1}^{\infty}$ that best approximate $u(x) = x$ at the interval $[0, \pi]$ is given by

$$u_n = \frac{2(-1)^{n-1} \sin nx}{n}$$



□

H.19

(a) Give an orthogonal base in $\mathcal{V}$ with the scalar product

$$(u \mid v) = \int_{-1}^1 u(x)v(x) \, dx$$

(b) Let $u(x) = x^2 - x$. Determine $v \in \mathcal{V}$ so that

$$\int_{-1}^1 |u(x) - v(x)|^2 \, dx$$

is minimized.

(c) Determine the lowest possible value of the integral in (b).

Solution:(a) $\square$ **H.20****Solution:**

For this setup, the scalar product is defined as

$$(u | v) = \int_0^1 u(x)v(x)w(x) \, dx = \int_0^1 u(x)v(x)x \, dx$$

Find an orthogonal base

$$\varphi_0 = 1$$

$$(\varphi_0 | \varphi_0) = \int_0^1 x \, dx = \frac{1}{2}$$

$$\psi_0 = \frac{\varphi_0}{(\varphi_0 | \varphi_0)} = 2$$

$$\varphi_1 = x - P_{[\varphi_0]}x = x - \frac{(\varphi_0 | x)}{(\varphi_0 | \varphi_0)}\varphi_0 = x - 2 \int_0^1 x^2 \, dx = x - \frac{2}{3}$$

$$(\varphi_1 | \varphi_1) = \int_0^1 \left(x - \frac{2}{3}\right)^2 x \, dx = \frac{1}{36}$$

$$\psi_1 = \frac{\varphi_1}{(\varphi_1 | \varphi_1)} = 36x - 24$$

Now $v \in \mathcal{M} = \{2, 36x - 24\}$

Are looking for

$$\int_0^1 (x^4 - a\psi_0 - b\psi_1)^2 x \, dx$$

From which we can calculate

$$a\psi_0 = \frac{(\psi_0 | x^4)}{(\psi_0 | \psi_0)}\psi_0$$

$$a = \frac{(\psi_0 | u)}{(\psi_0 | \psi_0)}$$

$$(\psi_0 | u) = \int_0^1 2x^5 \, dx = \frac{1}{3}$$

$$(\psi_0 | \psi_0) = \int_0^1 4x \, dx = 2$$

$$a = \frac{1}{6}$$

and

$$b\psi_1 = \frac{(\psi_1 | x^4)}{(\psi_1 | \psi_1)}\psi_1$$

$$b = \frac{(\psi_1 | x^4)}{(\psi_1 | \psi_1)}$$


$$(\psi_1 | x^4) = \int_0^1 (36x^6 - 24x^5) \, dx = \frac{36}{7} - \frac{24}{6} = \frac{8}{7}$$

$$(\psi_1 | \psi_1) = \int_0^1 (1296x^3 - 1728x^2 + 576x) \, dx = 36$$

$$b = \frac{2}{63}$$

The first degree polynomial that best approximates the function $u(x) = x^4$ in $I = [0, 1]$ is

$$\frac{1}{6}\psi_0 + \frac{2}{63}\psi_1 = \frac{8}{7}x - \frac{3}{7}$$



(b) We are now looking for a second degree polynomial which means our $\psi \in \mathcal{M} = \{1, x, x^2\}$ ψ_0 and ψ_1 were already calculated in (a) so

$$\varphi_2 = u_2 - P_{[\varphi_0, \varphi_1]} u_2 = x^2 - \frac{(\varphi_0 | x^2)}{(\varphi_0 | \varphi_0)} \varphi_0 - \frac{(\varphi_1 | x^2)}{(\varphi_1 | \varphi_1)} \varphi_1$$

$$\frac{(\varphi_0 | x^2)}{(\varphi_0 | \varphi_0)} \varphi_0 = 2(\varphi_0 | x^2) = 2 \int_0^1 x^3 dx = \frac{1}{2}$$

$$\frac{(\varphi_1 | x^2)}{(\varphi_1 | \varphi_1)} \varphi_1 = (\varphi_1 | x^2)(36x - 24)$$

Now calculating

$$(\varphi_1 | x^2) = \int_0^1 \left(x^4 - \frac{2x^3}{3} \right) dx = \frac{1}{30}$$

which gives finally

$$\varphi_2 = x^2 - \frac{6}{5}x + \frac{3}{10}$$

$$(\varphi_2 | \varphi_2) = \int_0^1 \left(x^2 - \frac{6x}{5} + \frac{3}{10} \right)^2 x \, dx = \frac{1}{600}$$

$$\psi_2 = 600x^2 - 720x + 180$$

We can now find the best approximation with

$$\int_0^1 (x^4 - a\psi_0 - b\psi_1 - c\psi_2)^2 x \, dx$$

$$c\psi_2 = \frac{(\psi_2 | x^4)}{(\psi_2 | \psi_2)} \psi_2$$

$$c = \frac{(\psi_2 | x^4)}{(\psi_2 | \psi_2)}$$

$$(\psi_2 | x^4) = \int_0^1 (600x^2 - 720x + 180)x^4 \, dx = \frac{15}{7}$$


$$(\psi_2 | \psi_2) = \int_0^1 (600x^2 - 720x + 180)^2 x \, dx = 600$$

$$\downarrow$$

$$c = \frac{15}{4200} = \frac{1}{280}$$

The best second degree polynomial to approximate $u(x) = x^4$ is then

$$a\psi_0 + b\psi_1 + c\psi_2 = \frac{15}{7}x^2 - \frac{10}{7}x + \frac{3}{14}$$



□

H.27

- (a) Show that p_n is orthogonal to all polynomials of degree $p < n$
- (b) Show that if p is a polynomial, not identically zero, such that $(p_k | p) = 0$ for $k = 0, 1, 2$, then degree of $p \geq 3$.

Solution:

- (a) All polynomials of degree m can be written

$$p = \sum_{k=0}^m c_k p_k$$

Gives

$$(p_n | p) = \left(p_n | \sum_{k=0}^m c_k p_k \right) = \sum_{k=0}^m c_k (p_n | p_k) = 0 \text{ for } m < n$$

since all polynomials are pairwise orthogonal

(b)

$$p = \sum_{k=0}^m c_k p_k$$

where

$$c_k = \frac{(p_k | p)}{(p_k | p_k)}$$

$$(p_k | p) = 0, \text{ for } k = 0, 1, 2 \rightarrow c_0 = c_1 = c_2 = 0$$

The restriction that $p \neq 0$ gives that $\deg p \geq 3$ $\square$

H.30

Solution:

$\square$

H.31

Solution:

$\square$

H.33

$$f_n(x) = x^n, \quad x \in I = [0, 1], \quad n = 0, 1, \dots$$

Determine $\|f_n\|_p$ for $p = 1, 2$ and ∞ . Is it for any of these norms true that $f_n \rightarrow 0$ when $n \rightarrow \infty$? What is the pointwise limit function?

Solution:

$$\|f_n\|_1 = \int_0^1 |x^n| \, dx = \left[\frac{x^{n+1}}{n+1} \right]_0^1 = \frac{1}{n+1} \rightarrow 0, \quad n \rightarrow \infty$$

$$\|f_n\|_2 = \left(\int_0^1 |x^n|^2 \, dx \right)^{\frac{1}{2}} = \left(\left[\frac{x^{2n+1}}{2n+1} \right]_0^1 \right)^{\frac{1}{2}} = \frac{1}{\sqrt{2n+1}} \rightarrow 0, \quad n \rightarrow \infty$$

$$\|f_n\|_\infty = \sup_{x \in [0,1]} |f_n| = \sup_{x \in [0,1]} |x^n| = 1 \rightarrow 1, \quad n \rightarrow \infty$$

□

H.36

Solution:

□

H.38

Solution:

□

H.39

Solution:

□

H.45**Solution:**

□

H.47**Solution:**

□

H.50

$$\mathcal{A} = -\sqrt{1-x^2} \frac{d}{dx} \left(\sqrt{1-x^2} \frac{d}{dx} \right)$$

$$D_{\mathcal{A}} = \{u \in \mathcal{C}^2(-1, 1) \mid u \text{ and } u' \text{ bounded/limited}\}$$

- (a) Determine a scalar product in which $\mathcal{A}$ is symmetric.
- (b) Show that $\mathcal{A}$ is symmetric in the scalar product from (a), using the definition.
- (c) Show that $1, x, 2x^2 - 1$ are eigenfunctions to $\mathcal{A}$ and determine the corresponding eigenvalues.
- (d) Evaluate

$$\int_{-1}^1 \frac{2x^2 - 1}{\sqrt{1-x^2}} dx$$

Solution:

- (a) The operator is a Sturm-Liouville operator

$$\mathcal{A}u = -\sqrt{1-x^2} \frac{d}{dx} \left(\sqrt{1-x^2} \frac{d}{dx} \right) u = -\frac{1}{w} (pu')', \quad w(x) = \frac{1}{\sqrt{1-x^2}}$$

and the scalar product is then

$$(u \mid v) = \int_{-1}^1 \overline{u(x)} v(x) \frac{1}{w(x)} dx$$

- (b) Using the scalar product definition

$$(u \mid v) = \int \overline{u(x)} v(x) w(x) \, dx, \quad w(x) = \frac{1}{\sqrt{1-x^2}}$$

we get

$$\begin{aligned}
 (u \mid \mathcal{A}v) &= \\
 &= \int_{-1}^1 \overline{u(x)} [\mathcal{A}v(x)] \frac{1}{w(x)} \, dx = \int_{-1}^1 \overline{u(x)} \left[-\sqrt{1-x^2} \frac{d}{dx} \left(\sqrt{1-x^2} \frac{d}{dx} v(x) \right) \right] \frac{1}{\sqrt{1-x^2}} \, dx = \\
 &= - \int_{-1}^1 \overline{u(x)} \left(\sqrt{1-x^2} v'(x) \right)' \, dx = - \underbrace{\left[\overline{u(x)} \sqrt{1-x^2} v'(x) \right]_{-1}^1}_{=0} + \int_{-1}^1 \overline{u'(x)} \sqrt{1-x^2} v'(x) \, dx = \\
 &= \int_{-1}^1 \left(\overline{u'(x)} \sqrt{1-x^2} \right) v'(x) \, dx = \underbrace{\left[\left(\overline{u'(x)} \sqrt{1-x^2} \right) v(x) \right]_{-1}^1}_{=0} - \int_{-1}^1 \left(\overline{u'(x)} \sqrt{1-x^2} \right)' v(x) \, dx = \\
 &= \int_{-1}^1 - \left(\overline{u'(x)} \sqrt{1-x^2} \right)' v(x) \, dx = \int_{-1}^1 -\sqrt{1-x^2} \left(\overline{u'(x)} \sqrt{1-x^2} \right)' v(x) \frac{1}{\sqrt{1-x^2}} \, dx = \\
 &= \int_{-1}^1 \left[-\sqrt{1-x^2} \frac{d}{dx} \left(\sqrt{1-x^2} \frac{d}{dx} \overline{u(x)} \right) \right] v(x) \frac{1}{\sqrt{1-x^2}} \, dx = \\
 &= \int_{-1}^1 [\mathcal{A}u(x)] v(x) \frac{1}{w(x)} \, dx = (\mathcal{A}u \mid v)
 \end{aligned}$$

(c)

$$\mathcal{A}u = \lambda u$$

$$u(x) = 1$$

$$\downarrow$$

$$\mathcal{A} \cdot 1 = \lambda_0 \cdot 1 \rightarrow -\sqrt{1-x^2} \frac{d}{dx} \left(\sqrt{1-x^2} \frac{d}{dx} \cdot 1 \right) = \lambda_0 \cdot 1 \rightarrow 0 = \lambda_0 \cdot 1 \rightarrow \boxed{\lambda_0 = 0}$$

$$u(x) = x$$

$$\downarrow$$

$$\mathcal{A}x = \lambda_1 x \rightarrow -\sqrt{1-x^2} \frac{d}{dx} \left(\sqrt{1-x^2} \frac{d}{dx} x \right) = \lambda_1 x \rightarrow -\sqrt{1-x^2} \frac{-2x}{2\sqrt{1-x^2}} = \lambda_1 x \rightarrow$$

$$\rightarrow x = \lambda_1 x \rightarrow \boxed{\lambda_1 = 1}$$

$$u(x) = 2x^2 - 1$$

$$\downarrow$$

$$\begin{aligned} \mathcal{A}(2x^2 - 1) &= \lambda_2(2x^2 - 1) \rightarrow -\sqrt{1-x^2} \frac{d}{dx} \left(\sqrt{1-x^2} \frac{d}{dx} (2x^2 - 1) \right) = \lambda_2(2x^2 - 1) \rightarrow \\ &\rightarrow -\sqrt{1-x^2} \frac{d}{dx} (4x\sqrt{1-x^2}) = -\sqrt{1-x^2} \left(4\sqrt{1-x^2} - \frac{8x^2}{2\sqrt{1-x^2}} \right) = -4 + 4x^2 + 4x^2 = \\ &= 8x^2 - 4 = \lambda_2(2x^2 - 1) \rightarrow 4(2x^2 - 1) = \lambda_2(2x^2 - 1) \rightarrow \boxed{\lambda_2 = 4} \end{aligned}$$

(d) The integral can be rewritten with the scalar product in (a) as

$$\int_{-1}^1 \frac{2x^2 - 1}{\sqrt{1-x^2}} dx = (1 \mid 2x^2 - 1), \quad w(x) = \frac{q}{\sqrt{1-x^2}}$$

Since the functions 1 and $2x^2 - 1$ are eigenfunctions, the scalar product between them are pairwise orthogonal and therefore = 0.

□

H.51

Solution:

□

H.52**Solution:**

□

Operator

$$\mathcal{A} = -\text{laplace} = -\frac{\partial^2}{\partial x^2} - \frac{\partial^2}{\partial y^2}$$

Solution:

$$\begin{aligned}
(u \mid \mathcal{A}v) &= \int_0^a \int_0^b \bar{u}(-v''_{xx} - v''_{yy}) \, dx \, dy = \\
(1) &= - \int_0^b \underbrace{\left(\int_0^a \bar{u} v''_{xx} \, dx \right)}_{(i-x)} \, dy - \int_0^a \underbrace{\left(\int_0^b \bar{u} v''_{yy} \, dy \right)}_{(i-y)} \, dx \\
(i-x) &= \underbrace{[\bar{u} v'_x]_0^a}_{=0} - \int_0^a \bar{u}'_x v'_x \, dx \\
(i-y) &= \underbrace{[\bar{u} v'_y]_0^b}_{=0} - \int_0^b \bar{u}'_y v'_y \, dy \\
&\downarrow \\
(1) &= \int_0^b \underbrace{\left(\int_0^a \bar{u}'_x v'_x \, dx \right)}_{(ii-x)} \, dy + \int_0^a \underbrace{\left(\int_0^b \bar{u}'_y v'_y \, dy \right)}_{(ii-y)} \, dx = (***) \\
(ii-x) &= \underbrace{[\bar{u}'_x v'_x]_0^a}_{=0} - \int_0^a \bar{u}''_{xx} v \, dx \\
(ii-y) &= \underbrace{[\bar{u}'_y v'_y]_0^b}_{=0} - \int_0^b \bar{u}''_{yy} v \, dy \\
&\downarrow \\
(1) &= - \int_0^b \int_0^a \bar{u}''_{xx} v \, dx \, dy - \int_0^a \int_0^b \bar{u}''_{yy} v \, dy \, dx = \int_0^a \int_0^b (-\bar{u}''_{xx} - \bar{u}''_{yy}) v \, dx \, dy = \\
&= \boxed{(\mathcal{A}u \mid v)}
\end{aligned}$$

To show positive semidefinite, use the form described on (***) and $u = v$:

$$(***) = \int_0^a \int_0^b (\bar{u}'_x u'_x + \bar{u}'_y u'_y) \, dx \, dy = \int_0^a \int_0^b (|u'_x|^2 + |u'_y|^2) \, dx \, dy \geq 0$$

To find the eigenvalues associated with the operator, use separation of variables

$$\begin{aligned}
u(x, y) &= X(x)Y(y) \\
\mathcal{A}u &= \lambda u \rightarrow -X''Y - XY'' = \lambda XY \rightarrow -\frac{X''}{X} - \frac{Y''}{Y} = \lambda \\
&\downarrow \\
\begin{cases} \frac{X''}{X} = -\gamma_x^2, & X(0) = X(a) = 0 \\ \frac{Y''}{Y} = -\gamma_y^2, & Y'(0) = Y'(b) = 0 \end{cases}
\end{aligned}$$

Starting with $X(x)$:

$$X'' = -\gamma_x^2 X \rightarrow X(x) = e^{i\gamma_x x} = A \cos(\gamma_x x) + B \sin(\gamma_x x)$$

$$X(0) = A = 0 \rightarrow A = 0$$

$$X(a) = B \sin(\gamma_x a) = 0 \rightarrow \gamma_x = \frac{\pi k_x}{a}, k_x = 1, 2, 3 \dots$$

↓

$$X(x) = B \sin\left(\frac{\pi k_x x}{a}\right) \quad \square$$

Now $Y(y)$:

$$Y'' = -\gamma_y^2 Y \rightarrow Y(y) = e^{i\gamma_y y} = A \cos(\gamma_y y) + B \sin(\gamma_y y)$$

$$Y'(x) = -\gamma_y A \sin(\gamma_y y) + \gamma_y B \cos(\gamma_y y)$$

$$Y'(0) = \gamma_y B = 0 \rightarrow B = 0$$

$$Y'(b) = -\gamma_y A \sin(\gamma_y b) = 0 \rightarrow \gamma_y = \frac{\pi k_y}{b}, k_y = 0, 1, 2 \dots$$

↓

$$Y(y) = A \cos\left(\frac{\pi k_y y}{b}\right) \quad \square$$

□

H.8 Operators in Hilbert Spaces

Linear avbildning

$$\mathcal{A} : \mathcal{H}_1 \rightarrow \mathcal{H}_2$$

where $\mathcal{H}_1$ and $\mathcal{H}_2$ are two pre-Hilbert Spaces. Linearity:

$$\mathcal{A}(\lambda u + \mu v) = \lambda \mathcal{A}u + \mu \mathcal{A}v$$

is termed a **linear operator**.

Example H.19

Let I be a compact interval on $\mathbb{R}$, and let $K(x, y)$ be continuous for $x, y \in I$.

$$v(x) = \int_I K(x, y) u(y) \, dy$$

This defines an **integral operator**

$$\mathcal{K} : L_2(I) \rightarrow L_2(I)$$

The function $K(x, y)$ is termed the operators **kernel**