

Potentials and Fields

Problem 10.1

Show that the differential equations for V and $\mathbf{A}$

$$\begin{aligned}\nabla^2 V + \frac{\partial}{\partial t}(\nabla \cdot \mathbf{A}) &= -\frac{1}{\epsilon_0} \rho \\ \left(\nabla^2 \mathbf{A} - \mu_0 \epsilon_0 \frac{\partial^2 \mathbf{A}}{\partial t^2} \right) - \nabla \left(\nabla \cdot \mathbf{A} + \mu_0 \epsilon_0 \frac{\partial V}{\partial t} \right) &= -\mu_0 \mathbf{J}\end{aligned}$$

can be written in the more symmetrical form

$$\begin{aligned}\square^2 V + \frac{\partial L}{\partial t} &= -\frac{1}{\epsilon_0} \rho \\ \square^2 \mathbf{A} - \nabla L &= -\mu_0 \mathbf{J}\end{aligned}$$

where

$$\square^2 \equiv \nabla^2 - \mu_0 \epsilon_0 \frac{\partial^2}{\partial t^2} \quad \text{and} \quad L \equiv \nabla \cdot \mathbf{A} + \mu_0 \epsilon_0 \frac{\partial V}{\partial t}$$

Solution:

Put the definitions of $\square^2$ and L into the equations for V :

$$\begin{aligned}\left(\nabla^2 - \mu_0 \epsilon_0 \frac{\partial^2}{\partial t^2} \right) V + \frac{\partial}{\partial t} \left(\nabla \cdot \mathbf{A} + \mu_0 \epsilon_0 \frac{\partial V}{\partial t} \right) &= \\ = \nabla^2 V - \mu_0 \epsilon_0 \frac{\partial^2 V}{\partial t^2} + \frac{\partial(\nabla \cdot \mathbf{A})}{\partial t} + \mu_0 \epsilon_0 \frac{\partial^2 V}{\partial t^2} &= \\ = \nabla^2 V + \frac{\partial}{\partial t}(\nabla \cdot \mathbf{A}) &= -\frac{1}{\epsilon_0} \rho \quad \# \end{aligned}$$

and for $\mathbf{A}$:

$$\begin{aligned} & \left(\nabla^2 - \mu_0 \epsilon_0 \frac{\partial^2}{\partial t^2} \right) \mathbf{A} - \nabla \left(\nabla \cdot \mathbf{A} + \mu_0 \epsilon_0 \frac{\partial V}{\partial t} \right) = \\ & = \left(\nabla^2 \mathbf{A} - \mu_0 \epsilon_0 \frac{\partial^2 \mathbf{A}}{\partial t^2} \right) - \nabla \left(\nabla \cdot \mathbf{A} + \mu_0 \epsilon_0 \frac{\partial V}{\partial t} \right) = -\mu_0 \mathbf{J} \quad \# \end{aligned}$$

□

Problem 10.2

Problem 10.3

(a) Find the fields, and the charge and current distributions, corresponding to

$$V(\mathbf{r}, t) = 0, \quad \mathbf{A}(\mathbf{r}, t) = -\frac{1}{4\pi\epsilon_0} \frac{qt}{r^2} \hat{\mathbf{r}}$$

(b) Use the gauge function $\lambda = -(1/4\pi\epsilon_0)(qt/r)$ to transform the potentials, and comment on the result.

Solution:

The $\mathbf{E}$ and $\mathbf{B}$ fields are

$$\begin{aligned} \mathbf{E} &= -\nabla V - \frac{\partial \mathbf{A}}{\partial t} \\ \mathbf{B} &= \nabla \times \mathbf{A} \end{aligned}$$

The derivative of $\mathbf{A}$ with respect to time is

$$\frac{\partial \mathbf{A}}{\partial t} = -\frac{1}{4\pi\epsilon_0} \frac{q}{r^2} \hat{\mathbf{r}},$$

while the curl of $\mathbf{A}$ in spherical coordinates is

$$\frac{1}{r \sin \theta} \left[\frac{\partial}{\partial \theta} (\sin \theta A_\phi) - \frac{\partial A_\theta}{\partial \phi} \right] \hat{\mathbf{r}} + \frac{1}{r} \left[\frac{1}{\sin \theta} \frac{\partial A_r}{\partial \phi} - \frac{\partial}{\partial r} (r v_\phi) \right] \hat{\boldsymbol{\theta}} + \frac{1}{r} \left[\frac{\partial}{\partial r} (r A_\theta) - \frac{\partial A_r}{\partial \theta} \right] \hat{\boldsymbol{\phi}}.$$

Since $A_\phi = A_\theta = 0$, and $\frac{\partial A_r}{\partial \phi} = \frac{\partial A_r}{\partial \theta} = 0$, the curl of $\mathbf{A}$ is zero.

(a)

$$\begin{aligned} \mathbf{E} &= -\frac{\partial \mathbf{A}}{\partial t} = \frac{1}{4\pi\epsilon_0} \frac{q}{r^2} \hat{\mathbf{r}} \\ \mathbf{B} &= \nabla \times \mathbf{A} = 0 \end{aligned}$$

(b) Using the gauge function, V and $\mathbf{A}$ are transformed into

$$\begin{aligned} V'(\mathbf{r}, t) &= V(\mathbf{r}, t) - \frac{\partial \lambda}{\partial t} \\ \mathbf{A}'(\mathbf{r}, t) &= \mathbf{A}(\mathbf{r}, t) + \nabla \lambda \end{aligned}$$

Writing out the derivative and gradient of λ

$$\begin{aligned} \frac{\partial \lambda}{\partial t} &= -\frac{1}{4\pi\epsilon_0} \frac{q}{r} \\ \nabla \lambda &= \frac{1}{4\pi\epsilon_0} \frac{qt}{r^2} \hat{\mathbf{r}} \end{aligned}$$

gives the transformed potentials

$$\begin{aligned} V'(\mathbf{r}, t) &= \frac{1}{4\pi\epsilon_0} \frac{q}{r} \\ \mathbf{A}'(\mathbf{r}, t) &= -\frac{1}{4\pi\epsilon_0} \frac{qt}{r^2} \hat{\mathbf{r}} + \frac{1}{4\pi\epsilon_0} \frac{qt}{r^2} = 0 \end{aligned}$$

$\mathbf{E}$ is now given by $-\nabla V$ (instead of by $-\frac{\partial \mathbf{A}}{\partial t}$), while $\mathbf{B}$ is again 0, this time directly because $\nabla \times \mathbf{A} = \nabla \times 0 = 0$.

$$\begin{aligned} \mathbf{E} &= -\nabla V = -\frac{1}{4\pi\epsilon_0} \frac{q}{r^2} \hat{\mathbf{r}} \\ \mathbf{B} &= \nabla \times \mathbf{A} = 0 \end{aligned}$$

The result is exactly the same as in (a).

□

Problem 10.4

Problem 10.5

Problem 10.6

Problem 10.7

Problem 10.8

Problem 10.9

Problem 10.10

Problem 10.11

Problem 10.12

Problem 10.13

Problem 10.14

Problem 10.15

Problem 10.16

Problem 10.17

Problem 10.18

Problem 10.19

Problem 10.20

Problem 10.21

Problem 10.22

Problem 10.23

Problem 10.24

Problem 10.25

Problem 10.26

Problem 10.27

Problem 10.28

Problem 10.29

Problem 10.30

Problem 10.31

Problem 10.32

Problem 10.33

Problem 10.34

Problem 10.35

Problem 10.36

Problem 10.37

Problem 10.38