

Electrodynamics and Relativity

Problem 12.1

Problem 12.2

Problem 12.3

Problem 12.4

As the outlaws escape in their getaway car, which goes $\frac{3}{4}c$, the police officer fires a bullet from the pursuit car, which only goes $\frac{1}{2}c$. The muzzle velocity of the bullet (relative to the gun) is $\frac{1}{3}c$. Does the bullet reach its target (a) according to Galileo, (b) according to Einstein?

Solution:

(a) Adding up the velocities with Galileo's velocity addition rule, the resulting velocity of the bullet is

$$\frac{1}{2}c + \frac{1}{3}c = \frac{5}{6}c > \frac{3}{4}c$$

so the bullet does reach its target.

(b) Using Einstein's velocity addition rule, the result is instead

$$v_{\text{bullet}} = \frac{v_{\text{car}} + v_{\text{muzzle}}}{1 + \frac{v_{\text{car}} v_{\text{muzzle}}}{c^2}} = \frac{\frac{5}{6}c}{1 + \frac{1}{6}} = \frac{5}{7}c < \frac{3}{4}c$$

so the bullet does not reach its target. $\square$

Problem 12.5**Problem 12.6****Problem 12.7**

In a laboratory experiment, a muon is observed to travel 800 m before disintegrating. A graduate student looks up the lifetime of a muon (2×10^{-6} s) and concludes that its speed was

$$v = \frac{800 \text{ m}}{2 \times 10^{-6} \text{ s}} = 4 \times 10^8 \text{ m/s}.$$

Faster than light! Identify the student's error, and find the actual speed of this muon.

Solution:

The student did not take into account the lifetime of the muon in the muons reference time. The muons lifetime in its own reference time is

$$\gamma t = \frac{1}{\sqrt{1 - \left(\frac{v}{c}\right)^2}} t$$

If it travels for 800m then its speed is

$$\frac{800 \text{ m}}{\gamma \times 2 \times 10^{-6} \text{ s}} = v$$

Simplifying the distance over time gives

$$\frac{800 \text{ m}}{2 \times 10^{-6} \text{ s}} = \frac{4}{3} c$$

Solving for v

$$\begin{aligned}
\frac{4}{3}c &= \gamma v = \frac{v}{\sqrt{1 - \left(\frac{v}{c}\right)^2}} \\
&\downarrow \\
\left(\frac{4}{3}c\right)^2 &= \frac{v^2}{1 - \left(\frac{v}{c}\right)^2} \\
&\downarrow \\
\left(\frac{4}{3}c\right)^2 \left(1 - \left(\frac{v}{c}\right)^2\right) &= v^2 \\
&\downarrow \\
\left(\frac{4}{3}c\right)^2 &= v^2 \left(1 + \left(\frac{4}{3}c\right)^2 \frac{1}{c^2}\right) = v^2 \left(1 + \frac{16}{9}\right) \\
&\downarrow \\
v &= \frac{4}{3}c \frac{1}{\sqrt{1 + \frac{16}{9}}} = \frac{4}{3}c \frac{1}{\sqrt{\frac{25}{9}}} = \frac{12}{15}c = \boxed{\frac{4}{5}c}
\end{aligned}$$

□

Problem 12.8

Problem 12.9

A Lincoln Continental is twice as long as a VW Beetle, when they are at rest. As the Continental overtakes the VW, going through a speed trap, a (stationary) policeman observes that they both have the same length. The VW is going at half the speed of light. How fast is the Lincoln going?

Solution:

The length of the cars while stationary is denoted L . With length contraction due to the velocity accounted for the length is denoted L' . With v_{vw} given and $L_{\text{lc}} = 2L_{\text{vw}}$, the solution is obtained by setting $L'_{\text{lc}} = L'_{\text{vw}}$ and solving for v_{lc} .

$$L_{lc} = 2L_{vw}$$

$$L'_{vw} = L_{vw} \sqrt{1 - \frac{v_{vw}^2}{c^2}} = L_{vw} \frac{\sqrt{3}}{2}$$

$$L'_{vw} = L'_{lc} = L_{lc} \sqrt{1 - \frac{v_{lc}^2}{c^2}} = L_{vw} \frac{\sqrt{3}}{2}$$

Using $L_{lc} = 2L_{vw}$ and solving for v_{lc} , the velocity for which the length of the Lincoln Continental equals the length of the VW Beetle is

$$v_{lc} = \frac{\sqrt{13}}{4}c$$

□

Problem 12.10

Problem 12.11

Problem 12.12

Solve

- (i) $\bar{x} = \gamma(x - vt)$
- (ii) $\bar{y} = y$
- (iii) $\bar{z} = z$
- (iv) $\bar{t} = \gamma\left(t - \frac{v}{c^2}x\right)$

for x, y, z, t in terms of $\bar{x}, \bar{y}, \bar{z}, \bar{t}$, and check that you recover

- (i') $x = \gamma(\bar{x} + v\bar{t})$
- (ii') $y = \bar{y}$
- (iii') $z = \bar{z}$
- (iv') $t = \gamma\left(\bar{t} + \frac{v}{c^2}\bar{x}\right)$

Solution:

From (i) and (iv) we have

$$(i) \quad \bar{x} = \gamma(x - vt) \rightarrow x = \frac{\bar{x}}{\gamma} + vt$$

$$(iv) \quad \bar{t} = \gamma\left(t - \frac{v}{c^2}x\right) \rightarrow t = \frac{\bar{t}}{\gamma} + \frac{v}{c^2}x$$

Inserting t from (iv) into the expression for x in (i) gives

$$x = \frac{\bar{x}}{\gamma} + v\left(\frac{\bar{t}}{\gamma} + \frac{v}{c^2}x\right) \rightarrow x\left(1 - \frac{v^2}{c^2}\right) = \frac{\bar{x}}{\gamma} + \frac{v\bar{t}}{\gamma}$$

↓

$$x \frac{1}{\gamma^2} = \frac{1}{\gamma}(\bar{x} + v\bar{t}) \rightarrow x = \gamma(\bar{x} + v\bar{t}) = (i')$$

Putting this expression for x into the expression for t in (iv) gives

$$t = \frac{\bar{t}}{\gamma} + \frac{v}{c^2}(\gamma(\bar{x} + v\bar{t})) = \gamma\left(\frac{\bar{t}}{\gamma^2} + \frac{v}{c^2}\bar{x} + \frac{v^2}{c^2}\bar{t}\right) = \gamma\left(\left(1 - \frac{v^2}{c^2} + \frac{v^2}{c^2}\right)\bar{t} + \frac{v}{c^2}\bar{x}\right)$$

↓

$$t = \gamma\left(\bar{t} + \frac{v}{c^2}\bar{x}\right) = (iv')$$

(ii) $\leftrightarrow$ (ii') and (iii) $\leftrightarrow$ (iii') follow directly from the equations.

□

Problem 12.13

Problem 12.14

Problem 12.15

Problem 12.16

Problem 12.17

Problem 12.18

Problem 12.19

Problem 12.20

Problem 12.21

Problem 12.22

Problem 12.23

Problem 12.24

Problem 12.25

Problem 12.26

Problem 12.27

Problem 12.28

Problem 12.29

Problem 12.30

Problem 12.31

Problem 12.32

Problem 12.33

Problem 12.34

Problem 12.35

Problem 12.36

Problem 12.37

Problem 12.38

Problem 12.39

Problem 12.40

Problem 12.41

Problem 12.42

Problem 12.43

Problem 12.44

Problem 12.45

Problem 12.46

Problem 12.47

Problem 12.48

Problem 12.49

Problem 12.50

Problem 12.51

Problem 12.52

Problem 12.53

Problem 12.54

Problem 12.55

Problem 12.56

Problem 12.57

Problem 12.58

Problem 12.59

Problem 12.60

Problem 12.61

Problem 12.62

Problem 12.63

Problem 12.64

Problem 12.65

Problem 12.66

Problem 12.67

Problem 12.68

Problem 12.69

Problem 12.70

Problem 12.71

Problem 12.72

Problem 12.73

Problem 12.74

Problem 12.75

Problem 12.76