

EFFECTFUL MEALY MACHINES :

Filippo Bonchi,
Elena Di Lascio and Mario Román

EFFECTFUL TRIPLES.

$$\mathcal{V} \rightarrow \mathcal{L} \rightarrow \mathcal{C}$$

cartesian \rightarrow monoidal \rightarrow premonoidal

values \rightarrow local computations \rightarrow effectful computations

BISIMULATION AND TRACE

Università di Pisa

University of Oxford

Ex. Set \rightarrow TRel \rightarrow Rel

Set \rightarrow Stoch \rightarrow Substoch


Set \rightarrow Stoch \rightarrow State(Stoch)

Set \rightarrow Kl(Z(T)) \rightarrow KlT

cart(c ℓ) \rightarrow Z(c ℓ) \rightarrow c ℓ

MEALY MACHINES.

- $\delta: M \otimes A \rightarrow M \otimes B$
 \rightsquigarrow transition morphism
- $\delta_0: I \rightarrow M$
 \rightsquigarrow initial state




THM. Mealy machines form a feedback effectful triple $\text{Mealy}(U, L, \mathcal{C})$.

EX. coalgebraic Mealy machines :

$$M \rightarrow T(M \otimes B)^A \quad \leftrightarrow \quad M \otimes A \rightarrow M \otimes B$$

BISIMULATION. $(g, s) \equiv (g, t)$ if there is



where morphisms $(g, s) \xrightarrow{u} (h, r)$ are


$$\begin{array}{c} M \\ A \\ R \end{array} = \boxed{g} \xrightarrow{u} \begin{array}{c} N \\ B \\ R \end{array} = \begin{array}{c} M \\ A \\ R \end{array} = \boxed{h} \begin{array}{c} N \\ B \\ R \end{array}$$

$$\boxed{s} \xrightarrow{u} \boxed{N} = \boxed{r} \boxed{N}$$


PROP. For coalgebraic Mealy machines, effectful bisimilarity is coalgebraic bisimilarity
(for T preserving weak pullbacks).

EFFECTFUL STREAMS. $\delta: A \rightarrow B$ is

- $M_\delta \in \text{objcl}$
- $\delta^\circ: A^\circ \rightarrow M_\delta \otimes B^\circ$
- $\delta^+: M_\delta \cdot A^+ \rightarrow B^+$




quotiented by sliding local computations:




THM. Effectful streams form a feedback effectful triple Stream(V, L, cl).

EFFECTFUL TRACES. Execution of Mealy machines

$\text{Tr}: \text{Mealy}(\mathcal{V}, \mathcal{L}, \mathcal{C}) \rightarrow \text{Stream}(\mathcal{V}, \mathcal{L}, \mathcal{C})$




THM. Trace is universal: $\text{Eff} \xrightarrow{\text{Mealy}} \text{FBR} \xrightarrow{\text{Eff}} \text{UFBR} \xrightarrow{\text{Unif}} \text{Eff}$



COR. Bisimilarity implies trace equivalence.

CAUSAL PROCESSES. $p : A \rightarrow B$ is

$\{p_m : A_0 \otimes \dots \otimes A_m \rightarrow B_0 \otimes \dots \otimes B_m\}$ such that



THM. \mathcal{C} copy-discard with conditionals


$\Rightarrow \text{Proc}(\mathcal{C})$ copy-discard.

THM. \mathcal{C} copy-discard with conditionals and ranges


$\Rightarrow \text{Proc}(\mathcal{C}) \simeq \text{Stream}(\text{cart}(\mathcal{C}), \text{tot}(\mathcal{C}), \mathcal{C})$.

TRACES ARE EFFECTFUL TRACES.

CONDITIONALS





CAUSALITY CONDITION





TRACE PREDICATE



$$\Delta_0 = \Delta \wedge \forall i \leq n \\ (\delta_{i+1}, b_i) = f(\delta_i, a_i)$$





$$\Delta_0 = \Delta \wedge \forall i \leq n \\ (\delta_{i+1}, b_i) = f(\delta_i, a_i)$$



$$\exists \Delta_0, \dots, \Delta_{m+1} \Delta_0 \in \Delta \\ \wedge \forall i \leq n (\delta_{i+1}, b_i) \in f(\delta_i, a_i)$$



$$\sum_{\Delta_0, \dots, \Delta_{m+1}} \Delta(\Delta_0) \\ \cdot \prod_{i \leq m} f(\delta_{i+1}, b_i | \Delta_i, a_i)$$



$$\sum_{\Delta_0, \dots, \Delta_{m+1}} \Delta(\Delta_0) \\ \cdot \prod_{i \leq m} f(\delta_{i+1}, b_i | \Delta_i, a_i)$$