

# ToothGrowth

Crescita dei denti nelle cavie in base a diversi livelli di  
vitamina C

Elena Rosa Zacheo

Università di Padova

## Indice

<i>Titolo 1</i>	<b>1</b>
<i>Titolo 2</i>	<b>1</b>
<i>Titolo 3</i>	<b>2</b>

### *Titolo 1*

### *Titolo 2*

La *cavia* è un roditore erbivoro che appartiene all'ordine degli isticomorfi insieme al degu e al cincilla. Le cavie diversamente dal coniglio, partoriscono prole atta (ecco perché la gravidanza è particolarmente lunga, circa 60 gg): che nasce già provvista di pelo, in grado di alimentarsi da

sola; questo non significa che i cuccioli possano essere subito tolti dalla mamma perché necessitano, come tutti i mammiferi delle cure parenterali.


Si tratta di una specie ad alto carattere sociale che richiede molta compagnia, per questo motivo in alcuni Stati, ad esempio in Svizzera, è vietato per legge detenerne un solo esemplare. I porcellini d'India devono avere a disposizione un ambiente ampio e pulito a superficie morbida, costituito ad esempio da uno spesso strato di fieno, e asciutta per evitare di causare lesioni alle zampe. Devono poter scorrazzare fuori dalla gabbia.

```
summary(cars)
```

```
##           speed           dist
##  Min.      : 4.0    Min.      :  2.00
##  1st Qu.:12.0    1st Qu.: 26.00
##  Median :15.0    Median : 36.00
##  Mean   :15.4    Mean    : 42.98
##  3rd Qu.:19.0    3rd Qu.: 56.00
##  Max.    :25.0    Max.     :120.00
```

### *Titolo 3*

You can also embed plots, for example:



Note that the `echo = FALSE` parameter was added to the code chunk to prevent printing of the R code that generated the plot.