

Models

1 Data

Adjacency matrix $\mathbf{Y} = [y_{i,i'}]$ corresponding to an undirected, binary network.

2 Erdos–Renyi Model

Likelihood

$$y_{i,i'} \mid \theta \stackrel{\text{ind}}{\sim} \text{Ber}(\theta)$$

Prior

$$\theta \sim \text{Beta}(a_\theta, b_\theta)$$

Parameters

$$\Upsilon = (\theta) .$$

Hyper-parameters

$$(a_\theta, b_\theta) .$$

Posterior

$$p(\theta \mid \mathbf{Y}) = p(\mathbf{Y} \mid \theta) p(\theta) = \theta^{a_\theta + s_y - 1} (1 - \theta)^{b_\theta + N - s_y - 1} = \text{Beta}(\theta \mid a_\theta + s_y, b_\theta + N - s_y) ,$$

where $s_y = \sum_{i=1}^{I-1} \sum_{i'=i+1}^I y_{i,i'}$ and $N = I(I-1)/2$.

Prior Elicitation

$$a_\theta = 1 , \quad b_\theta = 1 .$$

3 Distance Model

Likelihood

$$y_{i,i'} \mid \zeta, \mathbf{u}_i, \mathbf{u}_{i'} \stackrel{\text{ind}}{\sim} \text{Ber}(\text{expit}(\zeta - \|\mathbf{u}_i - \mathbf{u}_{i'}\|))$$

Prior

$$\begin{aligned} \mathbf{u}_i &\mid \sigma^2 \stackrel{\text{iid}}{\sim} \mathbf{N}(\mathbf{0}, \sigma^2 \mathbf{I}) \\ \sigma^2 &\sim \text{IGam}(a_\sigma, b_\sigma) \\ \zeta &\mid \omega^2 \sim \mathbf{N}(0, \omega^2) \\ \omega^2 &\sim \text{IGam}(a_\omega, b_\omega) \end{aligned}$$

Parameters

$$\Upsilon = (\mathbf{u}_1, \dots, \mathbf{u}_I, \zeta, \sigma^2, \omega^2),$$

where $\zeta \in \mathbb{R}$ and $\mathbf{u}_i = (u_{i,1}, \dots, u_{i,K}) \in \mathbb{R}^K$.

Hyper-parameters

$$(a_\sigma, b_\sigma, a_\omega, b_\omega).$$

Posterior

$$\begin{aligned} p(\Upsilon \mid \mathbf{Y}) &= p(\mathbf{Y} \mid \zeta, \{\mathbf{u}_i\}) p(\{\mathbf{u}_i\} \mid \sigma^2) p(\sigma^2) p(\zeta \mid \omega^2) p(\omega^2) \\ &\propto \prod_{i=1}^{I-1} \prod_{i'=i+1}^I \theta_{i,i'}^{y_{i,i'}} (1 - \theta_{i,i'})^{1-y_{i,i'}} \times \prod_{i=1}^I (\sigma^2)^{-K/2} \exp\left\{-\frac{1}{2\sigma^2} \|\mathbf{u}_i\|^2\right\} \times (\sigma^2)^{-(a_\sigma+1)} \exp\left\{-\frac{b_\sigma}{\sigma^2}\right\} \\ &\quad \times (\omega^2)^{-1/2} \exp\left\{-\frac{1}{2\omega^2} \zeta^2\right\} \times (\omega^2)^{-(a_\omega+1)} \exp\left\{-\frac{b_\omega}{\omega^2}\right\}, \end{aligned}$$

where $\theta_{i,i'} = \text{expit}(\zeta - \|\mathbf{u}_i - \mathbf{u}_{i'}\|)$.

MCMC Algorithm

The algorithm proceeds by generating a new state $\Upsilon^{(b+1)}$ from a current state $\Upsilon^{(b)}$, $b = 1, \dots, B$, as follows:

1. Sample $\mathbf{u}_i^{(b+1)}$, $i = 1, \dots, I$, according to a Metropolis–Hastings Algorithm, considering the fcd:

$$p(\mathbf{u}_i \mid \text{rest}) \propto \prod_{i'=i+1}^I \theta_{i,i'}^{y_{i,i'}} (1 - \theta_{i,i'})^{1-y_{i,i'}} \times \prod_{i'=1}^{i-1} \theta_{i',i}^{y_{i',i}} (1 - \theta_{i',i})^{1-y_{i',i}} \times \exp\left\{-\frac{1}{2\sigma^2} \|\mathbf{u}_i\|^2\right\}.$$

2. Sample $\zeta^{(b+1)}$ according to a Metropolis–Hastings Algorithm, considering the fd:

$$p(\zeta \mid \text{rest}) \propto \prod_{i=1}^{I-1} \prod_{i'=i+1}^I \theta_{i,i'}^{y_{i,i'}} (1 - \theta_{i,i'})^{1-y_{i,i'}} \times \exp\left\{-\frac{1}{2\omega^2} \zeta^2\right\}.$$

3. Sample $(\sigma^2)^{(b+1)}$ from $p(\sigma^2 \mid \text{rest}) = \text{IGam}\left(\sigma^2 \mid a_\sigma + \frac{IK}{2}, b_\sigma + \frac{1}{2} \sum_{i=1}^I \|\mathbf{u}_i\|^2\right)$.

4. Sample $(\omega^2)^{(b+1)}$ from $p(\omega^2 \mid \text{rest}) = \text{IGam}\left(\omega^2 \mid a_\omega + \frac{1}{2}, b_\omega + \frac{1}{2} \zeta^2\right)$.

Prior Elicitation

Setting a priori $\mathbb{E}[\sigma^2] = V_2(I^{1/K})$ and $\mathbb{E}[\omega^2] = 100$, with $\mathbb{CV}[\sigma^2] = \mathbb{CV}[\omega^2] = \infty$, it follows that

$$a_\sigma = 2, \quad b_\sigma = V_2(I^{1/K}), \quad a_\omega = 2, \quad b_\omega = 100,$$

where $V_2(I^{1/K})$ is the volume of a 2-dimensional Euclidean ball of radius $I^{1/K}$.

4 Class Model (Stochastic Block Model, SBM)

Likelihood

$$y_{i,i'} \mid \xi_i, \xi_{i'}, \{\eta_{k,\ell}\} \stackrel{\text{iid}}{\sim} \text{Ber}(\text{expit } \eta_{\phi(\xi_i, \xi_{i'})})$$

$$\begin{aligned} p(\mathbf{Y} \mid \{\xi_i\}, \{\eta_{k,\ell}\}) &= \prod_{i=1}^{I-1} \prod_{i'=i+1}^I (\text{expit } \eta_{\phi(\xi_i, \xi_{i'})})^{y_{i,i'}} (1 - \text{expit } \eta_{\phi(\xi_i, \xi_{i'})})^{1-y_{i,i'}} \\ &= \prod_{k=1}^K \prod_{\ell=k}^K (\text{expit } \eta_{k,\ell})^{s_{k,\ell}} (1 - \text{expit } \eta_{k,\ell})^{n_{k,\ell} - s_{k,\ell}} \end{aligned}$$

where $s_{k,\ell} = \sum_{\mathcal{S}_{k,\ell}} y_{i,i'}$, $n_{k,\ell} = \sum_{\mathcal{S}_{k,\ell}} 1$, with $\mathcal{S}_{k,\ell} = \{(i, i') : i < i' \text{ and } \phi(\xi_i, \xi_{i'}) = (k, \ell)\}$, and $\phi(x, y) = (\min\{x, y\}, \max\{x, y\})$ is a function to take into account that $\mathbf{Y} = [y_{i,i'}]$ is a symmetric matrix.

Prior

$$\begin{aligned} \eta_{k,\ell} \mid \zeta, \tau^2 &\stackrel{\text{iid}}{\sim} \text{N}(\zeta, \tau^2) \\ \zeta &\sim \text{N}(\mu_\zeta, \sigma_\zeta^2) \\ \tau^2 &\sim \text{IGam}(a_\tau, b_\tau) \\ \xi_i \mid \boldsymbol{\omega} &\stackrel{\text{iid}}{\sim} \text{Cat}(\boldsymbol{\omega}) \end{aligned}$$

$$\begin{aligned}\boldsymbol{\omega} \mid \alpha &\sim \text{Dir}\left(\frac{\alpha}{K}, \dots, \frac{\alpha}{K}\right) \\ \alpha &\sim \text{Gam}(a_\alpha, b_\alpha)\end{aligned}$$

Parameters

$$\boldsymbol{\Upsilon} = (\eta_{1,1}, \eta_{1,2}, \dots, \eta_{K,K}, \xi_1, \dots, \xi_I, \omega_1, \dots, \omega_K, \zeta, \tau^2, \alpha),$$

where $\xi_i \in \{1, \dots, K\}$, $i = 1, \dots, I$, are the cluster assignments ($\xi_i = k$ means that actor i belongs to cluster k), and $\boldsymbol{\omega} = (\omega_1, \dots, \omega_K)$ is a probability vector such that $\mathbb{Pr}[\xi_i = k \mid w_k] = w_k$, $k = 1, \dots, K$.

Hyper-parameters

$$(\mu_\zeta, \sigma_\zeta^2, a_\tau, b_\tau, a_\alpha, b_\alpha).$$

Posterior

$$\begin{aligned}p(\boldsymbol{\Upsilon} \mid \mathbf{Y}) &= p(\mathbf{Y} \mid \{\xi_i\}, \{\eta_{k,\ell}\}) p(\{\eta_{k,\ell}\} \mid \zeta, \tau^2) p(\zeta) p(\tau^2) p(\{\xi_i\} \mid \boldsymbol{\omega}) p(\boldsymbol{\omega} \mid \alpha) p(\alpha) \\ &\propto \prod_{i=1}^{I-1} \prod_{i'=i+1}^I \theta_{i,i'}^{y_{i,i'}} (1 - \theta_{i,i'})^{1-y_{i,i'}} \times \exp\left\{-\frac{1}{2\sigma_\zeta^2}(\zeta - \mu_\zeta)^2\right\} \times (\tau^2)^{-(a_\tau-1)} \exp\left\{-\frac{b_\tau}{\tau^2}\right\} \\ &\quad \times \prod_{k=1}^K \prod_{\ell=k}^K (\tau^2)^{-1/2} \exp\left\{-\frac{1}{2\tau^2}(\eta_{k,\ell} - \zeta)^2\right\} \times \prod_{i=1}^I \prod_{k=1}^K \omega_k^{[\xi_i=k]} \times \frac{\Gamma\left(\frac{\alpha}{K}\right)^K}{\Gamma(\alpha)} \prod_{k=1}^K \omega_k^{\frac{\alpha}{K}-1} \\ &\quad \times \alpha^{a_\alpha-1} \exp\{-b_\alpha \alpha\},\end{aligned}$$

where $\theta_{i,i'} = \text{expit}(\eta_{\phi(\xi_i, \xi_{i'})})$ and $[\cdot]$ is the Iverson bracket.

MCMC

The algorithm proceeds by generating a new state $\boldsymbol{\Upsilon}^{(b+1)}$ from a current state $\boldsymbol{\Upsilon}^{(b)}$, $b = 1, \dots, B$, as follows:

1. Sample $\eta_{k,\ell}^{(b+1)}$, $\ell = k, \dots, K$ and $k = 1, \dots, K$, according to a Metropolis–Hastings Algorithm, considering the fcd:

$$\begin{aligned}\log p(\eta_{k,\ell} \mid \text{rest}) &\propto s_{k,\ell} \log(\text{expit } \eta_{k,\ell}) + (n_{k,\ell} - s_{k,\ell}) \log(1 - \text{expit } \eta_{k,\ell}) - \frac{1}{2\tau^2}(\eta_{k,\ell} - \zeta)^2 \\ &= s_{k,\ell} \eta_{k,\ell} - n_{k,\ell} \log(1 + \exp \eta_{k,\ell}) - \frac{1}{2\tau^2}(\eta_{k,\ell} - \zeta)^2,\end{aligned}$$

where $s_{k,\ell} = \sum_{\mathcal{S}_{k,\ell}} y_{i,i'}$ and $n_{k,\ell} = \sum_{\mathcal{S}_{k,\ell}} 1$, with $\mathcal{S}_{k,\ell} = \{(i, i') : i < i' \text{ and } \phi(\xi_i, \xi_{i'}) = (k, \ell)\}$.

2. Sample $\xi_i^{(b+1)}$, $i = 1, \dots, I$, from a categorical distribution on $\{1, \dots, K\}$, such that:

$$\mathbb{Pr}[\xi_i = k \mid \text{rest}] \propto \omega_k \times \prod_{i'=i+1}^I \eta_{\phi(k, \xi_{i'})}^{y_{i,i'}} (1 - \eta_{\phi(k, \xi_{i'})})^{1-y_{i,i'}} \times \prod_{i'=1}^{i-1} \eta_{\phi(\xi_{i'}, k)}^{y_{i',i}} (1 - \eta_{\phi(\xi_{i'}, k)})^{1-y_{i',i}}.$$

3. Sample $\boldsymbol{\omega}^{(b+1)}$ from $p(\boldsymbol{\omega} \mid \text{rest}) = \text{Dir}(\boldsymbol{\omega} \mid \frac{\alpha}{K} + n_1, \dots, \frac{\alpha}{K} + n_K)$, where n_k is the number of actors in cluster $k \in \{1, \dots, K\}$.

4. Sample $\zeta^{(b+1)}$ from $\mathbf{N}(m, v^2)$, where

$$v^2 = \left(\frac{1}{\sigma_\zeta^2} + \frac{K(K+1)/2}{\tau^2} \right)^{-1} \quad \text{and} \quad m = v^2 \left(\frac{\mu_\zeta}{\sigma_\zeta^2} + \frac{1}{\tau^2} \sum_{k=1}^K \sum_{\ell=k}^K \eta_{k,\ell} \right).$$

5. Sample $(\sigma^2)^{(b+1)}$ from $p(\sigma^2 \mid \text{rest}) = \text{IGam}\left(\sigma^2 \mid a_\tau + \frac{K(K+1)}{4}, b_\tau + \frac{1}{2} \sum_{k=1}^K \sum_{\ell=k}^K (\eta_{k,\ell} - \zeta)^2\right)$.

6. Sample $\alpha^{(b+1)}$ according to a Metropolis–Hastings Algorithm, considering the fcd:

$$\log p(\alpha \mid \text{rest}) \propto \log \Gamma(\alpha) - K \log \Gamma(\alpha/K) + \frac{\alpha}{K} \sum_{k=1}^K \log \omega_k - (a_\beta - 1) \log \alpha - b_\alpha \alpha.$$

Prior Elicitation

$$\mu_\zeta = 0, \quad \sigma_\zeta^2 = 3, \quad a_\tau = 2, \quad b_\tau = 3, \quad a_\alpha = 1, \quad b_\alpha = 1.$$

5 Eigen Model

Likelihood

$$y_{i,i'} \mid \zeta, \mathbf{u}_i, \mathbf{u}_{i'}, \boldsymbol{\Lambda} \stackrel{\text{ind}}{\sim} \text{Ber}(\text{expit}(\zeta + \mathbf{u}_i^T \boldsymbol{\Lambda} \mathbf{u}_{i'}))$$

Prior

$$\begin{aligned} \mathbf{u}_i &\mid \sigma^2 \stackrel{\text{iid}}{\sim} \mathbf{N}(\mathbf{0}, \sigma^2 \mathbf{I}) \\ \sigma^2 &\sim \text{IGam}(a_\sigma, b_\sigma) \\ \lambda_k &\mid \kappa^2 \stackrel{\text{iid}}{\sim} \mathbf{N}(0, \kappa^2) \\ \kappa^2 &\sim \text{IGam}(a_\kappa, b_\kappa) \\ \zeta &\mid \omega^2 \sim \mathbf{N}(0, \omega^2) \\ \omega^2 &\sim \text{IGam}(a_\omega, b_\omega) \end{aligned}$$

Parameters

$$\boldsymbol{\Upsilon} = (\mathbf{u}_1, \dots, \mathbf{u}_I, \lambda_1, \dots, \lambda_K, \zeta, \sigma^2, \kappa^2, \omega^2),$$

where $\zeta \in \mathbb{R}$ and $\mathbf{u}_i = (u_{i,1}, \dots, u_{i,K}) \in \mathbb{R}^K$, and $\boldsymbol{\Lambda} = \text{diag}[\lambda_1, \dots, \lambda_K]$.

Hyper-parameters

$$(a_\sigma, b_\sigma, a_\kappa, b_\kappa, a_\omega, b_\omega).$$

Posterior

$$\begin{aligned} p(\Upsilon \mid \mathbf{Y}) &= p(\mathbf{Y} \mid \zeta, \{\mathbf{u}_i\}, \{\lambda_k\}) p(\{\mathbf{u}_i\} \mid \sigma^2) p(\sigma^2) p(\{\lambda_k\} \mid \kappa^2) p(\kappa^2) p(\zeta \mid \omega^2) p(\omega^2) \\ &\propto \prod_{i=1}^{I-1} \prod_{i'=i+1}^I \theta_{i,i'}^{y_{i,i'}} (1 - \theta_{i,i'})^{1-y_{i,i'}} \times \prod_{i=1}^I (\sigma^2)^{-K/2} \exp\left\{-\frac{1}{2\sigma^2} \|\mathbf{u}_i\|^2\right\} \times (\sigma^2)^{-(a_\sigma+1)} \exp\left\{-\frac{b_\sigma}{\sigma^2}\right\} \\ &\quad \times \prod_{k=1}^K (\kappa^2)^{-1/2} \exp\left\{-\frac{1}{2\kappa^2} \lambda_k^2\right\} \times (\kappa^2)^{-(a_\kappa+1)} \exp\left\{-\frac{b_\kappa}{\kappa^2}\right\} \times (\omega^2)^{-1/2} \exp\left\{-\frac{1}{2\omega^2} \zeta^2\right\} \\ &\quad \times (\omega^2)^{-(a_\omega+1)} \exp\left\{-\frac{b_\omega}{\omega^2}\right\}, \end{aligned}$$

where $\theta_{i,i'} = \text{expit}(\zeta + \mathbf{u}_i \mathbf{\Lambda} \mathbf{u}_{i'})$.

MCMC

The algorithm proceeds by generating a new state $\Upsilon^{(b+1)}$ from a current state $\Upsilon^{(b)}$, $b = 1, \dots, B$, as follows:

1. Sample $\mathbf{u}_i^{(b+1)}$, $i = 1, \dots, I$, according to a Metropolis–Hastings Algorithm, considering the fcd:

$$p(\mathbf{u}_i \mid \text{rest}) \propto \prod_{i'=i+1}^I \theta_{i,i'}^{y_{i,i'}} (1 - \theta_{i,i'})^{1-y_{i,i'}} \times \prod_{i'=1}^{i-1} \theta_{i',i}^{y_{i',i}} (1 - \theta_{i',i})^{1-y_{i',i}} \times \exp\left\{-\frac{1}{2\sigma^2} \|\mathbf{u}_i\|^2\right\}.$$

2. Sample $\lambda_k^{(b+1)}$ according to a Metropolis–Hastings Algorithm, considering the fcd:

$$p(\lambda_k \mid \text{rest}) \propto \prod_{i=1}^{I-1} \prod_{i'=i+1}^I \theta_{i,i'}^{y_{i,i'}} (1 - \theta_{i,i'})^{1-y_{i,i'}} \times \exp\left\{-\frac{1}{2\kappa^2} \lambda_k^2\right\}.$$

3. Sample $\zeta^{(b+1)}$ according to a Metropolis–Hastings Algorithm, considering the fcd:

$$p(\zeta \mid \text{rest}) \propto \prod_{i=1}^{I-1} \prod_{i'=i+1}^I \theta_{i,i'}^{y_{i,i'}} (1 - \theta_{i,i'})^{1-y_{i,i'}} \times \exp\left\{-\frac{1}{\omega^2} \zeta^2\right\}.$$

4. Sample $(\sigma^2)^{(b+1)}$ from $p(\sigma^2 \mid \text{rest}) = \text{IGam}\left(\sigma^2 \mid a_\sigma + \frac{IK}{2}, b_\sigma + \frac{1}{2} \sum_{i=1}^I \|\mathbf{u}_i\|^2\right)$.

5. Sample $(\kappa^2)^{(b+1)}$ from $p(\kappa^2 \mid \text{rest}) = \text{IGam}\left(\kappa^2 \mid a_\kappa + \frac{Q}{2}, b_\kappa + \frac{1}{2} \sum_{k=1}^K \lambda_k^2\right)$.

6. Sample $(\omega^2)^{(b+1)}$ from $p(\omega^2 \mid \text{rest}) = \text{IGam}(\omega^2 \mid a_\sigma + \frac{1}{2}, b_\sigma + \frac{1}{2}\zeta^2)$.

Prior Elicitation

Setting a priori $\mathbb{E}[\sigma^2] = \mathbb{E}[\kappa^2] = \mathbb{E}[\omega^2] = 3$, with $\mathbb{CV}[\sigma^2] = \mathbb{CV}[\kappa^2] = \mathbb{CV}[\omega^2] = \infty$, it follows that

$$a_\sigma = 2, \quad b_\sigma = 3, \quad a_\kappa = 2, \quad b_\kappa = 3, \quad a_\omega = 2, \quad b_\omega = 3.$$

6 Class-Distance Model

Likelihood

$$y_{i,i'} \mid \zeta, \{\mathbf{u}_k\}, \xi_i, \xi_{i'} \stackrel{\text{ind}}{\sim} \text{Ber}(\text{expit} \eta_{\phi(\xi_i, \xi_{i'})})$$

where

$$\eta_{k,\ell} = \zeta - \|\mathbf{u}_k - \mathbf{u}_\ell\|$$

Prior

$$\begin{aligned} \mathbf{u}_k &\mid \sigma^2 \stackrel{\text{iid}}{\sim} \text{N}(\mathbf{0}, \sigma^2 \mathbf{I}) \\ \sigma^2 &\sim \text{IGam}(a_\sigma, b_\sigma) \\ \zeta &\mid \omega^2 \sim \text{N}(0, \omega^2) \\ \omega^2 &\sim \text{IGam}(a_\omega, b_\omega) \\ \xi_i &\mid \boldsymbol{\omega} \stackrel{\text{iid}}{\sim} \text{Cat}(\boldsymbol{\omega}) \\ \boldsymbol{\omega} &\mid \alpha \sim \text{Dir}\left(\frac{\alpha}{K}, \dots, \frac{\alpha}{K}\right) \\ \alpha &\sim \text{Gam}(a_\alpha, b_\alpha) \end{aligned}$$

Parameters

$$\boldsymbol{\Upsilon} = (\mathbf{u}_1, \dots, \mathbf{u}_K, \xi_1, \dots, \xi_I, \omega_1, \dots, \omega_K, \sigma^2, \omega^2, \alpha),$$

where $\zeta \in \mathbb{R}$ and $\mathbf{u}_k = (u_{k,1}, \dots, u_{k,Q}) \in \mathbb{R}^Q$.

Hyper-parameters

$$(a_\sigma, b_\sigma, a_\omega, b_\omega, a_\alpha, b_\alpha).$$

Posterior

$$p(\boldsymbol{\Upsilon} \mid \mathbf{Y}) = p(\mathbf{Y} \mid \zeta, \{\mathbf{u}_k\}, \{\xi_i\}, \{\xi_{i'}\}) p(\{\mathbf{u}_k\} \mid \sigma^2) p(\sigma^2) p(\zeta \mid \omega^2) p(\omega^2) p(\{\xi_i\} \mid \boldsymbol{\omega}) p(\boldsymbol{\omega} \mid \alpha) p(\alpha)$$

$$\begin{aligned}
& \propto \prod_{i=1}^{I-1} \prod_{i'=i+1}^I \theta_{i,i'}^{y_{i,i'}} (1 - \theta_{i,i'})^{1-y_{i,i'}} \times \prod_{k=1}^K (\sigma^2)^{-K/2} \exp\left\{-\frac{1}{2\sigma^2} \|\mathbf{u}_k\|^2\right\} \times (\sigma^2)^{-(a_\sigma+1)} \exp\left\{-\frac{b_\sigma}{\sigma^2}\right\} \\
& \times (\omega^2)^{-1/2} \exp\left\{-\frac{1}{2\omega^2} \zeta^2\right\} \times (\omega^2)^{-(a_\omega+1)} \exp\left\{-\frac{b_\omega}{\omega^2}\right\} \times \prod_{i=1}^I \prod_{k=1}^K \omega_k^{[\xi_i=k]} \times \frac{\Gamma\left(\frac{\alpha}{K}\right)^K}{\Gamma(\alpha)} \prod_{k=1}^K \omega_k^{\frac{\alpha}{K}-1} \\
& \times \alpha^{a_\alpha-1} \exp\{-b_\alpha \alpha\}
\end{aligned}$$

where $\theta_{i,i'} = \text{expit } \eta_{\phi(\xi_i, \xi_{i'})}$ and $[\cdot]$ is the Iverson bracket.

MCMC

The algorithm proceeds by generating a new state $\Upsilon^{(b+1)}$ from a current state $\Upsilon^{(b)}$, $b = 1, \dots, B$, as follows:

1. Sample $\mathbf{u}_k^{(b+1)}$, $k = 1, \dots, K$, according to a Metropolis–Hastings Algorithm, considering the fcd:

$$p(\mathbf{u}_k \mid \text{rest}) \propto \prod_{\substack{i,i': i < i' \\ \xi_i = k \text{ or } \xi_{i'} = k}} \theta_{i,i'}^{y_{i,i'}} (1 - \theta_{i,i'})^{1-y_{i,i'}} \times \exp\left\{-\frac{1}{2\sigma^2} \|\mathbf{u}_k\|^2\right\}.$$

2. Sample $\zeta^{(b+1)}$ according to a Metropolis–Hastings Algorithm, considering the fcd:

$$p(\zeta \mid \text{rest}) \propto \prod_{i=1}^{I-1} \prod_{i'=i+1}^I \theta_{i,i'}^{y_{i,i'}} (1 - \theta_{i,i'})^{1-y_{i,i'}} \times \exp\left\{-\frac{1}{\omega^2} \zeta^2\right\}.$$

3. Sample $\xi_i^{(b+1)}$, $i = 1, \dots, I$, from a categorical distribution on $\{1, \dots, K\}$, such that:

$$\mathbb{P}r[\xi_i = k \mid \text{rest}] \propto \omega_k \times \prod_{i'=i+1}^I \eta_{\phi(k, \xi_{i'})}^{y_{i,i'}} (1 - \eta_{\phi(k, \xi_{i'})})^{1-y_{i,i'}} \times \prod_{i'=1}^{i-1} \eta_{\phi(\xi_{i'}, k)}^{y_{i',i}} (1 - \eta_{\phi(\xi_{i'}, k)})^{1-y_{i',i}}.$$

4. Sample $\boldsymbol{\omega}^{(b+1)}$ from $p(\boldsymbol{\omega} \mid \text{rest}) = \text{Dir}(\boldsymbol{\omega} \mid \frac{\alpha}{K} + n_1, \dots, \frac{\alpha}{K} + n_K)$, where n_k is the number of actors in cluster $k \in \{1, \dots, K\}$.
5. Sample $(\sigma^2)^{(b+1)}$ from $p(\sigma^2 \mid \text{rest}) = \text{IGam}\left(\sigma^2 \mid a_\sigma + \frac{KQ}{2}, b_\sigma + \frac{1}{2} \sum_{k=1}^K \|\mathbf{u}_k\|^2\right)$.
6. Sample $(\omega^2)^{(b+1)}$ from $p(\omega^2 \mid \text{rest}) = \text{IGam}\left(\omega^2 \mid a_\omega + \frac{1}{2}, b_\omega + \frac{1}{2} \zeta^2\right)$.
7. Sample $\alpha^{(b+1)}$ according to a Metropolis–Hastings Algorithm, considering the fcd:

$$\log p(\alpha \mid \text{rest}) \propto \log \Gamma(\alpha) - K \log \Gamma(\alpha/K) + \frac{\alpha}{K} \sum_{k=1}^K \log \omega_k - (a_\alpha - 1) \log \alpha - b_\alpha \alpha.$$

Prior Elicitation Setting a priori $\mathbb{E}[\sigma^2] = V_2(K^{1/Q})$, $\mathbb{E}[\omega^2] = 3$ and $\mathbb{E}[\alpha] = 1$, with $\mathbb{CV}[\sigma^2] = \mathbb{CV}[\omega^2] = \infty$ and $\mathbb{CV}[\alpha] = 1$, it follows that

$$a_\sigma = 2, \quad b_\sigma = V_2(K^{1/Q}), \quad a_\omega = 2, \quad b_\omega = 3, \quad a_\alpha = 1, \quad b_\alpha = 1,$$

where $V_2(K^{1/Q})$ is the volume of a 2-dimensional Euclidean ball of radius $I^{1/Q}$.

7 Class-Eigen Model

Likelihood

$$y_{i,i'} \mid \zeta, \{\mathbf{u}_k\}, \mathbf{\Lambda}, \xi_i, \xi_{i'} \stackrel{\text{ind}}{\sim} \text{Ber}(\text{expit } \eta_{\phi(\xi_i, \xi_{i'})})$$

where

$$\eta_{k,\ell} = \zeta + \mathbf{u}_k^T \mathbf{\Lambda} \mathbf{u}_\ell$$

Prior

$$\begin{aligned} \mathbf{u}_k &\mid \sigma^2 \stackrel{\text{iid}}{\sim} \mathbf{N}(\mathbf{0}, \sigma^2 \mathbf{I}) \\ \sigma^2 &\sim \text{IGam}(a_\sigma, b_\sigma) \\ \lambda_q &\mid \kappa^2 \stackrel{\text{iid}}{\sim} \mathbf{N}(0, \kappa^2) \\ \kappa^2 &\sim \text{IGam}(a_\kappa, b_\kappa) \\ \zeta &\mid \omega^2 \sim \mathbf{N}(0, \omega^2) \\ \omega^2 &\sim \text{IGam}(a_\omega, b_\omega) \\ \xi_i &\mid \omega \stackrel{\text{iid}}{\sim} \text{Cat}(\omega) \\ \omega &\mid \alpha \sim \text{Dir}\left(\frac{\alpha}{K}, \dots, \frac{\alpha}{K}\right) \\ \alpha &\sim \text{Gam}(a_\alpha, b_\alpha) \end{aligned}$$

Parameters

$$\mathbf{\Upsilon} = (\mathbf{u}_1, \dots, \mathbf{u}_K, \lambda_1, \dots, \lambda_Q, \xi_1, \dots, \xi_I, \omega_1, \dots, \omega_K, \sigma^2, \kappa^2, \omega^2, \alpha),$$

where $\zeta \in \mathbb{R}$ and $\mathbf{u}_k = (u_{k,1}, \dots, u_{k,Q}) \in \mathbb{R}^Q$, and $\mathbf{\Lambda} = \text{diag}[\lambda_1, \dots, \lambda_Q]$.

Hyper-parameters

$$(a_\sigma, b_\sigma, a_\kappa, b_\kappa, a_\omega, b_\omega, a_\alpha, b_\alpha).$$

Posterior

$$p(\mathbf{\Upsilon} \mid \mathbf{Y}) = p(\mathbf{Y} \mid \zeta, \{\mathbf{u}_k\}, \{\lambda_q\}, \{\xi_i\}) p(\{\mathbf{u}_k\} \mid \sigma^2) p(\sigma^2) p(\{\lambda_q\} \mid \kappa^2) p(\kappa^2) p(\zeta \mid \omega^2) p(\omega^2)$$

$$\begin{aligned}
& p(\{\xi_i\} \mid \boldsymbol{\omega}) p(\boldsymbol{\omega} \mid \alpha) p(\alpha) \\
& \propto \prod_{i=1}^{I-1} \prod_{i'=i+1}^I \theta_{i,i'}^{y_{i,i'}} (1 - \theta_{i,i'})^{1-y_{i,i'}} \times \prod_{k=1}^K (\sigma^2)^{-K/2} \exp\left\{-\frac{1}{2\sigma^2} \|\mathbf{u}_k\|^2\right\} \times (\sigma^2)^{-(a_\sigma+1)} \exp\left\{-\frac{b_\sigma}{\sigma^2}\right\} \\
& \times \prod_{q=1}^Q (\kappa^2)^{-1/2} \exp\left\{-\frac{1}{2\kappa^2} \lambda_q^2\right\} \times (\kappa^2)^{-(a_\kappa+1)} \exp\left\{-\frac{b_\kappa}{\kappa^2}\right\} \times (\omega^2)^{-1/2} \exp\left\{-\frac{1}{2\omega^2} \zeta^2\right\} \\
& \times (\omega^2)^{-(a_\omega+1)} \exp\left\{-\frac{b_\omega}{\omega^2}\right\} \times \prod_{i=1}^I \prod_{k=1}^K \omega_k^{[\xi_i=k]} \times \frac{\Gamma\left(\frac{\alpha}{K}\right)^K}{\Gamma(\alpha)} \prod_{k=1}^K \omega_k^{\frac{\alpha}{K}-1} \times \alpha^{a_\alpha-1} \exp\{-b_\alpha \alpha\},
\end{aligned}$$

where $\theta_{i,i'} = \text{expit } \eta_{\phi(\xi_i, \xi_{i'})}$ and $[\cdot]$ is the Iverson bracket.

MCMC

The algorithm proceeds by generating a new state $\boldsymbol{\Upsilon}^{(b+1)}$ from a current state $\boldsymbol{\Upsilon}^{(b)}$, $b = 1, \dots, B$, as follows:

1. Sample $\mathbf{u}_k^{(b+1)}$, $k = 1, \dots, K$, according to a Metropolis–Hastings Algorithm, considering the fcd:

$$p(\mathbf{u}_k \mid \text{rest}) \propto \prod_{\substack{i < i' \\ \xi_i = k \text{ or } \xi_{i'} = k}} \theta_{i,i'}^{y_{i,i'}} (1 - \theta_{i,i'})^{1-y_{i,i'}} \times \exp\left\{-\frac{1}{2\sigma^2} \|\mathbf{u}_k\|^2\right\}.$$

2. Sample $\lambda_q^{(b+1)}$, $q = 1, \dots, Q$, according to a Metropolis–Hastings Algorithm, considering the fcd:

$$p(\lambda_q \mid \text{rest}) \propto \prod_{i=1}^{I-1} \prod_{i'=i+1}^I \theta_{i,i'}^{y_{i,i'}} (1 - \theta_{i,i'})^{1-y_{i,i'}} \times \exp\left\{-\frac{1}{2\kappa^2} \lambda_q^2\right\}.$$

3. Sample $\zeta^{(b+1)}$ according to a Metropolis–Hastings Algorithm, considering the fcd:

$$p(\zeta \mid \text{rest}) \propto \prod_{i=1}^{I-1} \prod_{i'=i+1}^I \theta_{i,i'}^{y_{i,i'}} (1 - \theta_{i,i'})^{1-y_{i,i'}} \times \exp\left\{-\frac{1}{\omega^2} \zeta^2\right\}.$$

4. Sample $\xi_i^{(s+1)}$, $i = 1, \dots, I$, from a categorical distribution on $\{1, \dots, K\}$, such that:

$$\mathbb{Pr}[\xi_i = k \mid \text{rest}] \propto \omega_k \times \prod_{i'=i+1}^I \eta_{\phi(k, \xi_{i'})}^{y_{i,i'}} (1 - \eta_{\phi(k, \xi_{i'})})^{1-y_{i,i'}} \times \prod_{i'=1}^{i-1} \eta_{\phi(\xi_{i'}, k)}^{y_{i',i}} (1 - \eta_{\phi(\xi_{i'}, k)})^{1-y_{i',i}}.$$

5. Sample $\boldsymbol{\omega}^{(b+1)}$ from $p(\boldsymbol{\omega} \mid \text{rest}) = \text{Dir}(\boldsymbol{\omega} \mid \frac{\alpha}{K} + n_1, \dots, \frac{\alpha}{K} + n_K)$, where n_k is the number of actors in cluster $k \in \{1, \dots, K\}$.

6. Sample $(\sigma^2)^{(b+1)}$ from $p(\sigma^2 \mid \text{rest}) = \text{IGam}\left(\sigma^2 \mid a_\sigma + \frac{KQ}{2}, b_\sigma + \frac{1}{2} \sum_{k=1}^K \|\mathbf{u}_k\|^2\right)$.
7. Sample $(\kappa^2)^{(b+1)}$ from $p(\kappa^2 \mid \text{rest}) = \text{IGam}\left(\kappa^2 \mid a_\kappa + \frac{Q}{2}, b_\kappa + \frac{1}{2} \sum_{q=1}^Q \lambda_q^2\right)$.
8. Sample $(\omega^2)^{(b+1)}$ from $p(\omega^2 \mid \text{rest}) = \text{IGam}\left(\omega^2 \mid a_\omega + \frac{1}{2}, b_\omega + \frac{1}{2}\zeta^2\right)$.
9. Sample $\alpha^{(b+1)}$ according to a Metropolis–Hastings Algorithm, considering the fcd:

$$\log p(\alpha \mid \text{rest}) \propto \log \Gamma(\alpha) - K \log \Gamma(\alpha/K) + \frac{\alpha}{K} \sum_{k=1}^K \log \omega_k - (a_\alpha - 1) \log \alpha - b_\alpha \alpha.$$

Prior Elicitation

Setting a priori $\mathbb{E}[\sigma^2] = \mathbb{E}[\kappa^2] = \mathbb{E}[\omega^2] = 3$, $\mathbb{E}[\alpha^2] = 1$, with $\mathbb{CV}[\sigma^2] = \mathbb{CV}[\kappa^2] = \mathbb{CV}[\omega^2] = \infty$ and $\mathbb{CV}[\alpha] = 1$, it follows that

$$a_\sigma = 2, \quad b_\sigma = 3, \quad a_\kappa = 2, \quad b_\kappa = 3, \quad a_\omega = 2, \quad b_\omega = 3, \quad a_\alpha = 1, \quad b_\alpha = 1.$$

8 Multilevel–Class Model

Likelihood

$$y_{i,i'} \mid \xi_i, \xi_{i'}, \{\eta_{k,\ell}\} \stackrel{\text{ind}}{\sim} \text{Ber}(\text{expit } \eta_{\phi(\xi_i, \xi_{i'})})$$

$$\begin{aligned} p(\mathbf{Y} \mid \{\xi_i\}, \{\eta_{k,\ell}\}) &= \prod_{i=1}^{I-1} \prod_{i'=i+1}^I (\text{expit } \eta_{\phi(\xi_i, \xi_{i'})})^{y_{i,i'}} (1 - \text{expit } \eta_{\phi(\xi_i, \xi_{i'})})^{1-y_{i,i'}} \\ &= \prod_{k=1}^K \prod_{\ell=k}^K (\text{expit } \eta_{k,\ell})^{s_{k,\ell}} (1 - \text{expit } \eta_{k,\ell})^{n_{k,\ell} - s_{k,\ell}} \end{aligned}$$

where $s_{k,\ell} = \sum_{\mathcal{S}_{k,\ell}} y_{i,i'}$ and $n_{k,\ell} = \sum_{\mathcal{S}_{k,\ell}} 1$, with $\mathcal{S}_{k,\ell} = \{(i, i') : i < i' \text{ and } \phi(\xi_i, \xi_{i'}) = (k, \ell)\}$.

Prior

$$\begin{aligned} \eta_{k,\ell} \mid \gamma_k, \gamma_\ell, \{\mu_{q,r}\}, \sigma^2 &\sim \mathbf{N}(\mu_{\phi(\gamma_k, \gamma_\ell)}, \sigma^2) \\ \mu_{q,r} \mid \zeta, \tau^2 &\stackrel{\text{ind}}{\sim} \mathbf{N}(\zeta, \tau^2) \\ \zeta &\sim \mathbf{N}(\mu_\zeta, \sigma_\zeta^2) \\ \tau^2 &\sim \text{IGam}(a_\tau, b_\tau) \\ \sigma^2 &\sim \text{IGam}(a_\sigma, b_\sigma) \\ \gamma_k &\mid \boldsymbol{\vartheta} \stackrel{\text{iid}}{\sim} \text{Cat}(\boldsymbol{\vartheta}) \end{aligned}$$

$$\begin{aligned}
\boldsymbol{\vartheta} \mid \boldsymbol{\beta} &\sim \text{Dir}\left(\frac{\beta}{Q}, \dots, \frac{\beta}{Q}\right) \\
\boldsymbol{\beta} &\sim \text{Gam}(a_\beta, b_\beta) \\
\xi_i \mid \boldsymbol{\omega} &\stackrel{\text{iid}}{\sim} \text{Cat}(\boldsymbol{\omega}) \\
\boldsymbol{\omega} \mid \boldsymbol{\alpha} &\sim \text{Dir}\left(\frac{\alpha}{K}, \dots, \frac{\alpha}{K}\right) \\
\boldsymbol{\alpha} &\sim \text{Gam}(a_\alpha, b_\alpha)
\end{aligned}$$

Parameters

$$\boldsymbol{\Upsilon} = (\eta_{1,1}, \eta_{1,2}, \dots, \eta_{K,K}, \mu_{1,1}, \mu_{1,2}, \dots, \mu_{Q,Q}, \zeta, \tau^2, \sigma^2, \gamma_1, \dots, \gamma_K, \vartheta_1, \dots, \vartheta_Q, \beta, \xi_1, \dots, \xi_I, \omega_1, \dots, \omega_K, \alpha),$$

where $\boldsymbol{\xi} = (\xi_1, \dots, \xi_I)$, $\xi_i \in \{1, \dots, K\}$, are the actors-cluster assignments ($\xi_i = k$ means that actor i belongs to cluster k), $\boldsymbol{\gamma} = (\gamma_1, \dots, \gamma_K)$, $\gamma_k \in \{1, \dots, Q\}$, are the super-cluster assignments ($\gamma_k = q$ means that cluster k belongs to super-cluster q), $\phi(x, y) = (\min\{x, y\}, \max\{x, y\})$ is a function to take into account that both $\mathbf{Y} = [y_{i,i'}]$ and $\boldsymbol{\eta} = [\eta_{k,\ell}]$ are symmetric matrices, and $\boldsymbol{\omega} = (\omega_1, \dots, \omega_K)$ and $\boldsymbol{\vartheta} = (\vartheta_1, \dots, \vartheta_Q)$ are probability vectors such that $\Pr[\xi_i = k \mid \omega_k] = \omega_k$ and $\Pr[\gamma_k = q \mid \vartheta_q] = \vartheta_q$, respectively.

Hyper-parameters

$$(\mu_\zeta, \sigma_\zeta^2, a_\tau, b_\tau, a_\sigma, b_\sigma, a_\beta, b_\beta, a_\alpha, b_\alpha).$$

Posterior

$$\begin{aligned}
p(\boldsymbol{\Upsilon} \mid \mathbf{Y}) &\propto \prod_{i=1}^{I-1} \prod_{i'=i+1}^I \theta_{i,i'}^{y_{i,i'}} (1 - \theta_{i,i'})^{1-y_{i,i'}} \times \prod_{k=1}^K \prod_{\ell=k}^K (\sigma^2)^{-1/2} \exp\left\{-\frac{1}{2\sigma^2} (\eta_{k,\ell} - \mu_{\phi(\gamma_k, \gamma_\ell)})^2\right\} \\
&\times \prod_{q=1}^Q \prod_{r=q}^Q (\tau^2)^{-\frac{1}{2}} \exp\left\{-\frac{1}{2\tau^2} (\mu_{q,r} - \zeta)^2\right\} \times \exp\left\{-\frac{1}{2\sigma_\zeta^2} (\zeta - \mu_\zeta)^2\right\} \times (\tau^2)^{-(a_\tau-1)} \exp\left\{-\frac{b_\tau}{\tau^2}\right\} \\
&\times (\sigma^2)^{-(a_\sigma-1)} \exp\left\{-\frac{b_\sigma}{\sigma^2}\right\} \times \prod_{k=1}^K \prod_{q=1}^Q \vartheta_q^{[\gamma_k=q]} \times \frac{\Gamma\left(\frac{\beta}{Q}\right)^Q}{\Gamma(\beta)} \prod_{q=1}^Q \vartheta_q^{\frac{\beta}{Q}-1} \times \beta^{a_\beta-1} \exp\{-b_\beta \beta\} \\
&\times \prod_{i=1}^I \prod_{k=1}^K \omega_k^{[\xi_i=k]} \times \frac{\Gamma\left(\frac{\alpha}{K}\right)^K}{\Gamma(\alpha)} \prod_{k=1}^K \omega_k^{\frac{\alpha}{K}-1} \times \alpha^{a_\alpha-1} \exp\{-b_\alpha \alpha\},
\end{aligned}$$

where $\theta_{i,i'} = \text{expit} \eta_{\phi(\xi_i, \xi_{i'})}$ and $[\cdot]$ is the Iverson bracket.

MCMC

The algorithm proceeds by generating a new state $\boldsymbol{\Upsilon}^{(b+1)}$ from a current state $\boldsymbol{\Upsilon}^{(b)}$, $b = 1, \dots, B$, as follows:

1. Sample $\eta_{k,\ell}^{(b+1)}$ according to a Metropolis–Hastings Algorithm, considering the fd:

$$\begin{aligned}\log p(\eta_{k,\ell} \mid \text{rest}) &\propto s_{k,\ell} \log(\text{expit } \eta_{k,\ell}) + (n_{k,\ell} - s_{k,\ell}) \log(1 - \text{expit } \eta_{k,\ell}) - \frac{1}{2\sigma^2} (\eta_{k,\ell} - \mu_{\phi(\gamma_k, \gamma_\ell)})^2 \\ &= s_{k,\ell} \eta_{k,\ell} - n_{k,\ell} \log(1 + \exp \eta_{k,\ell}) - \frac{1}{2\sigma^2} (\eta_{k,\ell} - \mu_{\phi(\gamma_k, \gamma_\ell)})^2,\end{aligned}$$

where $s_{k,\ell} = \sum_{\mathcal{S}_{k,\ell}} y_{i,i'}$ and $n_{k,\ell} = \sum_{\mathcal{S}_{k,\ell}} 1$, with $\mathcal{S}_{k,\ell} = \{(i, i') : i < i' \text{ and } \phi(\xi_i, \xi_{i'}) = (k, \ell)\}$.

2. Sample $\mu_{q,r}^{(b+1)}$ from $\mathbf{N}(m, v^2)$, where

$$v^2 = \left(\frac{1}{\tau^2} + \frac{n_{q,r}}{\sigma^2} \right)^{-1} \quad \text{and} \quad m = v^2 \left(\frac{\zeta}{\tau^2} + \frac{s_{q,r}}{\sigma^2} \right),$$

where $s_{q,r} = \sum_{\mathcal{S}_{q,r}} \eta_{k,\ell}$ and $n_{q,r} = \sum_{\mathcal{S}_{q,r}} 1$, with $\mathcal{S}_{q,r} = \{(k, \ell) : k \leq \ell \text{ and } \phi(\gamma_k, \gamma_\ell) = (q, r)\}$.

3. Sample $\zeta^{(b+1)}$ from $\mathbf{N}(m, v^2)$, where

$$v^2 = \left(\frac{1}{\sigma_\zeta^2} + \frac{n_Q}{\tau^2} \right)^{-1} \quad \text{and} \quad m = v^2 \left(\frac{\mu_\zeta}{\sigma_\zeta^2} + \frac{\mu_{..}}{\tau^2} \right),$$

where $n_Q = Q(Q+1)/2$ and $\mu_{..} = \sum_{q=1}^Q \sum_{r=q}^Q \mu_{q,r}$.

4. Sample $(\tau^2)^{(b+1)}$ from $\text{IGam}(a, b)$, where

$$a = a_\tau + n_Q/2 \quad \text{and} \quad b = b_\tau + \frac{1}{2} \sum_{q=1}^Q \sum_{r=q}^Q (\mu_{q,r} - \zeta)^2,$$

where $n_Q = Q(Q+1)/2$.

5. Sample $(\sigma^2)^{(b+1)}$ from $\text{IGam}(a, b)$, where

$$a = a_\sigma + n_K/2 \quad \text{and} \quad b = b_\sigma + \frac{1}{2} \sum_{k=1}^K \sum_{\ell=k}^K (\eta_{k,\ell} - \mu_{\phi(\gamma_k, \gamma_\ell)})^2,$$

where $n_K = K(K+1)/2$.

6. Sample $\gamma_k^{(b+1)}$ from a categorical distribution on $\{1, \dots, Q\}$, such that:

$$\mathbb{P}r[\gamma_k = q \mid \text{rest}] \propto \vartheta_q \times \prod_{\ell=k}^K \mathbf{N}(\eta_{k,\ell} \mid \mu_{\phi(q, \gamma_\ell)}, \sigma^2) \times \prod_{\ell=1}^{k-1} \mathbf{N}(\eta_{\ell,k} \mid \mu_{\phi(q, \gamma_\ell)}, \sigma^2),$$

for $q \in \{1, \dots, Q\}$.

7. Sample $\boldsymbol{\vartheta}^{(b+1)}$ from $p(\boldsymbol{\vartheta} \mid \text{rest}) = \text{Dir}\left(\boldsymbol{\vartheta} \mid \frac{\beta}{Q} + n_1, \dots, \frac{\beta}{Q} + n_Q\right)$, where n_q is the number of clusters in super-cluster $q \in \{1, \dots, Q\}$.

8. Sample $\beta^{(b+1)}$ according to a Metropolis–Hastings Algorithm, considering the fcd:

$$p(\beta \mid \text{rest}) \propto \log \Gamma(\beta) - K \log \Gamma(\beta/Q) + \frac{\beta}{Q} \sum_{q=1}^Q \log \vartheta_q - (a_\beta - 1) \log \beta - b_\beta \beta.$$

9. Sample $\xi_i^{(b+1)}$ from a categorical distribution on $\{1, \dots, K\}$, such that:

$$\mathbb{Pr} [\xi_i = k \mid \text{rest}] \propto \omega_k \times \prod_{i'=i+1}^I \eta_{\phi(k, \xi_{i'})}^{y_{i, i'}} (1 - \eta_{\phi(k, \xi_{i'})})^{1-y_{i, i'}} \times \prod_{i'=1}^{i-1} \eta_{\phi(k, \xi_{i'})}^{y_{i', i}} (1 - \eta_{\phi(k, \xi_{i'})})^{1-y_{i', i}}.$$

for $k \in \{1, \dots, K\}$.

10. Sample $\omega^{(b+1)}$ from $p(\omega \mid \text{rest}) = \text{Dir}(\omega \mid \frac{\alpha}{K} + n_1, \dots, \frac{\alpha}{K} + n_K)$, where n_k is the number of actors in cluster $k \in \{1, \dots, K\}$.

11. Sample $\alpha^{(b+1)}$ according to a Metropolis–Hastings Algorithm, considering the fcd:

$$p(\alpha \mid \text{rest}) \propto \log \Gamma(\alpha) - K \log \Gamma(\alpha/K) + \frac{\alpha}{K} \sum_{k=1}^K \log \omega_k - (a_\alpha - 1) \log \alpha - b_\alpha \alpha.$$

Prior Elicitation

$$\mu_\zeta = 0, \quad \sigma_\zeta^2 = 3, \quad a_\tau = 2, \quad b_\tau = 3, \quad a_\sigma = 2, \quad b_\sigma = 3, \quad a_\beta = 1, \quad b_\beta = 3, \quad a_\alpha = 1, \quad b_\alpha = 1.$$

9 WAIC

$$\text{lppd} = \sum_{i=1}^n \log \left(\frac{1}{B} \sum_{b=1}^B p(y_i \mid \theta^b) \right)$$

$$p_{\text{WAIC}_1} = 2 \sum_{i=1}^n \left[\log \left(\frac{1}{B} \sum_{b=1}^B p(y_i \mid \theta^b) \right) - \frac{1}{B} \sum_{b=1}^B \log p(y_i \mid \theta^b) \right]$$

$$p_{\text{WAIC}_2} = \frac{1}{B-1} \sum_{i=1}^n \sum_{b=1}^B (a_{i,b} - \bar{a}_i)^2, \quad \bar{a}_i = \frac{1}{B} \sum_{b=1}^B a_{i,b}, \quad a_{i,b} = \log p(y_i \mid \theta^b)$$

$$\text{WAIC} = -2 \text{lppd} + 2 p_{\text{WAIC}}$$