Examination: Histopathological examination

Patient: XXX	PESEL: XXX	Age:	Gender:		
Material: Total organ resection – segment of the large intestine Unit in charge:					
Physician in charge:					
Material collected on:	Material received	on:			
Expected time of examination:					
Clinical diagnosis: Tumour of the hepatic flexure.					
Examination performed	l on:				

Macroscopic description:

An 18 cm length of the large intestine with a piece of the mesentery sized 20 x 7 x 5 cm, with a 7.5 cm length of the small intestine, and a fragment of the omentum sized 11 x 8 x l,5. A cauliflower-shaped, ulcerous tumour sized 5.5 x 4 x 2.2 cm found in the mucosa. The lesion surrounds 100% of the intestine circumference, narrowing its lumen, is located 10.5 cm from the proximal incision line, 7 cm from the distal incision line, and 3 cm from the Bauhnin's valve. Minimum side margin 6.5 cm. Microscopic description:

Adenocarcinoma tubulopapillare partim mucinosum (G3).

Infiltratio carcinomatosa tuncae muscularis propriae et telae adiposae pericolicae.

Incision lines clear of neoplastic lesions.

Metastases carcinomatosae in lymphonodis (No III/VI). Infiltratio carcinomatosa telae adiposae perinodalis. Lymphonodulitis reactiva No XVIII).

Adjacent omentum fragment free of cancer.

Histopathological diagnosis:

Adenocarcinoma tubulopapillare partim mucinosum coli. Tubulopapillar and partially mucinous adenocarcinoma of the colon. (G3, Dukes B, Astler-Coller B2, pT3, pNO).

Compliance validated by:

CONTACT YOUR DOCTOR WITH THIS REPORT!