Block 2

Weeks 5–8 · 3 sessions a week · pyramid style

When: 3 sessions a week, with a rest day between (e.g. Mon, Wed, Fri)

How long: 15–20 minutes, plus warm-up and cool-down

Effort: 6–7 out of 10

Each session

1. **Warm up, 5 minutes.** Gentle mobility and marching on the spot.
2. **Main workout, 15–20 minutes.** The exercises below, in this order.
3. **Cool down, 5 minutes.** Walk it out and stretch lightly.

Your workout

> Do every exercise for 12 reps, then every exercise for 11 reps, and keep counting down to 1. Move straight down the pyramid and rest as needed between exercises.

1	**Sit-to-stand** Feet flat, lean forward slightly, stand tall, then sit back down with control.	12 down to 1
2	**Incline push-up using a chair** Hands on the chair seat, body in one line, chest to the edge.	12 down to 1 (lower incline)
3	**Glute bridges** Drive through your heels, lift hips, lower slowly.	12 down to 1
4	**Dead bug (legs only)** Lower back pressed to the floor, lower one leg at a time slowly.	12 down to 1
5	**Bird dog (legs only)** On hands and knees, extend one leg back, hips level, back flat.	12 down to 1

Rest: Move straight down the pyramid, rest as needed between exercises.

Watch the demo for each exercise before your first session. The link is in your email.

Walking this block

Walk 3–4 times a week for 20–25 minutes, plus a 10-minute walk after one meal each day.

Walking between your sessions makes a real difference to blood sugar, especially a short walk after a meal.

How hard should it feel?

Rate your effort from 1 (very easy) to 10 (flat out). Aim for **6–7** this block.

1–2	**3–4**	**5–6**	**7–8**	**9–10**
Very easy, could go all day	Easy, chatting freely	Working, can still talk	Hard, short sentences only	Very hard to flat out

Too easy? Slow each rep down or add a round. Too hard? Use the easier option shown in the demo and rest a little longer.

Tick off your sessions

	Session 1	Session 2	Session 3	Notes (effort, how you felt)
Week 5				
Week 6				
Week 7				
Week 8				

Stay safe. Stop any exercise that causes sharp pain (normal muscle tiredness is fine). If you take medicine that lowers blood glucose, such as insulin or a sulfonylurea, keep a fast-acting carbohydrate (like jelly beans or juice) nearby while you exercise, and know the signs of a low: shakiness, sweating, dizziness, confusion. If you're unsure whether an exercise is right for you, check with your doctor or nurse first.