Block 1

Weeks 1–4 · 3 sessions a week · circuit style

When: 3 sessions a week, with a rest day between (e.g. Mon, Wed, Fri)

How long: 15–20 minutes, plus warm-up and cool-down

Effort: 5–6 out of 10

Each session

1. **Warm up, 5 minutes.** Gentle mobility and marching on the spot.
2. **Main workout, 15–20 minutes.** The exercises below, in this order.
3. **Cool down, 5 minutes.** Walk it out and stretch lightly.

Your workout

> Do the 5 exercises in order. Rest 1 minute after the last one. That's one round. Repeat rounds until your 15–20 minutes are up.

1	**Bodyweight squat** Sit back like into a chair, chest up, knees tracking over toes.	12 reps (hold a chair if needed)
2	**Push-up** Body in one straight line, lower chest towards the floor, push away.	12 reps (on knees if needed)
3	**Bird dog** Reach opposite arm and leg long, keep hips square to the floor.	16 reps (8 each side)
4	**Marching glute bridge** Hold hips high and level as you lift one foot at a time.	16 reps (8 each side)
5	**Forearm plank** Elbows under shoulders, body straight, squeeze glutes and brace your tummy.	30 seconds

Rest: Rest 1 minute after all 5 exercises.

Watch the demo for each exercise before your first session. The link is in your email.

Walking this block

Walk briskly 4 times a week for 25 minutes (add a hill or incline if you can), plus two short walks after meals each day.

Walking between your sessions makes a real difference to blood sugar, especially a short walk after a meal.

How hard should it feel?

Rate your effort from 1 (very easy) to 10 (flat out). Aim for **5–6** this block.

1–2	**3–4**	**5–6**	**7–8**	**9–10**
Very easy, could go all day	Easy, chatting freely	Working, can still talk	Hard, short sentences only	Very hard to flat out

Too easy? Slow each rep down or add a round. Too hard? Use the easier option shown in the demo and rest a little longer.

Tick off your sessions

	Session 1	Session 2	Session 3	Notes (effort, how you felt)
Week 1				
Week 2				
Week 3				
Week 4				

Stay safe. Stop any exercise that causes sharp pain (normal muscle tiredness is fine). If you take medicine that lowers blood glucose, such as insulin or a sulfonylurea, keep a fast-acting carbohydrate (like jelly beans or juice) nearby while you exercise, and know the signs of a low: shakiness, sweating, dizziness, confusion. If you're unsure whether an exercise is right for you, check with your doctor or nurse first.