Block 2

Weeks 5–8 · 3 sessions a week · pyramid style

When: 3 sessions a week, with a rest day between (e.g. Mon, Wed, Fri)

How long: 15–20 minutes, plus warm-up and cool-down

Effort: 6–7 out of 10

Each session

1. **Warm up, 5 minutes.** Gentle mobility and marching on the spot.
2. **Main workout, 15–20 minutes.** The exercises below, in this order.
3. **Cool down, 5 minutes.** Walk it out and stretch lightly.

Your workout

> Do every exercise for 10 reps, then every exercise for 9 reps, and keep counting down to 1. Move straight down the pyramid.

1	**Air squat with calf raise** Squat down, then stand and rise onto your toes in one smooth motion.	10 down to 1 (one fluid motion)
2	**Inchworm into push-up** Walk hands out to a plank, do a push-up, walk back.	10 down to 1 (one fluid motion)
3	**Marching glute bridge** Hold hips high and level as you lift one foot at a time.	10 down to 1 each side
4	**Bird dog, knee and elbow touch** Extend opposite arm and leg, then draw elbow and knee together underneath.	10 down to 1 each side
5	**Plank step-outs** Hold a strong plank, step one foot out and back, hips steady.	10 down to 1 each side

Rest: Move straight down the pyramid.

Watch the demo for each exercise before your first session. The link is in your email.

Walking this block

Twice a week, walk intervals: 3 minutes brisk, 1 minute fast, 5 times (about 20 minutes). Steady walks on other days.

Walking between your sessions makes a real difference to blood sugar, especially a short walk after a meal.

How hard should it feel?

Rate your effort from 1 (very easy) to 10 (flat out). Aim for **6–7** this block.

1–2 Very easy, could go all day	**3–4** Easy, chatting freely	**5–6** Working, can still talk	**7–8** Hard, short sentences only	**9–10** Very hard to flat out

Too easy? Slow each rep down or add a round. Too hard? Use the easier option shown in the demo and rest a little longer.

Tick off your sessions

	Session 1	Session 2	Session 3	Notes (effort, how you felt)
Week 5				
Week 6				
Week 7				
Week 8				

Stay safe. Stop any exercise that causes sharp pain (normal muscle tiredness is fine). If you take medicine that lowers blood glucose, such as insulin or a sulfonylurea, keep a fast-acting carbohydrate (like jelly beans or juice) nearby while you exercise, and know the signs of a low: shakiness, sweating, dizziness, confusion. If you're unsure whether an exercise is right for you, check with your doctor or nurse first.