Block 3

Weeks 9–12 · 3 sessions a week · circuit style

When: 3 sessions a week, with a rest day between (e.g. Mon, Wed, Fri)

How long: 15–20 minutes, plus warm-up and cool-down

Effort: 7–8 out of 10

Each session

1. **Warm up, 5 minutes.** Gentle mobility and marching on the spot.
2. **Main workout, 15–20 minutes.** The exercises below, in this order.
3. **Cool down, 5 minutes.** Walk it out and stretch lightly.

Your workout

> Do the 4 exercises in order, 30 seconds each. Rest 1 minute after the last one. That's one round. Repeat rounds until your 15–20 minutes are up.

1	**Reverse lunges** Step back, lower the back knee towards the floor, front knee over ankle.	30 seconds
2	**Push-up with knee taps** Do a push-up, then draw one knee towards the elbow each side.	30 seconds
3	**Plank with steps in and out** From a plank, step feet in towards hands and back out, hips steady.	30 seconds (one fluid motion)
4	**Bird dog knee hover (legs only)** On hands and knees, lift knees just off the floor, hold still.	30 seconds

Rest: Rest 1 minute after all 4 exercises.

Watch the demo for each exercise before your first session. The link is in your email.

Walking this block

2–3 times a week, do a 25–30 minute interval walk. Walk every day, even if only 10 minutes after a meal.

Walking between your sessions makes a real difference to blood sugar, especially a short walk after a meal.

How hard should it feel?

Rate your effort from 1 (very easy) to 10 (flat out). Aim for **7–8** this block.

1–2	**3–4**	**5–6**	**7–8**	**9–10**
Very easy, could go all day	Easy, chatting freely	Working, can still talk	Hard, short sentences only	Very hard to flat out

Too easy? Slow each rep down or add a round. Too hard? Use the easier option shown in the demo and rest a little longer.

Tick off your sessions

	Session 1	Session 2	Session 3	Notes (effort, how you felt)
Week 9				
Week 10				
Week 11				
Week 12				

Stay safe. Stop any exercise that causes sharp pain (normal muscle tiredness is fine). If you take medicine that lowers blood glucose, such as insulin or a sulfonylurea, keep a fast-acting carbohydrate (like jelly beans or juice) nearby while you exercise, and know the signs of a low: shakiness, sweating, dizziness, confusion. If you're unsure whether an exercise is right for you, check with your doctor or nurse first.