Block 2

Weeks 5–8 · 3 sessions a week · pyramid style

When: 3 sessions a week, with a rest day between (e.g. Mon, Wed, Fri)

How long: 15–20 minutes, plus warm-up and cool-down

Effort: 7 out of 10

Each session

1. **Warm up, 5 minutes.** Gentle mobility and marching on the spot.
2. **Main workout, 15–20 minutes.** The exercises below, in this order.
3. **Cool down, 5 minutes.** Walk it out and stretch lightly.

Your workout

> Do every exercise for 5 reps, then every exercise for 4 reps, and keep counting down to 1. Rest 1 minute after each full pyramid.

1	**Bodyweight squat** Sit back like into a chair, chest up, knees tracking over toes.	5 down to 1
2	**Single-leg RDL** Hinge at the hips, back flat, reach down as your back leg lifts.	5 down to 1 each side
3	**Curtsy lunges** Step one leg behind and across, bend both knees, chest tall.	5 down to 1
4	**Inchworm with 2 push-ups** Walk hands out to a plank, do two push-ups, walk back.	5 down to 1
5	**Sit-ups** Feet planted, curl up with control, lower back down slowly.	5 down to 1
6	**Leg raises** Lower back pressed down, lift straight legs up, lower slowly.	5 down to 1

Rest: Rest 1 minute after each completed pyramid.

Watch the demo for each exercise before your first session. The link is in your email.

Walking this block

2–3 times a week, do a 20–25 minute interval walk (hard and easy bursts). Steady walks on other days.

Walking between your sessions makes a real difference to blood sugar, especially a short walk after a meal.

How hard should it feel?

Rate your effort from 1 (very easy) to 10 (flat out). Aim for **7** this block.

1–2	**3–4**	**5–6**	**7–8**	**9–10**
Very easy, could go all day	Easy, chatting freely	Working, can still talk	Hard, short sentences only	Very hard to flat out

Too easy? Slow each rep down or add a round. Too hard? Use the easier option shown in the demo and rest a little longer.

Tick off your sessions

	Session 1	Session 2	Session 3	Notes (effort, how you felt)
Week 5				
Week 6				
Week 7				
Week 8				

Stay safe. Stop any exercise that causes sharp pain (normal muscle tiredness is fine). If you take medicine that lowers blood glucose, such as insulin or a sulfonylurea, keep a fast-acting carbohydrate (like jelly beans or juice) nearby while you exercise, and know the signs of a low: shakiness, sweating, dizziness, confusion. If you're unsure whether an exercise is right for you, check with your doctor or nurse first.