Block 3

Weeks 9–12 · 3 sessions a week · circuit style

When: 3 sessions a week, with a rest day between (e.g. Mon, Wed, Fri)

How long: 15–20 minutes, plus warm-up and cool-down

Effort: 7 out of 10

Each session

1. **Warm up, 5 minutes.** Gentle mobility and marching on the spot.
2. **Main workout, 15–20 minutes.** The exercises below, in this order.
3. **Cool down, 5 minutes.** Walk it out and stretch lightly.

Your workout

> Do the 6 exercises in order. Rest 1 minute after the last one. That's one round. Repeat rounds until your 15–20 minutes are up.

1	**Bodyweight squat** Sit back like into a chair, chest up, knees tracking over toes.	12 reps (chair-assisted if needed)
2	**Floor push-up on knees** Hands under shoulders, straight line from knees to head, lower chest down.	10 reps
3	**Marching glute bridge** Hold hips high and level as you lift one foot at a time.	16 reps
4	**Hollow hold** Lower back pressed down, arms and legs reaching long, ribs tucked in.	20 seconds
5	**Ankle taps** Shoulders slightly lifted, reach side to side to tap each heel.	20 reps (10 each side)
6	**Bird dog** Reach opposite arm and leg long, keep hips square to the floor.	12 reps

Rest: Rest 1 minute after all 6 exercises.

Watch the demo for each exercise before your first session. The link is in your email.

Walking this block

Walk 4 times a week for 25–30 minutes, plus a 10-minute walk after meals each day.

Walking between your sessions makes a real difference to blood sugar, especially a short walk after a meal.

How hard should it feel?

Rate your effort from 1 (very easy) to 10 (flat out). Aim for **7** this block.

1–2	**3–4**	**5–6**	**7–8**	**9–10**
Very easy, could go all day	Easy, chatting freely	Working, can still talk	Hard, short sentences only	Very hard to flat out

Too easy? Slow each rep down or add a round. Too hard? Use the easier option shown in the demo and rest a little longer.

Tick off your sessions

	Session 1	Session 2	Session 3	Notes (effort, how you felt)
Week 9				
Week 10				
Week 11				
Week 12				

Stay safe. Stop any exercise that causes sharp pain (normal muscle tiredness is fine). If you take medicine that lowers blood glucose, such as insulin or a sulfonylurea, keep a fast-acting carbohydrate (like jelly beans or juice) nearby while you exercise, and know the signs of a low: shakiness, sweating, dizziness, confusion. If you're unsure whether an exercise is right for you, check with your doctor or nurse first.