



## Section 1.1

# 1 **GRADED** [2 points]



[Correct drawing of vector field + 1 point] Regardless of the initial condition, the solution will tend towards  $y = 3/2$ . [Long-term behavior + 1 point]

# 3 **GRADED** [2 points]



[Correct drawing of vector field + 1 point] Regardless of the initial condition, the solution tend toward  $\infty$ . [Long-term behavior + 1 point]

# 22 **GRADED** [6 points]

Let  $V(t)$  be the volume of the raindrop and  $S(t)$  the surface area. The question tells us that

$$\frac{d}{dt}V(t) = -\gamma S(t) \quad [+1 \text{ points}] \quad (1)$$

for some number  $\gamma$  (the rate per area). If  $R$  is the radius, the formula for the volume of a sphere is  $V(t) = \frac{4}{3}\pi R^3$  and  $S(t) = 4\pi R^2$  [These formulas +2 points] it follows that

$$S(t) = 4\pi \left( \frac{3}{4\pi} \right)^{2/3} V^{2/3} \quad [+1 \text{ points}] \quad (2)$$

hence

$$\frac{d}{dt}V(t) = -\mu V^{2/3} \quad [+2 \text{ points}] \quad (3)$$

for some constant  $\mu$ .

## Section 1.2

### # 7 GRADED [9 points]

(a) The solution is

$$p(t) = 900 + ce^{t/2} \quad [\text{identify solution +2 points}] \quad (4)$$

To find  $c$  we set  $p(0) = 850$ :

$$850 = 900 + c \implies c = -50 \quad [\text{solve for } c +1 \text{ point}] \quad (5)$$

To find the time the population goes extinct, we set  $p(t_{\text{ex}}) = 0$  and solve for  $t_{\text{ex}}$ :

$$900 - 50e^{t_{\text{ex}}/2} = 0 \implies e^{t_{\text{ex}}/2} = \frac{900}{50} = 18 \quad (6)$$

$$\implies t_{\text{ex}} = 2 \ln 18 \quad [\text{solve for } t_{\text{ex}} +2 \text{ point}] \quad (7)$$

(b) The same idea as (a): First find  $c$

$$p_0 = 900 + c \implies c = p_0 - 900 \quad (8)$$

and find  $t_{\text{ex}}$

$$900 - (p_0 - 900)e^{t_{\text{ex}}/2} = 0 \implies e^{t_{\text{ex}}/2} = \frac{900}{p_0 - 900} \quad (9)$$

$$\implies t_{\text{ex}} = 2 \ln \frac{900}{p_0 - 900} \quad [\text{repeat (a) with general } p_0 +2 \text{ point}] \quad (10)$$

(c) If  $t_{\text{ex}} = 1$ ,

$$t_{\text{ex}} = 1 = 2 \ln \frac{900}{p_0 - 900} \implies \frac{900}{p_0 - 900} = e^{1/2} \quad (11)$$

$$\implies p_0 - 900 = \frac{900}{e^{1/2}} \quad (12)$$

$$\implies p_0 = 900 + \frac{900}{\ln 2} = 900 \left( 1 + \frac{1}{e^{1/2}} \right) \quad [+2 \text{ point}] \quad (13)$$

### # 10

### # 12 GRADED

(a) Solving the differential equation for  $Q$  yields exponential decay

$$Q = Q(0)e^{-rt} \quad (14)$$

For this problem  $Q(0) = 100\text{mg}$  and at  $t = 1$  (assuming we measure  $t$  in weeks)  $Q(t) = 82.04$ . Plugging these values in to the solutions yields an equation for  $r$ :

$$82.04 = 100e^{-r} \implies r = \ln \frac{100}{82.04} \approx 1.97$$

# 17

**Section 1.3**# 11 **GRADED [6 points]**Starting with  $y_1$ , the derivatives are

$$y' = \frac{1}{2}t^{1/2-1} = \frac{1}{2t^{1/2}} \quad [+1 \text{ point}] \quad (15)$$

$$y'' = -\frac{1}{2} \frac{1}{2t^{1/2+1}} = -\frac{1}{4t^{3/2}} \quad [+1 \text{ point}] \quad (16)$$

Hence

$$-2t^2y'' + 3ty' - y = 2\frac{t^2}{4t^{3/2}} + \frac{3t}{2t^{1/2}} - y \quad (17)$$

$$= -\frac{1}{2}t^{1/2} + \frac{3}{2}t^{1/2} - t^{1/2} = t^{1/2} - t^{1/2} \quad [+1 \text{ point}] \quad (18)$$

For  $t_2$  we have

$$y' = -t^{-2} \quad [+1 \text{ point}] \quad (19)$$

$$y'' = 2t^{-3} \quad [+1 \text{ point}] \quad (20)$$

hence

$$-4t^{-1} - 3t^{-1} - t^{-1} = 0 \quad [+1 \text{ point}] \quad (21)$$

**Additional Questions**