

Lecture 2 (9/16)

Independence, Marginalization, and Conditioning

Example

$P(C, G)$	$G = M$	$G = F$
$C = 1$ (diagnosed)	0.0021	0.0019
$C = 0$ (not diagnosed)	0.4899	0.5061

Example

$P(C, G)$	$G = M$	$G = F$
$C = 1$ (diagnosed)	0.0021	0.0019
$C = 0$ (not diagnosed)	0.4899	0.5061

For example, if these numbers are obtained from a population of N people,

$$0.0021 = \frac{N(\{\text{Diagnosed \& Male}\})}{N}, \quad N(E) = \# \text{ samples in event } E$$

Example

$P(C, G)$	$G = M$	$G = F$
$C = 1$ (diagnosed)	0.0021	0.0019
$C = 0$ (not diagnosed)	0.4899	0.5061

For example, if these numbers are obtained from a population of N people,

$$0.0021 = \frac{N(\{\text{Diagnosed \& Male}\})}{N}, \quad N(E) = \# \text{ samples in event } E$$

What do we care about?

- Hospital management: What is the chance a random person (M or F) is diagnosed? (marginalization)
- Dr and patient: What is the chance my patient (known to be M or F) will be diagnosed? (conditional probability)

Marginalization

If we don't care about M/F distinction:

$$\begin{aligned}P(C = 1) &= P(C = 1, G = M) + P(C = 1, G = F) \\&= 0.0021 + 0.0019 = 0.004\end{aligned}$$

Marginalization

If we don't care about M/F distinction:

$$\begin{aligned}P(C = 1) &= P(C = 1, G = M) + P(C = 1, G = F) \\&= 0.0021 + 0.0019 = 0.004\end{aligned}$$

We used that probability of disjoint events as: **(Countable) additivity**: for any countable collection (meaning we can list them as $E_1, E_2, \dots$) of pairwise disjoint sets $E_1, E_2, \dots$,

$$P\left(\bigcup_i E_i\right) = \sum_i P(E_i).$$

Marginalization

If we don't care about M/F distinction:

$$\begin{aligned}P(C = 1) &= P(C = 1, G = M) + P(C = 1, G = F) \\&= 0.0021 + 0.0019 = 0.004\end{aligned}$$

We used that probability of disjoint events as: **(Countable) additivity**: for any countable collection (meaning we can list them as $E_1, E_2, \dots$) of pairwise disjoint sets $E_1, E_2, \dots$,

$$P\left(\bigcup_i E_i\right) = \sum_i P(E_i).$$

Marginalization: Sum out the variable you don't care about:

$$P_X(x) = \sum_y P(x, y).$$

Independence

Random variables X, Y are **independent** if

$$P_{X,Y}(x, y) = P_X(x) P_Y(y) \quad \text{for all } x, y.$$

Independence

Random variables X, Y are **independent** if

$$P_{X,Y}(x, y) = P_X(x) P_Y(y) \quad \text{for all } x, y.$$

Example

$X, Y \in \{0, 1\}$ have joint distribution

$$P(X = x, Y = y) = \begin{cases} 1/2 & (0, 0) \\ 1/8 & (0, 1) \\ 1/8 & (1, 0) \\ 1/4 & (1, 1) \end{cases}$$

Are they independent?

Example

$X, Y \in \{0, 1\}$ have joint distribution

$$P(X = x, Y = y) = \begin{cases} 1/2 & (0, 0) \\ 1/8 & (0, 1) \\ 1/8 & (1, 0) \\ 1/4 & (1, 1) \end{cases}$$

Are they independent?

Example

$X, Y \in \{0, 1\}$ have joint distribution

$P(X, Y)$	$Y = 0$	$Y = 1$
$X = 0$	1/8	3/8
$X = 1$	1/8	3/8

Are they independent?

How many checks for a 2×2 table?

Let X and Y have (marginal) distributions

$$X \sim \text{Bernoulli}(p)$$

$$Y \sim \text{Bernoulli}(q)$$

How many checks for a 2×2 table?

Let X and Y have (marginal) distributions

$$X \sim \text{Bernoulli}(p)$$

$$Y \sim \text{Bernoulli}(q)$$

Suppose

$$P(X = 1, Y = 1) = pq.$$

Does this mean X and Y are independent?

How many checks for a 2×2 table?

Let X and Y have (marginal) distributions

$$X \sim \text{Bernoulli}(p)$$

$$Y \sim \text{Bernoulli}(q)$$

Suppose

$$P(X = 1, Y = 1) = pq.$$

Does this mean X and Y are independent?

The other three cells are not free: rows and columns must sum to the marginals, so

$$\begin{aligned} P(X = 1, Y = 0) + P(X = 1, Y = 1) &= P(X=1) = p \\ \implies P(X = 1, Y = 0) &= p - P(X = 1, Y = 1) \\ &= p(1 - q) = P(X = 1)P(Y = 0) \end{aligned}$$

Same for the other entries. In general: for variables X and Y with n and m outcomes respectively, we need to check $(m - 1)(n - 1)$ entries.

Example

$X \in \{0, 1\}$ and $Y \in \{1, 2, 3, 4\}$ have joint distribution

$P(X, Y)$	$Y = 1$	$Y = 2$	$Y = 3$	$Y = 4$
$X = 0$	1/16	1/16	1/16	1/16
$X = 1$	3/16	3/16	3/16	3/16

Are X and Y independent?

Example

$X \in \{0, 1\}$ and $Y \in \{1, 2, 3, 4\}$ have joint distribution

$P(X, Y)$	$Y = 1$	$Y = 2$	$Y = 3$	$Y = 4$
$X = 0$	1/16	1/16	1/16	1/16
$X = 1$	3/16	3/16	3/16	3/16

Are X and Y independent?

Example

$P(X, Y)$	$Y = 1$	$Y = 2$	$Y = 3$	$Y = 4$
$X = 0$	1/4	1/8	1/8	0
$X = 1$	0	1/8	1/8	1/4

Are X and Y independent?

Cancer diagnosis data: compare the joint probabilities to the products of the marginals:

(C, G)	Joint $P(C, G)$	Product $P(C)P(G)$
$(1, M)$	0.0021	0.001968
$(1, F)$	0.0019	0.002032
$(0, M)$	0.4899	0.490032
$(0, F)$	0.5061	0.505968

Cancer diagnosis data: compare the joint probabilities to the products of the marginals:

(C, G)	Joint $P(C, G)$	Product $P(C)P(G)$
$(1, M)$	0.0021	0.001968
$(1, F)$	0.0019	0.002032
$(0, M)$	0.4899	0.490032
$(0, F)$	0.5061	0.505968

- Are they independent? Or could it be that the difference is due to having a small number of samples?

Cancer diagnosis data: compare the joint probabilities to the products of the marginals:

(C, G)	Joint $P(C, G)$	Product $P(C)P(G)$
$(1, M)$	0.0021	0.001968
$(1, F)$	0.0019	0.002032
$(0, M)$	0.4899	0.490032
$(0, F)$	0.5061	0.505968

- Are they independent? Or could it be that the difference is due to having a small number of samples?
- One way to approach: collect more data and see if the numbers get closer.

Cancer diagnosis data: compare the joint probabilities to the products of the marginals:

(C, G)	Joint $P(C, G)$	Product $P(C)P(G)$
$(1, M)$	0.0021	0.001968
$(1, F)$	0.0019	0.002032
$(0, M)$	0.4899	0.490032
$(0, F)$	0.5061	0.505968

- Are they independent? Or could it be that the difference is due to having a small number of samples?
- One way to approach: collect more data and see if the numbers get closer.
- If we can't do this, we need to quantify statistical uncertainty (Unit 3)

Cancer diagnosis data: compare the joint probabilities to the products of the marginals:

(C, G)	Joint $P(C, G)$	Product $P(C)P(G)$
$(1, M)$	0.0021	0.001968
$(1, F)$	0.0019	0.002032
$(0, M)$	0.4899	0.490032
$(0, F)$	0.5061	0.505968

- Are they independent? Or could it be that the difference is due to having a small number of samples?
- One way to approach: collect more data and see if the numbers get closer.
- If we can't do this, we need to quantify statistical uncertainty (Unit 3)
- Are men more or less likely to be diagnosed?

Conditioning

The conditional probability of X given Y is

$$P(X = x \mid Y = y) = \frac{P(X = x, Y = y)}{P(Y = y)}$$

Conditioning

The conditional probability of X given Y is

$$P(X = x \mid Y = y) = \frac{P(X = x, Y = y)}{P(Y = y)}$$

Think of it in terms of proportions:

$P(x \mid y)$ is the fraction of times x is observed among outcomes for which y is observed.

Conditioning

The conditional probability of X given Y is


$$P(X = x \mid Y = y) = \frac{P(X = x, Y = y)}{P(Y = y)}$$

Think of it in terms of proportions:


$P(x \mid y)$ is the fraction of times x is observed among outcomes for which y is observed.

- Marginalize over X means: I no longer see X
- Conditioning on X : I fix X to a certain value

Joint, marginal, and conditional probability: a population




Joint, marginal, and conditional probability: a population



A **joint probability**: brown eyes *and* sad.

$$P(\text{brown, sad}) = \frac{4}{24} = \frac{1}{6}.$$


Joint, marginal, and conditional probability: a population



Marginalizing over eye color.

$$P(\text{sad}) = \frac{8}{24} = \frac{1}{3}.$$


Joint, marginal, and conditional probability: a population




Conditioning on brown eyes.

$$P(\text{sad} \mid \text{brown}) = \frac{4}{16} = \frac{1}{4}.$$

Marginalizing vs. conditioning: a tree




Marginalizing vs. conditioning: a tree



Marginalizing over eye color: lump the two highlighted branches together.

$$P(\text{sad}) = \underbrace{\frac{4}{24}}_{\text{sad, brown}} + \underbrace{\frac{4}{24}}_{\text{sad, blue}} = \frac{8}{24} = \frac{1}{3}.$$

Marginalizing vs. conditioning: a tree



Conditioning on brown eyes: forget the blue-eyed branch, read off proportions within what's left.

$$P(\text{sad} | \text{brown}) = \frac{4}{16} = \frac{1}{4}.$$

Example

$X, Y \in \{0, 1\}$ have joint distribution

$$P(X = x, Y = y) = \begin{cases} 3/8 & (0, 0) \\ 1/8 & (0, 1) \\ 1/4 & (1, 0) \\ 1/4 & (1, 1) \end{cases}$$

Find $P(Y = 1 \mid X = 0)$.

Example

$X, Y \in \{0, 1\}$ have joint distribution

$$P(X = x, Y = y) = \begin{cases} 3/8 & (0, 0) \\ 1/8 & (0, 1) \\ 1/4 & (1, 0) \\ 1/4 & (1, 1) \end{cases}$$

Find $P(Y = 1 \mid X = 0)$.

Example

Now suppose $X \in \{0, 1\}$ and $Y \in \{1, 2, 3, 4\}$ have joint distribution

$P(X, Y)$	$Y = 1$	$Y = 2$	$Y = 3$	$Y = 4$
$X = 0$	$1/8$	$1/8$	$1/4$	$1/8$
$X = 1$	$1/8$	0	$1/8$	$1/8$

Find $P(X = 1 \mid Y = 3)$.

Example

Cancer diagnosis data again

$C \mid G$	Marginal $P(C)$	Conditional $P(C \mid G)$
$1 \mid M$	0.0040	0.00427
$1 \mid F$	0.0040	0.00374
$0 \mid M$	0.9960	0.9957
$0 \mid F$	0.9960	0.9963

Example

Cancer diagnosis data again

$C \mid G$	Marginal $P(C)$	Conditional $P(C \mid G)$
$1 \mid M$	0.0040	0.00427
$1 \mid F$	0.0040	0.00374
$0 \mid M$	0.9960	0.9957
$0 \mid F$	0.9960	0.9963

- Are they independent? Same problem as marginal table.

Example

Cancer diagnosis data again

$C \mid G$	Marginal $P(C)$	Conditional $P(C \mid G)$
$1 \mid M$	0.0040	0.00427
$1 \mid F$	0.0040	0.00374
$0 \mid M$	0.9960	0.9957
$0 \mid F$	0.9960	0.9963

- Are they independent? Same problem as marginal table.
- But it is much easier to see the dependence:

$$\text{effect of being male} = P(1|M) - P(1|F) = 0.00427 - 0.00374 = 0.00053$$

Taste of regression models

- Regression models¹ are essentially models given to us as conditional probability models.

¹Some people use regression models more specifically to refer to models with real valued output

Taste of regression models

- Regression models¹ are essentially models given to us as conditional probability models.
- Thus it is natural to consider models given to us in conditional form: For a pair of Bernoullis (X, Y) , instead of defining a joint distribution, we can write

$$\begin{aligned} X &\sim \text{Bernoulli}(q_X) \\ Y|X &\sim \text{Bernoulli}(q_1X + q_0(1 - X)) \quad q_1, q_0 \in [0, 1] \end{aligned}$$

¹Some people use regression models more specifically to refer to models with real valued output

Taste of regression models

- Regression models¹ are essentially models given to us as conditional probability models.
- Thus it is natural to consider models given to us in conditional form: For a pair of Bernoullis (X, Y) , instead of defining a joint distribution, we can write

$$\begin{aligned} X &\sim \text{Bernoulli}(q_X) \\ Y|X &\sim \text{Bernoulli}(q_1X + q_0(1 - X)) \quad q_1, q_0 \in [0, 1] \end{aligned}$$

- $Y|X$ means the distribution of Y conditioned on a value of X

¹Some people use regression models more specifically to refer to models with real valued output

Taste of regression models

- Regression models¹ are essentially models given to us as conditional probability models.
- Thus it is natural to consider models given to us in conditional form: For a pair of Bernoullis (X, Y) , instead of defining a joint distribution, we can write

$$\begin{aligned} X &\sim \text{Bernoulli}(q_X) \\ Y|X &\sim \text{Bernoulli}(q_1X + q_0(1 - X)) \quad q_1, q_0 \in [0, 1] \end{aligned}$$

- $Y|X$ means the distribution of Y conditioned on a value of X
- In general, a regression model takes the form

$$Y|X \sim \text{Name}(\text{Parameters})$$

where now “Parameters” depends on X (if they are not independent).

¹Some people use regression models more specifically to refer to models with real valued output

Example

From the cancer diagnosis data we saw (coding $G = 1$ for male, $G = 0$ for female)

$$q_1 = P(C = 1 \mid G = 1) = 0.00427, \quad q_0 = P(C = 1 \mid G = 0) = 0.00374.$$

Example

From the cancer diagnosis data we saw (coding $G = 1$ for male, $G = 0$ for female)

$$q_1 = P(C = 1 \mid G = 1) = 0.00427, \quad q_0 = P(C = 1 \mid G = 0) = 0.00374.$$

- $0.00053 = q_1 - q_0$ is the “effect” of being male on the diagnosis probability.

Example

From the cancer diagnosis data we saw (coding $G = 1$ for male, $G = 0$ for female)

$$q_1 = P(C = 1 \mid G = 1) = 0.00427, \quad q_0 = P(C = 1 \mid G = 0) = 0.00374.$$

- $0.00053 = q_1 - q_0$ is the “effect” of being male on the diagnosis probability.

Example

Consider

$$P(X = x, Y = y) = \begin{cases} 1/2 & (0, 0) \\ 1/8 & (0, 1) \\ 1/8 & (1, 0) \\ 1/4 & (1, 1) \end{cases}$$

Write a conditional model for $Y|X$.

Example

$$X \sim \text{Bernoulli}(1/3)$$
$$Y|X \sim \text{Bernoulli}(X/2 + 1/3)$$

Find all the joint probabilities and the marginal of Y .

Condition on an even involving a sequence

Example

We are betting on a series of events, each modeled as independent Bernoulli variable $X_i \sim \text{Bernoulli}(q)$ for $i = 1, 2, 3, \dots$. Let^a

$$K = \min\{i : X_i = 0\}$$

- Find $P(K = 4)$
- Find $P(K = 10 | K > 6)$.

Condition on an even involving a sequence

Example

We are betting on a series of events, each modeled as independent Bernoulli variable $X_i \sim \text{Bernoulli}(q)$ for $i = 1, 2, 3, \dots$. Let^a

$$K = \min\{i : X_i = 0\}$$

- Find $P(K = 4)$
- Find $P(K = 10 | K > 6)$.

$$P(K = 4) = P(X_1 = X_2 = X_3 = 1, X_4 = 0) = q^3(1 - q)$$

$$P(K = 10 | K > 6) = \frac{P(K = 10)}{P(K > 6)} = \frac{q^9(1 - q)}{q^6} = q^3(1 - q)$$

^aWe say $K \sim \text{Geometric}(q)$, but you don't need to remember this.

A Markov Chain: conditioning on a history

Example

$$X_0 \sim \text{Bernoulli}(1/2)$$

$$X_{t+1} \mid (X_t, X_{t-1}, \dots, X_0) \sim \text{Bernoulli}(q_1 X_t + q_0(1 - X_t))$$

and assume $X_0 = 1$. This says $X_{t+1} \mid X_t$ is independent of $X_{t-1}, \dots, X_0$, since those don't appear in the Bernoulli random variable.

A Markov Chain: conditioning on a history

Example

$$X_0 \sim \text{Bernoulli}(1/2)$$

$$X_{t+1} \mid (X_t, X_{t-1}, \dots, X_0) \sim \text{Bernoulli}(q_1 X_t + q_0(1 - X_t))$$

and assume $X_0 = 1$. This says $X_{t+1} \mid X_t$ is independent of $X_{t-1}, \dots, X_0$, since those don't appear in the Bernoulli random variable.

Let $p_t = P(X_t = 1)$.

A Markov Chain: conditioning on a history

Example

$$X_0 \sim \text{Bernoulli}(1/2)$$

$$X_{t+1} \mid (X_t, X_{t-1}, \dots, X_0) \sim \text{Bernoulli}(q_1 X_t + q_0(1 - X_t))$$

and assume $X_0 = 1$. This says $X_{t+1} \mid X_t$ is independent of $X_{t-1}, \dots, X_0$, since those don't appear in the Bernoulli random variable.

Let $p_t = P(X_t = 1)$. Then

$$\begin{aligned} p_t &= P(X_t = 1 \mid X_{t-1} = 1)P(X_{t-1} = 1) + P(X_t = 1 \mid X_{t-1} = 0)P(X_{t-1} = 0) \\ &= q_1 p_{t-1} + q_0(1 - p_{t-1}) = (q_1 - q_0)p_{t-1} + q_0. \end{aligned}$$

Steady-state: The probabilities no longer change when $p_t = p_{t-1}$,

$$p_t = \frac{q_0}{1 - q_1 + q_0}.$$

Bayes' formula

Bayes' formula relates the two conditional probabilities for a pair of random variables:

$$P(x | y) = \frac{P(y | x) P(x)}{P(y)}.$$

Bayes' formula

Bayes' formula relates the two conditional probabilities for a pair of random variables:

$$P(x | y) = \frac{P(y | x) P(x)}{P(y)}.$$

Proof: By definition of conditional probability,

$$P(x | y) = \frac{P(x, y)}{P(y)} \quad \text{and} \quad P(y | x) = \frac{P(x, y)}{P(x)}.$$

Bayes' formula

Bayes' formula relates the two conditional probabilities for a pair of random variables:

$$P(x | y) = \frac{P(y | x) P(x)}{P(y)}.$$

Proof: By definition of conditional probability,

$$P(x | y) = \frac{P(x, y)}{P(y)} \quad \text{and} \quad P(y | x) = \frac{P(x, y)}{P(x)}.$$

The second equation gives $P(x, y) = P(y | x) P(x)$. Substituting this into the first,

$$P(x | y) = \frac{P(x, y)}{P(y)} = \frac{P(y | x) P(x)}{P(y)}. \quad \blacksquare$$

Example

A disease affects 1% of a population. A test is positive 95% of the time if you have the disease, and 10% of the time if you don't. Find $P(D \mid +)$.

Example

It rains on 20% of days. When it rains, the forecast correctly predicts rain 90% of the time. When it doesn't rain, the forecast incorrectly predicts rain 15% of the time. Find $P(\text{rain} \mid \text{forecast rain})$.

Answer key

- $\{1/2, 1/8, 1/8, 1/4\}$: independent? **No**
- 2×2 table $\{1/8, 3/8, 1/8, 3/8\}$: independent? **Yes**
- 2×4 table (1/16s and 3/16s): independent? **Yes**
- 2×4 table (with zeros): independent? **No**
- $P(Y=1 \mid X=0)$ for $\{3/8, 1/8, 1/4, 1/4\}$: **1/4**
- $P(X=1 \mid Y=3)$ for the 2×4 conditional example: **1/3**
- Conditional model for $\{1/2, 1/8, 1/8, 1/4\}$: $Y \mid X=0 \sim \text{Bernoulli}(1/5)$,
 $Y \mid X=1 \sim \text{Bernoulli}(2/3)$
- $X \sim \text{Bernoulli}(1/3)$, $Y \mid X \sim \text{Bernoulli}(X/2 + 1/3)$: joint = $(\frac{4}{9}, \frac{2}{9}, \frac{1}{18}, \frac{5}{18})$ for
 $(0, 0), (0, 1), (1, 0), (1, 1)$; marginal $P(Y=1) = \mathbf{1/2}$
- Disease test: $P(D \mid +) \approx \mathbf{0.0876}$
- Weather forecast: $P(\text{rain} \mid \text{forecast rain}) = \mathbf{0.6}$