# Lecture 25 (11/13)

*Additional topics and practice*

**Today:**

1) A practice problem on.
2) Double descent: what happens past the interpolation threshold?
3) Bayesian inference and the relationship to regularization.
4) The kernel trick

## Question

Consider

$$Y = \beta_1 X + \epsilon, \quad \beta_0 = 0 \text{ and } E[X] = 0$$

so the true regression function is $f(x) = E[Y|X = x]$. Treat the $X_i$ as fixed. Recall that the least squared estimate $\hat{\beta}_1$ has standard error $\text{se}(\hat{\beta}_1) = \sigma_\epsilon / \sqrt{N\hat{\sigma}_X^2}$.

(a) What are the bias and the variance of $\hat{\beta}_1$?

(b) Compute $\text{MSE}_{\hat{y}(3,D)} = E_D\left[(\hat{y}(3, D) - f(3))^2\right]$. What is $E\left[(\hat{y}(3, D) - Y)^2\right]$?

(c) What is the expected squared error $E\left[(\hat{y}(3, D) - Y)^2\right]$ for a new point at $x = 3$?

# Solution

The prediction is $\hat{y}(x, D) = \hat{\beta}_1 x$.
(a) Least squares is unbiased, $E[\hat{\beta}_1] = \beta_1$, so the bias is 0. The variance is the square of the standard error:

$$\text{var}(\hat{\beta}_1) = \frac{\sigma_\epsilon^2}{N\hat{\sigma}_X^2}.$$

# Solution

The prediction is $\hat{y}(x, D) = \hat{\beta}_1 x$.
(a) Least squares is unbiased, $E[\hat{\beta}_1] = \beta_1$, so the bias is 0. The variance is the square of the standard error:

$$\text{var}(\hat{\beta}_1) = \frac{\sigma_\epsilon^2}{N\hat{\sigma}_X^2}.$$

(b) $\hat{y}(3, D) - f(3) = 3\hat{\beta}_1 - 3\beta_1 = 3(\hat{\beta}_1 - \beta_1)$, so

$$\text{MSE}_{\hat{y}(3,D)} = E_D\left[9(\hat{\beta}_1 - \beta_1)^2\right] = 9\,\text{var}(\hat{\beta}_1) = \frac{9\sigma_\epsilon^2}{N\hat{\sigma}_X^2}.$$

(The bias is 0, so the MSE is all variance.)

# Solution

The prediction is $\hat{y}(x, D) = \hat{\beta}_1 x$.
(a) Least squares is unbiased, $E[\hat{\beta}_1] = \beta_1$, so the bias is 0. The variance is the square of the standard error:

$$\text{var}(\hat{\beta}_1) = \frac{\sigma_\epsilon^2}{N\hat{\sigma}_X^2}.$$

(b) $\hat{y}(3, D) - f(3) = 3\hat{\beta}_1 - 3\beta_1 = 3(\hat{\beta}_1 - \beta_1)$, so

$$\text{MSE}_{\hat{y}(3,D)} = E_D\left[9(\hat{\beta}_1 - \beta_1)^2\right] = 9\,\text{var}(\hat{\beta}_1) = \frac{9\sigma_\epsilon^2}{N\hat{\sigma}_X^2}.$$

(The bias is 0, so the MSE is all variance.)
(c) A new point is $Y = f(3) + \epsilon$, with $\epsilon$ independent of the training set $D$, so the cross term has mean 0:

$$E\left[(\hat{y}(3, D) - Y)^2\right] = E_D\left[(\hat{y}(3, D) - f(3))^2\right] + E[\epsilon^2] = \frac{9\sigma_\epsilon^2}{N\hat{\sigma}_X^2} + \sigma_\epsilon^2.$$

PNAS

# Reconciling modern machine-learning practice and the classical bias–variance trade-off

Mikhail Belkin, Daniel Hsu, Siyuan Ma, and Soumik Mandal Authors Info & Affiliations

Edited by Peter J. Bickel, University of California, Berkeley, CA, and approved July 2, 2019 (received for review February 21, 2019)

**July 24, 2019** | 116 (32) 15849-15854 | https://doi.org/10.1073/pnas.1903070116

*"While breakthroughs in machine learning and artificial intelligence are changing society, our fundamental understanding has lagged behind. It is traditionally believed that fitting models to the training data exactly is to be avoided as it leads to poor performance on unseen data. However, powerful modern classifiers frequently have near-perfect fit in training. . ."*

PNAS


## Reconciling modern machine-learning practice and the classical bias–variance trade-off


Mikhail Belkin, Daniel Hsu, Siyuan Ma, and Soumik Mandal Authors Info & Affiliations

Edited by Peter J. Bickel, University of California, Berkeley, CA, and approved July 2, 2019 (received for review February 21, 2019)

**July 24, 2019** | 116 (32) 15849-15854 | https://doi.org/10.1073/pnas.1903070116


*"While breakthroughs in machine learning and artificial intelligence are changing society, our fundamental understanding has lagged behind. It is traditionally believed that fitting models to the training data exactly is to be avoided as it leads to poor performance on unseen data. However, powerful modern classifiers frequently have near-perfect fit in training. . ."*




# Double descent in polynomial regression

Consider polynomial regression with degree $K$.




# Double descent in polynomial regression

Consider polynomial regression with degree $K$.




# Double descent in polynomial regression

Consider polynomial regression with degree $K$.




# Double descent in polynomial regression

Consider polynomial regression with degree $K$.




# Double descent in polynomial regression


Consider polynomial regression with degree $K$.




# Double descent in polynomial regression

Consider polynomial regression with degree $K$.






$K = 10$
$K = 19$
$K = 40$
$y$
$x$
4
2
0
−2
−4
0.0
0.5
1.0
fits to new samples of $Y$
true $f(x)$
fit to the data
data

# Double descent in a Fourier (sin/cos) model

Consider a Fourier model with $K$ features.



# Double descent in a Fourier (sin/cos) model



# Bayesian inference: The idea

- So far, we have thought of parameters as a fixed quantities which are estimated from data. The sample distribution provides the uncertainty in these estimates. **The estimator is thought of as a random variable, but NOT the parameter itself.**


# Bayesian inference: The idea

- So far, we have thought of parameters as a fixed quantities which are estimated from data. The sample distribution provides the uncertainty in these estimates. **The estimator is thought of as a random variable, but NOT the parameter itself.**
- In Bayesian inference, the idea is to treat parameters themselves as random variables *before* fitting the data, and then update this distribution based on the data.

# Bayesian inference: The idea

- So far, we have thought of parameters as a fixed quantities which are estimated from data. The sample distribution provides the uncertainty in these estimates. **The estimator is thought of as a random variable, but NOT the parameter itself.**
- In Bayesian inference, the idea is to treat parameters themselves as random variables *before* fitting the data, and then update this distribution based on the data.
- The distribution of a parameter before we fit the data is a our prior.

# Bayesian inference



# Bayesian: Simple example

$$\mu \sim \text{Normal}(0, \tau^2) \quad \text{(prior)}$$
$$Y \mid \mu \sim \text{Normal}(\mu, \sigma_\epsilon^2) \quad \text{(likelihood)}$$

# Bayesian: Simple example

$$\mu \sim \text{Normal}(0, \tau^2) \qquad \text{(prior)}$$
$$Y \mid \mu \sim \text{Normal}(\mu, \sigma_\epsilon^2) \qquad \text{(likelihood)}$$

## Question

Assuming $\sigma_\epsilon^2$ is fixed (meaning we don't model it with priors), what is the posterior of $\mu$?

# Finding the posterior for the mean

Both $\mu$ and $Y$ have mean 0, and

$$\text{var}(Y) = \tau^2 + \sigma_\epsilon^2, \qquad \text{cov}(\mu, Y) = \text{cov}(\mu, \mu + \epsilon) = \tau^2.$$

# Finding the posterior for the mean

Both $\mu$ and $Y$ have mean 0, and

$$\text{var}(Y) = \tau^2 + \sigma_\epsilon^2, \qquad \text{cov}(\mu, Y) = \text{cov}(\mu, \mu + \epsilon) = \tau^2.$$

Regress $\mu$ (response) on $Y$ (predictor). The slope is

$$\frac{\text{cov}(\mu, Y)}{\text{var}(Y)} = \frac{\tau^2}{\tau^2 + \sigma_\epsilon^2}, \qquad \text{and} \qquad \rho^2 = \frac{\text{cov}(\mu, Y)^2}{\text{var}(\mu)\text{var}(Y)} = \frac{\tau^2}{\tau^2 + \sigma_\epsilon^2}.$$

# Finding the posterior for the mean

Both $\mu$ and $Y$ have mean 0, and

$$\text{var}(Y) = \tau^2 + \sigma_\epsilon^2, \qquad \text{cov}(\mu, Y) = \text{cov}(\mu, \mu + \epsilon) = \tau^2.$$

Regress $\mu$ (response) on $Y$ (predictor). The slope is

$$\frac{\text{cov}(\mu, Y)}{\text{var}(Y)} = \frac{\tau^2}{\tau^2 + \sigma_\epsilon^2}, \qquad \text{and} \qquad \rho^2 = \frac{\text{cov}(\mu, Y)^2}{\text{var}(\mu)\text{var}(Y)} = \frac{\tau^2}{\tau^2 + \sigma_\epsilon^2}.$$

The noise variance is $\text{var}(\mu)(1 - \rho^2)$, and $(\mu, Y)$ are jointly Normal, so

$$\mu \mid Y \sim \text{Normal}\Big(\frac{\tau^2}{\tau^2 + \sigma_\epsilon^2}\, Y,\ \frac{\tau^2 \sigma_\epsilon^2}{\tau^2 + \sigma_\epsilon^2}\Big).$$

# Bayesian: Many observations

$$\mu \sim \text{Normal}(0, \tau^2) \qquad \text{(prior)}$$

$$Y_1, \ldots, Y_N \mid \mu \sim \text{Normal}(\mu, \sigma_\epsilon^2) \text{ independent} \qquad \text{(likelihood)}$$

# Bayesian: Many observations

$$\mu \sim \text{Normal}(0, \tau^2) \qquad \text{(prior)}$$

$$Y_1, \ldots, Y_N \mid \mu \sim \text{Normal}(\mu, \sigma_\epsilon^2) \text{ independent} \qquad \text{(likelihood)}$$

## Question

What is the posterior of $\mu$ given $Y_1, \ldots, Y_N$?

# Finding the posterior with many observations

The posterior depends on the data only through $\overline{Y}$, and $\overline{Y} \mid \mu \sim \text{Normal}(\mu, \sigma_\epsilon^2/N)$.

# Finding the posterior with many observations

The posterior depends on the data only through $\overline{Y}$, and $\overline{Y} \mid \mu \sim \mathrm{Normal}(\mu, \sigma_\epsilon^2/N)$. So $\overline{Y}$ plays the role of $Y$, with $\sigma_\epsilon^2/N$ in place of $\sigma_\epsilon^2$:

$$\mathrm{var}(\overline{Y}) = \tau^2 + \frac{\sigma_\epsilon^2}{N}, \qquad \mathrm{cov}(\mu, \overline{Y}) = \tau^2.$$

# Finding the posterior with many observations

The posterior depends on the data only through $\overline{Y}$, and $\overline{Y} \mid \mu \sim \text{Normal}(\mu, \sigma_\epsilon^2/N)$. So $\overline{Y}$ plays the role of $Y$, with $\sigma_\epsilon^2/N$ in place of $\sigma_\epsilon^2$:


$$\text{var}(\overline{Y}) = \tau^2 + \frac{\sigma_\epsilon^2}{N}, \qquad \text{cov}(\mu, \overline{Y}) = \tau^2.$$

Regressing $\mu$ (response) on $\overline{Y}$ (predictor) as before,

$$\mu \mid Y_1, \ldots, Y_N \sim \text{Normal}\Big(\frac{\tau^2}{\tau^2 + \sigma_\epsilon^2/N}\,\overline{Y},\ \frac{\tau^2 \sigma_\epsilon^2/N}{\tau^2 + \sigma_\epsilon^2/N}\Big).$$


# Prior, sample and posterior

$\sigma_\epsilon = 1.$




# Prior, sample and posterior

$\sigma_\epsilon = 1.$




# Prior, sample and posterior

$\sigma_\epsilon = 1.$




# Prior, sample and posterior

$\sigma_\epsilon = 1.$



# Prior, sample and posterior

$\sigma_\epsilon = 1.$



# Laplace's rule of succession from a prior

Take a uniform prior on $q$ and $N$ Bernoulli trials:

$$q \sim \text{Uniform}(0, 1)$$

$$Y_i \mid q \sim \text{Bernoulli}(q), \quad i = 1, \ldots, N, \qquad S = \sum_i Y_i$$

# Laplace's rule of succession from a prior

Take a uniform prior on $q$ and $N$ Bernoulli trials:

$$q \sim \text{Uniform}(0, 1)$$

$$Y_i \mid q \sim \text{Bernoulli}(q), \quad i = 1, \ldots, N, \qquad S = \sum_i Y_i$$

The posterior mean is Laplace's rule of succession:

$$E[q \mid S] = \frac{S + 1}{N + 2}.$$

# Laplace's rule of succession from a prior


Take a uniform prior on $q$ and $N$ Bernoulli trials:

$$q \sim \text{Uniform}(0, 1)$$

$$Y_i \mid q \sim \text{Bernoulli}(q), \quad i = 1, \ldots, N, \qquad S = \sum_i Y_i$$

The posterior mean is Laplace's rule of succession:

$$E[q \mid S] = \frac{S + 1}{N + 2}.$$



# Regularization from priors

Put a prior on each coefficient:

$$\beta_j \sim \text{Normal}(0, \tau_j^2), \quad j = 1, \ldots, K$$

$$Y \mid X, \beta \sim \text{Normal}\Big( \sum_j \beta_j \phi_j(X),\ \sigma_\epsilon^2 \Big)$$

# Regularization from priors

Put a prior on each coefficient:

$$\beta_j \sim \text{Normal}(0, \tau_j^2), \quad j = 1, \ldots, K$$
$$Y \mid X, \beta \sim \text{Normal}\Big( \sum_j \beta_j \phi_j(X),\ \sigma_\epsilon^2 \Big)$$

The posterior mean of $\beta$ minimizes

$$\hat{\sigma}^2_{\text{train}} + \frac{1}{N} \sum_j \left( \frac{\sigma_\epsilon}{\tau_j} \right)^2 \beta_j^2,$$

so a prior is a penalty: with all $\tau_j = \tau$ this is ridge regression with $\lambda = \sigma_\epsilon^2 / (N\tau^2)$.

# Regularization from priors

Put a prior on each coefficient:

$$\beta_j \sim \text{Normal}(0, \tau_j^2), \quad j = 1, \ldots, K$$
$$Y \mid X, \beta \sim \text{Normal}\Big( \sum_j \beta_j \phi_j(X),\ \sigma_\epsilon^2 \Big)$$


The posterior mean of $\beta$ minimizes

$$\hat{\sigma}^2_{\text{train}} + \frac{1}{N} \sum_j \left( \frac{\sigma_\epsilon}{\tau_j} \right)^2 \beta_j^2,$$

so a prior is a penalty: with all $\tau_j = \tau$ this is ridge regression with $\lambda = \sigma_\epsilon^2 / (N\tau^2)$.

# Posterior distribution of the regression function

Now let's "forget" about the $\beta_j$ (that is marginalize over them), and examine the distribution of the function $f(x)$. Here are some random samples for different choices of $\tau_j$.



# Covariance of a random function

Since $E[\beta_j] = 0$, we have $E[f(x)] = 0$. Since the $\beta_j$ are independent,

$$\operatorname{cov}\big(f(x), f(x')\big) = \sum_{j=1}^{J} \tau_j^2 \, \phi_j(x)\phi_j(x') = K(x, x').$$

$K$ is called the kernel.

## Question

Suppose we observe the value of $f$ at a single point $x'$. What is $E\big[f(x) \mid f(x')\big]$?

# Solution

Both $f(x)$ and $f(x')$ have mean 0, and

$$\text{var}\big(f(x')\big) = K(x', x'), \qquad \text{cov}\big(f(x), f(x')\big) = K(x, x').$$

# Solution

Both $f(x)$ and $f(x')$ have mean 0, and

$$\text{var}\big(f(x')\big) = K(x', x'), \qquad \text{cov}\big(f(x), f(x')\big) = K(x, x').$$

Regress $f(x)$ (response) on $f(x')$ (predictor). The intercept is 0 and the slope is

$$\frac{\text{cov}\big(f(x), f(x')\big)}{\text{var}\big(f(x')\big)} = \frac{K(x, x')}{K(x', x')}.$$

# Solution

Both $f(x)$ and $f(x')$ have mean 0, and

$$\text{var}\big(f(x')\big) = K(x', x'), \qquad \text{cov}\big(f(x), f(x')\big) = K(x, x').$$

Regress $f(x)$ (response) on $f(x')$ (predictor). The intercept is 0 and the slope is

$$\frac{\text{cov}\big(f(x), f(x')\big)}{\text{var}\big(f(x')\big)} = \frac{K(x, x')}{K(x', x')}.$$

Since $f(x)$ and $f(x')$ are sums of Normals, they are jointly Normal, so the regression line is the conditional mean:

$$E\big[f(x) \mid f(x')\big] = \frac{K(x, x')}{K(x', x')}\, f(x').$$