
Lecture 18 (10/26)


Review for Midterm 2

Derive-athon: population formulas



Unit notes: ① §2.3 the slope ② §2.3 regression to the mean ③ §2.4 coeff. of determination ④ §2.4 correlation ⑤ §4.2 omitted variable bias

Derive-athon: estimators



Unit notes: 6 §3.2 bias-variance tradeoff 7 §3.4 sample distribution of $\hat{\beta}_1$ 8 §2.1, §3.1 CLT 9 §1.4, §3.1; $\text{var}(Y_i) = \frac{1}{12}$ in the 9/25 slides

True or False

1) If an estimator is unbiased, its mean-squared error equals its variance.

True or False

1) If an estimator is unbiased, its mean-squared error equals its variance.
2) Doubling the sample size halves the standard error of $\hat{\beta}_1$.

True or False

1) If an estimator is unbiased, its mean-squared error equals its variance.
2) Doubling the sample size halves the standard error of $\hat{\beta}_1$.
3) If adding X_2 to a model leaves the coefficient on X_1 unchanged, then X_2 is uncorrelated with X_1.

True or False

1) If an estimator is unbiased, its mean-squared error equals its variance.
2) Doubling the sample size halves the standard error of $\hat{\beta}_1$.
3) If adding X_2 to a model leaves the coefficient on X_1 unchanged, then X_2 is uncorrelated with X_1.
4) If X_1 and X_2 are uncorrelated, they are also independent.

True or False

1) If an estimator is unbiased, its mean-squared error equals its variance.
2) Doubling the sample size halves the standard error of $\hat{\beta}_1$.
3) If adding X_2 to a model leaves the coefficient on X_1 unchanged, then X_2 is uncorrelated with X_1.
4) If X_1 and X_2 are uncorrelated, they are also independent.
5) If X_1 and X_2 are positively correlated and both have positive coefficients in the two-predictor model, then leaving out either predictor increases the coefficient on the other.

True or False

1) If an estimator is unbiased, its mean-squared error equals its variance.
2) Doubling the sample size halves the standard error of $\hat{\beta}_1$.
3) If adding X_2 to a model leaves the coefficient on X_1 unchanged, then X_2 is uncorrelated with X_1.
4) If X_1 and X_2 are uncorrelated, they are also independent.
5) If X_1 and X_2 are positively correlated and both have positive coefficients in the two-predictor model, then leaving out either predictor increases the coefficient on the other.
6) A coefficient that is statistically significant in a single-predictor model stays significant after another predictor is added.

True or False

1) If an estimator is unbiased, its mean-squared error equals its variance.
2) Doubling the sample size halves the standard error of $\hat{\beta}_1$.
3) If adding X_2 to a model leaves the coefficient on X_1 unchanged, then X_2 is uncorrelated with X_1.
4) If X_1 and X_2 are uncorrelated, they are also independent.
5) If X_1 and X_2 are positively correlated and both have positive coefficients in the two-predictor model, then leaving out either predictor increases the coefficient on the other.
6) A coefficient that is statistically significant in a single-predictor model stays significant after another predictor is added.
7) In a stationary AR(1) process, $\mathrm{cov}(Y_t, Y_{t+k})$ doesn't depend on k

True or False

1) If an estimator is unbiased, its mean-squared error equals its variance.
2) Doubling the sample size halves the standard error of $\hat{\beta}_1$.
3) If adding X_2 to a model leaves the coefficient on X_1 unchanged, then X_2 is uncorrelated with X_1.
4) If X_1 and X_2 are uncorrelated, they are also independent.
5) If X_1 and X_2 are positively correlated and both have positive coefficients in the two-predictor model, then leaving out either predictor increases the coefficient on the other.
6) A coefficient that is statistically significant in a single-predictor model stays significant after another predictor is added.
7) In a stationary AR(1) process, $\mathrm{cov}(Y_t, Y_{t+k})$ doesn't depend on k
8) For a categorical predictor with three categories, changing which category is the baseline changes the R^2.

True or False

1) If an estimator is unbiased, its mean-squared error equals its variance.
2) Doubling the sample size halves the standard error of $\hat{\beta}_1$.
3) If adding X_2 to a model leaves the coefficient on X_1 unchanged, then X_2 is uncorrelated with X_1.
4) If X_1 and X_2 are uncorrelated, they are also independent.
5) If X_1 and X_2 are positively correlated and both have positive coefficients in the two-predictor model, then leaving out either predictor increases the coefficient on the other.
6) A coefficient that is statistically significant in a single-predictor model stays significant after another predictor is added.
7) In a stationary AR(1) process, $\mathrm{cov}(Y_t, Y_{t+k})$ doesn't depend on k
8) For a categorical predictor with three categories, changing which category is the baseline changes the R^2.
9) If data $(Y_i, X_{1,i}, X_{2,i})$ satisfy the regression assumptions for the two-predictor model, then $(Y_i, X_{1,i})$ satisfy them for the single-predictor model.

True or False

1) If an estimator is unbiased, its mean-squared error equals its variance.
2) Doubling the sample size halves the standard error of $\hat{\beta}_1$.
3) If adding X_2 to a model leaves the coefficient on X_1 unchanged, then X_2 is uncorrelated with X_1.
4) If X_1 and X_2 are uncorrelated, they are also independent.
5) If X_1 and X_2 are positively correlated and both have positive coefficients in the two-predictor model, then leaving out either predictor increases the coefficient on the other.
6) A coefficient that is statistically significant in a single-predictor model stays significant after another predictor is added.
7) In a stationary AR(1) process, $\mathrm{cov}(Y_t, Y_{t+k})$ doesn't depend on k
8) For a categorical predictor with three categories, changing which category is the baseline changes the R^2.
9) If data $(Y_i, X_{1,i}, X_{2,i})$ satisfy the regression assumptions for the two-predictor model, then $(Y_i, X_{1,i})$ satisfy them for the single-predictor model.

True or False: answers

1) **True.** $\text{MSE} = \text{se}(\hat{\theta})^2 + \text{bias}^2$, and the bias is zero.

True or False: answers

1) **True.** $\text{MSE} = \text{se}(\hat{\theta})^2 + \text{bias}^2$, and the bias is zero.
2) **False.** $\text{se}(\hat{\beta}_1) \approx \sigma_\epsilon / (\sigma_X \sqrt{N})$, so it shrinks by a factor of $1/\sqrt{2}$.

True or False: answers

1) **True.** $\text{MSE} = \text{se}(\hat{\theta})^2 + \text{bias}^2$, and the bias is zero.
2) **False.** $\text{se}(\hat{\beta}_1) \approx \sigma_\epsilon / (\sigma_X \sqrt{N})$, so it shrinks by a factor of $1/\sqrt{2}$.
3) **False.** By omitted variable bias the change is $\hat{\beta}_2 \hat{\beta}_{1,2}$, which is zero when $\hat{\beta}_2 = 0$, however correlated X_1 and X_2 are.

True or False: answers

1) **True.** $\text{MSE} = \text{se}(\hat{\theta})^2 + \text{bias}^2$, and the bias is zero.
2) **False.** $\text{se}(\hat{\beta}_1) \approx \sigma_\epsilon / (\sigma_X \sqrt{N})$, so it shrinks by a factor of $1/\sqrt{2}$.
3) **False.** By omitted variable bias the change is $\hat{\beta}_2 \hat{\beta}_{1,2}$, which is zero when $\hat{\beta}_2 = 0$, however correlated X_1 and X_2 are.
4) **False.** Uncorrelated only rules out a linear relationship: if $X_1 \sim \text{Normal}(0, 1)$ and $X_2 = X_1^2$, then $\textbf{cov}(X_1, X_2) = E[X_1^3] = 0$, but X_2 is determined by X_1. (Independent variables are always uncorrelated.)

True or False: answers

1) **True.** $\text{MSE} = \text{se}(\hat{\theta})^2 + \text{bias}^2$, and the bias is zero.
2) **False.** $\text{se}(\hat{\beta}_1) \approx \sigma_\epsilon/(\sigma_X\sqrt{N})$, so it shrinks by a factor of $1/\sqrt{2}$.
3) **False.** By omitted variable bias the change is $\hat{\beta}_2\hat{\beta}_{1,2}$, which is zero when $\hat{\beta}_2 = 0$, however correlated X_1 and X_2 are.
4) **False.** Uncorrelated only rules out a linear relationship: if $X_1 \sim \text{Normal}(0, 1)$ and $X_2 = X_1^2$, then $\text{cov}(X_1, X_2) = E[X_1^3] = 0$, but X_2 is determined by X_1. (Independent variables are always uncorrelated.)
5) **True.** $\beta_1' = \beta_1 + \beta_2\beta_{1,2}$ and $\beta_2' = \beta_2 + \beta_1\beta_{2,1}$; $\beta_{1,2}$ and $\beta_{2,1}$ have the sign of $\text{cov}(X_1, X_2)$, so both added terms are positive.

True or False: answers

1) **True.** $\text{MSE} = \text{se}(\hat{\theta})^2 + \text{bias}^2$, and the bias is zero.
2) **False.** $\text{se}(\hat{\beta}_1) \approx \sigma_\epsilon/(\sigma_X\sqrt{N})$, so it shrinks by a factor of $1/\sqrt{2}$.
3) **False.** By omitted variable bias the change is $\hat{\beta}_2\hat{\beta}_{1,2}$, which is zero when $\hat{\beta}_2 = 0$, however correlated X_1 and X_2 are.
4) **False.** Uncorrelated only rules out a linear relationship: if $X_1 \sim \text{Normal}(0, 1)$ and $X_2 = X_1^2$, then $\text{cov}(X_1, X_2) = E[X_1^3] = 0$, but X_2 is determined by X_1. (Independent variables are always uncorrelated.)
5) **True.** $\beta_1' = \beta_1 + \beta_2\beta_{1,2}$ and $\beta_2' = \beta_2 + \beta_1\beta_{2,1}$; $\beta_{1,2}$ and $\beta_{2,1}$ have the sign of $\text{cov}(X_1, X_2)$, so both added terms are positive.
6) **False.** Adding a correlated predictor can increase the standard error and hence the p value.

True or False: answers

1) **True.** $\text{MSE} = \text{se}(\hat{\theta})^2 + \text{bias}^2$, and the bias is zero.
2) **False.** $\text{se}(\hat{\beta}_1) \approx \sigma_\epsilon/(\sigma_X\sqrt{N})$, so it shrinks by a factor of $1/\sqrt{2}$.
3) **False.** By omitted variable bias the change is $\hat{\beta}_2\hat{\beta}_{1,2}$, which is zero when $\hat{\beta}_2 = 0$, however correlated X_1 and X_2 are.
4) **False.** Uncorrelated only rules out a linear relationship: if $X_1 \sim \text{Normal}(0, 1)$ and $X_2 = X_1^2$, then $\text{cov}(X_1, X_2) = E[X_1^3] = 0$, but X_2 is determined by X_1. (Independent variables are always uncorrelated.)
5) **True.** $\beta_1' = \beta_1 + \beta_2\beta_{1,2}$ and $\beta_2' = \beta_2 + \beta_1\beta_{2,1}$; $\beta_{1,2}$ and $\beta_{2,1}$ have the sign of $\text{cov}(X_1, X_2)$, so both added terms are positive.
6) **False.** Adding a correlated predictor can increase the standard error and hence the p value.
7) **False.** $\text{cov}(Y_t, Y_{t+k}) = \rho^k\sigma_\epsilon^2/(1-\rho^2)$ shrinks as k grows. Stationary means it does not depend on t.

True or False: answers

1) **True.** $\text{MSE} = \text{se}(\hat{\theta})^2 + \text{bias}^2$, and the bias is zero.
2) **False.** $\text{se}(\hat{\beta}_1) \approx \sigma_\epsilon/(\sigma_X\sqrt{N})$, so it shrinks by a factor of $1/\sqrt{2}$.
3) **False.** By omitted variable bias the change is $\hat{\beta}_2\hat{\beta}_{1,2}$, which is zero when $\hat{\beta}_2 = 0$, however correlated X_1 and X_2 are.
4) **False.** Uncorrelated only rules out a linear relationship: if $X_1 \sim \text{Normal}(0, 1)$ and $X_2 = X_1^2$, then $\text{cov}(X_1, X_2) = E[X_1^3] = 0$, but X_2 is determined by X_1. (Independent variables are always uncorrelated.)
5) **True.** $\beta_1' = \beta_1 + \beta_2\beta_{1,2}$ and $\beta_2' = \beta_2 + \beta_1\beta_{2,1}$; $\beta_{1,2}$ and $\beta_{2,1}$ have the sign of $\text{cov}(X_1, X_2)$, so both added terms are positive.
6) **False.** Adding a correlated predictor can increase the standard error and hence the p value.
7) **False.** $\text{cov}(Y_t, Y_{t+k}) = \rho^k\sigma_\epsilon^2/(1-\rho^2)$ shrinks as k grows. Stationary means it does not depend on t.
8) **False.** Either way the model has one mean per category, so the fitted values and R^2 are the same.

True or False: answers

1) **True.** $\text{MSE} = \text{se}(\hat{\theta})^2 + \text{bias}^2$, and the bias is zero.
2) **False.** $\text{se}(\hat{\beta}_1) \approx \sigma_\epsilon/(\sigma_X\sqrt{N})$, so it shrinks by a factor of $1/\sqrt{2}$.
3) **False.** By omitted variable bias the change is $\hat{\beta}_2\hat{\beta}_{1,2}$, which is zero when $\hat{\beta}_2 = 0$, however correlated X_1 and X_2 are.
4) **False.** Uncorrelated only rules out a linear relationship: if $X_1 \sim \text{Normal}(0, 1)$ and $X_2 = X_1^2$, then $\text{cov}(X_1, X_2) = E[X_1^3] = 0$, but X_2 is determined by X_1. (Independent variables are always uncorrelated.)
5) **True.** $\beta_1' = \beta_1 + \beta_2\beta_{1,2}$ and $\beta_2' = \beta_2 + \beta_1\beta_{2,1}$; $\beta_{1,2}$ and $\beta_{2,1}$ have the sign of $\text{cov}(X_1, X_2)$, so both added terms are positive.
6) **False.** Adding a correlated predictor can increase the standard error and hence the p value.
7) **False.** $\text{cov}(Y_t, Y_{t+k}) = \rho^k\sigma_\epsilon^2/(1-\rho^2)$ shrinks as k grows. Stationary means it does not depend on t.
8) **False.** Either way the model has one mean per category, so the fitted values and R^2 are the same.
9) **False.** If $X_2 = X_1^2$, then $E[Y \mid X_1] = \beta_0 + \beta_1 X_1 + \beta_2 X_1^2$ is not linear in X_1. It holds if $X_2 = a + \beta_{1,2}X_1 + \eta$ with η Normal and independent of X_1 (noise $\beta_2\eta + \epsilon$).

Switching the axes

Suppose

$$Y = \beta_0 + \beta_1 X + \epsilon$$

with ϵ independent of X, and let ρ be the correlation between X and Y. Now fit the single-predictor model with X as the response and Y as the predictor:

$$X = \gamma_0 + \gamma_1 Y + \eta.$$

Question

Write γ_1 in terms of β_1 and ρ. When is $\gamma_1 = 1/\beta_1$?

See also Problem 2.5 (swapping response and predictor variables) in the Unit 2 notes.