

☆ THE ART OF COMPUTER PROGRAMMING ☆

☆☆☆☆ ERRATA TO VOLUME 1 (3rd edition) ☆☆☆☆

This document is a transcript of the notes that I have been making in my personal copy of *The Art of Computer Programming*, Volume 1 (third edition) since it was first printed in 1997.

Four levels of updates — “errors,” “amendments,” “plans,” and “improvements” — appear, indicated by four different typographic conventions:

► **Page 666** line 1 _____ 04 Jul 1776

Technical or typographical errors (aka bugs) are the most critical items, so they are flagged with a ‘►’ preceding the page number. The date on which I first was told about the bug is shown; this is the effective date on which I paid the finder’s fee. The necessary corrections are indicated in a straightforward way. If, for example, the book says ‘ n ’ where it should have said ‘ $n + 1$ ’, the change is shown thus:

$n \rightsquigarrow n + 1$

Page 666 line 2 _____ 14 Jul 1789

Amendments to the text appear in the same format as bugs, but without the ‘►’. These are things I wish I had known about or thought of when I wrote the original text, so I added them later. The date is the date I drafted the new text.

Page 666 line 3 20 Nov 1917


Plans for the future represent a third kind of item. In such notes I sketched my intentions about things that I wasn’t ready to flesh out further when I wrote them down. You can identify these items because they’re written in slanted type, and preceded by a bunch of dots ‘.....’ leading to the date on which I recorded the plan in my files.

Page 666 line 4 _____ 10 Jan 1938

The fourth and final category — indicated by page and line number in smaller, slanted type — consists of minor corrections or improvements that most readers don’t want to know about, because they are so trivial. You wouldn’t even be seeing these items if you hadn’t specifically chosen to print the complete errata list in all its gory details. Are you sure you wanted to do that?

My shelves at home are bursting with preprints and reprints of significant research results that I want to digest and summarize, where appropriate, in the ultimate edition of Volume 1. I didn’t do that in the third edition because I would surely have to do it over again later: New results continue to pour forth at a great rate, and I will have time to rewrite that volume only once. Volumes 4 and 5 need to be finished first. So I’ve put most of my effort so far into writing up those parts of the total picture that seem to have converged to their near-final form. It follows, somewhat paradoxically, that the updates in this document are most current in the areas where there has been least activity.

On the other hand I do believe that the changes listed here bring Volume 1 completely up to date in two respects: (1) All of the research problems in the previous edition — i.e., all exercises that were rated 46 and above — have received new ratings of 45 or less whenever I learned of a solution; and in such cases, the answer now refers to that solution. (2) All of the historical information about pioneering developments has been amended whenever new details have come to my attention.

 *The ultimate, glorious, future editions of Volumes 1–3 are works in progress. Please let me know of any improvements that you think I ought to make. Send your comments either by snail mail to D. E. Knuth, Computer Science, Gates Building 4B, Stanford University, Stanford CA 94305-9045, or by email to taocp@cs.stanford.edu. (Use email for book suggestions only, please — all other correspondence is returned unread to the sender, or discarded, because I have no time to read ordinary email.) Although I'm working full time on Volume 4B these days, I will try to reply to all such messages within a year of receipt. Current news about The Art of Computer Programming is posted on*

<http://www-cs-faculty.stanford.edu/~knuth/taocp.html>

and updated regularly.

—Don Knuth, April 1997

*What happened?
The subject took the bit in its teeth and ran away with it,
that's what happened.
I know now how Sir James Frazer felt when,
after setting out to dash off a brief monograph
on a single obscure rite, he found himself
in the embarrassing possession of
the 12 volumes of "The Golden Bough."
— WAVERLEY ROOT (1974)*



FUNDAMENTAL ALGORITHMS



Copyright © 1997, 1998, 1999, 2000, 2001, 2002, 2003, 2004, 2005,
2006, 2007, 2008, 2009, 2010, Addison–Wesley; all rights reserved
Last updated 28 February 2011

Most of these corrections have already been made in recent printings.

Page iv line 4 of the Library of Congress Data _____ 07 Jan 2011
xx,650p. ↗ xx,652p.

Page vii lines 13 and 14 _____ 05 Jan 2011
three separate books (Volumes 4A, 4B, and 4C) ↗ several separate books
(Volumes 4A, 4B, and so on)

Page vii line 15 _____ 30 Dec 1997
Volume 6, *The Theory of Languages*; Volume 7, *Compilers*. ↗ Volume 6, *The
Theory of Languages* (Chapter 11); Volume 7, *Compilers* (Chapter 12).

Page xi line –7 _____ 28 Oct 2008
cheerfully pay \$2.56 ↗ cheerfully award \$2.56

► **Page xii** step 2 of the flowchart _____ 16 Jul 1997
pp. xvii–xix ↗ pp. xv–xvii

Page xv line 11 _____ 09 Oct 2010
This is sometimes unfortunate ↗ A motley mixture is, however, often unfortunate

Page xvi line –6 _____ 09 Oct 2010
This is not ↗ This distinction is not

Page xvii line 9 _____ 09 Oct 2010
Later editions ↗ Later printings

Page xix replacement for the bottom line _____ 07 Jan 2011

Appendix C — Index to Algorithms and Theorems 628

Index and Glossary 630

► **Page xviii** lines 12–19 _____ 01 Jul 1998
1.2.8 ↗ 1.2.8. 1.3 ↗ 1.3.

► **Page 1** line 11 _____ 26 Feb 2007
(1844) ↗ (1843)

Page 1 line –5 _____ 10 Apr 2001
celebrated book ↗ celebrated Arabic text

► **Page 1** line –4 _____ 01 Aug 1997
Kitab al jabr ↗ *Kitāb al-jabr*

Page 3 line 3 after Fig. 1 _____ 03 Feb 2000

Step E3 \rightsquigarrow step E3

Page 4 line -15 _____ 09 Apr 2003

$$\frac{544}{119} = 4\frac{68}{119} \rightsquigarrow 544/119 = 4 + 68/119$$

Page 6 lines 1 and 2 _____ 02 Mar 2000

set of divisors \rightsquigarrow set of common divisors (twice)

►Page 8 three places in (3) _____ 05 Aug 2000

$$f(\sigma, j) \rightsquigarrow f((\sigma, j)) \quad [\text{twice}]$$

$$f(\sigma, N) \rightsquigarrow f((\sigma, N)) \quad [\text{once}]$$

►Page 9 line 1 of exercise 5 _____ 25 Dec 2010

appears in \rightsquigarrow appears after

Page 9 line 2 of exercise 5 _____ 25 Dec 2010

three \rightsquigarrow at least three

Page 9 lines 6 and 7 of exercise 9 _____ 02 May 2008

I_2 might be the initial state ... values of m and n . \rightsquigarrow I_2 might be the set of initial states, each including the program that determines the greatest common divisor as well as the particular values of m and n .

Page 12 line -5 _____ 06 Aug 2002

8.] \rightsquigarrow 8. See also Section 7.2.1.4.]

Page 13 line 23 _____ 19 Mar 2008

$$\phi^0 = \phi^{n-1} \rightsquigarrow \phi^0 = \phi^{1-1}$$

Page 13 line -3 _____ 21 May 2005

d and two integers a and b , \rightsquigarrow d , and we also compute two not-necessarily-positive integers a and b

►Page 15 replacement for (7) _____ 26 Jul 1997

*if an assertion attached to any arrow leading into the box is true
before the operation in that box is performed, then all of the
assertions on relevant arrows leading away from the box are true
after the operation.* (7)

►Page 16 line -7 _____ 24 Mar 1999

$m, n, c,$ and $r \rightsquigarrow m, n, c, d,$ and r

Page 16 near the bottom _____ 24 Mar 1999

line -5: the algorithm with \rightsquigarrow the method with

line -2: of the algorithm \rightsquigarrow of the procedure

line -1: start the procedure with \rightsquigarrow start with the values

Page 17 lines 26 and 27 _____ 29 Mar 2005

[See AMM 24 \rightsquigarrow [See *The Penny Cyclopædia* 11 (1838), 465–466; AMM 24

Page 19 add clarification to line 3 _____ 23 Nov 1997

(A prime number is considered to be the “product” of a single prime, namely itself.)

Page 19 lines 11–12 _____ 05 Aug 2004

suggested to the author by R. W. Floyd, is shown in Fig. 5. \rightsquigarrow suggested by Warren Lushbaugh, is shown in Fig. 5; see *Math. Gazette* **49** (1965), 200.

Page 19 lines 18 and 20 _____ 20 Sep 2002

$(1 \rightsquigarrow \cdot (1$

►Page 19 in the Area part of Fig. 5 _____ 30 Jul 1997

$4 \cdot 3 \cdot 3^2 + 4 \cdot 5 \cdot 5^2 \rightsquigarrow 4 \cdot 3 \cdot 3^2 + 4 \cdot 4 \cdot 4^2 + 4 \cdot 5 \cdot 5^2$

►Page 19 line 4 of exercise 13 _____ 27 Nov 1997

“ $m > 0, n = 0, T = 0.$ ” \rightsquigarrow “ $m > 0, n > 0, T = 0.$ ”

►Page 22 line 8 _____ 28 Jul 1997

m th root of v , \rightsquigarrow m th root of u ,

Page 22 line 10 _____ 28 Jul 1997

numbers $r \rightsquigarrow$ numbers $r = p/q$

Page 22 line –12 _____ 02 Oct 1998

$1^x = 1 \rightsquigarrow b^x = 1$

Page 23 line –1 _____ 08 Aug 2002

the return \rightsquigarrow the annual return

Page 25 line 1 _____ 21 Apr 1999

$0.0100110100010000011 \dots \rightsquigarrow (0.0100110100010000011 \dots)_2$

Page 26 first line of exercise 26 _____ 20 Mar 2001

Determine upper bounds on the accuracy of \rightsquigarrow Find a rigorous upper bound on the error made by

Page 27 new exercise for Section 1.2.2 _____ 28 Jun 2003

30. [12] Simplify the expression $(\ln n)^{\ln n / \ln \ln n}$, assuming that $n > 1$.

Page 27 the line following (1) _____ 04 Jan 2010

If n is zero, \dots zero. In general, \rightsquigarrow

If n is zero, the value of $a_1 + a_2 + \dots + a_n = \sum_{j=1}^n a_j = \sum_{1 \leq j \leq n} a_j$ is defined to be zero. Our convention of using “three dots” in sums such as $a_1 + a_2 + \dots + a_n$ therefore has some slightly peculiar, but sensible, behavior in borderline cases (see exercise 1).

In general,

Page 27 lines 9 and 10 from the bottom _____ 20 Mar 2001

The example in the preceding sentence \rightsquigarrow A sum such as $\sum_{j \leq k} \binom{j+k}{2j-k}$

►Page 27 line 4 from the bottom _____ 16 Jul 1997

$$\sum_{j=1}^{\infty} = \rightsquigarrow \sum_{j=1}^{\infty} a_j =$$

►Page 27 replacement for the bottom line _____ 16 Jul 1997

$$\sum_{R(j)} a_j = \left(\lim_{n \rightarrow \infty} \sum_{\substack{R(j) \\ -n \leq j < 0}} a_j \right) + \left(\lim_{n \rightarrow \infty} \sum_{\substack{R(j) \\ 0 \leq j < n}} a_j \right), \quad (3)$$

Page 28 line -11 _____ 06 Aug 2001

the range \rightsquigarrow the relevant values

Page 33 near the top _____ 06 Aug 2001

line 3: range \rightsquigarrow relevant values

line 4: a permutation of the range \rightsquigarrow such a permutation

►Page 33 line -5 _____ 29 Jul 1997

$$[1 \leq i \leq n][1 \leq j \leq n] \rightsquigarrow [1 \leq i \leq n][1 \leq j \leq i]$$

Page 34 replacements for exercises 1-3 _____ 04 Jan 2010

►1. [10] The text says that $a_1 + a_2 + \cdots + a_0 = 0$. What, then, is $a_2 + \cdots + a_0$?

2. [01] What does the notation $\sum_{1 \leq j \leq n} a_j$ mean, if $n = 3.14$?

►3. [13] Without using the \sum -notation, write out the equivalent of

$$\sum_{0 \leq n \leq 5} \frac{1}{2n+1},$$

and also the equivalent of

$$\sum_{0 \leq n^2 \leq 5} \frac{1}{2n^2+1}.$$

Explain why the two results are different, in spite of rule (b).

Page 34 line 1 of exercise 15 _____ 12 Jun 2002

$$n2^n \rightsquigarrow n \times 2^n$$

►Page 35 exercise 21 _____ 07 Feb 2004

from ... (16). \rightsquigarrow from (8) and (17).

►Page 37 line -2 _____ 15 Jul 1997

$$a_{ij} = 1/(x_i + y_i) \rightsquigarrow a_{ij} = 1/(x_i + y_j)$$

Page 39 line 6 of section 1.2.4 _____ 25 Feb 1999

because both \rightsquigarrow because

►Page 40 line -12 _____ 02 Aug 2000

(modulo m), \rightsquigarrow (modulo m).

Page 41 line 1 of exercise 6 _____ 12 Aug 1997

real numbers \rightsquigarrow real numbers

► **Page 42** line 3 of exercise 21 _____ 16 Jul 1997
factors. \rightsquigarrow factors).

Page 44 line 1 of exercise 45 _____ 15 Oct 2007
implies that \rightsquigarrow implies that, when m and n are positive integers,


► **Page 45** line 9 of exercise 47 _____ 23 Jun 2010
(modulo 2). \rightsquigarrow (modulo 2), unless q is a multiple of p .

► **Page 45** line 1 of exercise 49 _____ 07 Jan 1998
the function $f(x)$ \rightsquigarrow the integer-valued function $f(x)$

Page 48 line 8 _____ 27 Feb 2005
 $n!$. \rightsquigarrow $n!$. [See A. M. Legendre, *Essai sur la Théorie des Nombres*, second edition (Paris: 1808), page 8.]

Page 49 line 16 _____ 19 Jan 2005
1808) \rightsquigarrow 1808), page 219

Page 50 better labels on Figure 7 _____ 12 Aug 1998



Page 50 in Eq. (19) _____ 01 Jan 2006
 $(x + j)$; \rightsquigarrow $(x + j)$.

Page 50 lines -6 and -5 _____ 12 Jun 2003

[This notation ... 313.] \rightsquigarrow [The notations $x^{\bar{k}}$ and x^k are due respectively to A. Capelli, *Giornale di Mat. di Battaglini* **31** (1893), 291–313, and L. Toscano, *Comment. Accademia della Scienze* **3** (1939), 721–757.]

- Page 53** near the bottom _____ 30 Nov 2000
 line -5: Shih-Chieh Chu \rightsquigarrow Chu Shih-Chieh
 line -4: invention. \rightsquigarrow invention. Yang Hui, in 1261, credited them to Chia Hsien, whose work (c. 1000) is now lost.
- **Page 53** line -2 or -3, 12th thru 14th printings only _____ 28 Aug 2003
 Halāyuddha \rightsquigarrow Halāyudha
- Page 53* line -1 or -2 _____ 21 Sep 2004
Chandaḥ-sūtra \rightsquigarrow *Chandaḥśāstra*
- Page 54** opening lines _____ 23 Jan 2005
 In about 1150 ... were known \rightsquigarrow Another Indian mathematician, Mahāvīra, had previously explained rule (3) for computing $\binom{r}{k}$ in Chapter 6 of his *Gaṇita Sāra Saṅgraha*, written about 850; and in 1150 Bhāskara repeated Mahāvīra's rule near the end of his famous book *Līlāvati*. For small values of k , binomial coefficients were known
- Page 54** penultimate line of first paragraph _____ 23 Jan 2005
 Ettingshausen in his book \rightsquigarrow Ettingshausen in §31 of his book
- Page 54* caption to Figure 8 _____ 23 Jun 2001
 [the second line should be in 9-point type, not 10-point]
- Page 59* line -4 _____ 22 Aug 2002
Paris (1772) \rightsquigarrow (Paris, 1772)
- Page 59** line -3 _____ 19 Jan 2005
 489–498 \rightsquigarrow part 1, 489–498.
- Page 59* line -3 _____ 30 Nov 2000
 Shih-Chieh Chu's \rightsquigarrow Chu Shih-Chieh's
- **Page 59** line -2 _____ 27 Oct 2004
Civilization \rightsquigarrow *Civilisation*
- Page 60** line -13 _____ 30 Dec 2004
 with $n = -1$ \rightsquigarrow with $m = 0$ and $n = -1$
- **Page 61** line 13 _____ 19 Oct 1997
 $\binom{n+1-(n+1)}{n+1-1+0} \rightsquigarrow \binom{n+1-(n+1)}{n+1-1+0} \frac{1}{n+1}$
- **Page 63** line 12 _____ 21 Jul 2003
 233–248 \rightsquigarrow 233–247
- **Page 64** line 11 _____ 18 Dec 1999
 r of less \rightsquigarrow r or less

Page 69 line 6 of exercise 10 _____ 26 Jul 1997

(E. Lucas, 1877.) \rightsquigarrow (É. Lucas, 1877.)

Page 70 line 1 of exercise 17 _____ 30 Nov 2000

Chu-Vandermonde \rightsquigarrow Chu-Vandermonde

Page 71 line 3 of exercise 31 _____ 11 Oct 2001

nonnegative integers \rightsquigarrow integers

Page 71 replacement for exercise 33 _____ 19 Jan 2005

33. [M20] (A. Vandermonde, 1772.) Show that the binomial formula is valid also when it involves factorial powers instead of the ordinary powers. In other words, prove that

$$(x+y)^n = \sum_k \binom{n}{k} x^k y^{n-k}; \quad (x+y)^{\overline{n}} = \sum_k \binom{n}{k} x^{\overline{k}} y^{\overline{n-k}}.$$

Page 73 line 2 of exercise 56 _____ 31 May 2002

$n = \binom{a}{1} + \binom{b}{2} + \binom{c}{3}$ and $0 \leq a < b < c$. \rightsquigarrow $n = \binom{a}{3} + \binom{b}{2} + \binom{c}{1}$ and $a > b > c \geq 0$.

Page 74 replacement for exercise 66 _____ 08 Nov 2005

66. [HM30] Suppose x , y , and z are real numbers satisfying

$$\binom{x}{n} = \binom{y}{n} + \binom{z}{n-1},$$

where $x \geq n-1$, $y \geq n-1$, $z > n-2$, and n is an integer ≥ 2 . Prove that

$$\begin{aligned} \binom{x}{n-1} &\leq \binom{y}{n-1} + \binom{z}{n-2} && \text{if and only if} && y \geq z; \\ \binom{x}{n+1} &\leq \binom{y}{n+1} + \binom{z}{n} && \text{if and only if} && y \leq z. \end{aligned}$$

Page 74 line 1 of exercise 68 _____ 17 Mar 1998

De Moivre \rightsquigarrow de Moivre

Page 75 lines 6–9 _____ 05 May 2007

(Besides H_n, \dots the harmonic series.) \rightsquigarrow Besides H_n, \dots the harmonic series. Chinese bamboo strips written before 186 B.C. already explained how to compute $H_{10} = 7381/2520$, as an exercise in arithmetic. [See C. Cullen, *Historia Math.* **34** (2007), 10–44.]

Page 77 the line after (10) _____ 25 Feb 2000

the infinite series \rightsquigarrow the infinite series 1.2.9–(17)

Page 80 line 3 _____ 26 Jul 1997

sequence F_n \rightsquigarrow sequence $\langle F_n \rangle$

Page 80 line 7 _____ 20 Jul 2000

Hemachandra \rightsquigarrow Hemacandra

Page 80 line 10 _____ 16 Dec 2001

Johann Kepler \rightsquigarrow Johannes Kepler

- **Page 80** line 20 _____ 04 Jan 2006
 at most $k + 1$ \wedge at most $k - 1$
- Page 81** line 8 _____ 18 Nov 2008
 sculpture. \wedge sculpture. [See T. A. Cook, *The Curves of Life* (1914), 420.]
- Page 81* line 16 _____ 22 Aug 2002
Paris 1 (1680) \wedge *Sci. 1* (Paris, 1680)
- Page 81* line -8 _____ 27 Jul 1999
 multiple of F_k \wedge multiple of F_n
- Page 82* line 7 _____ 26 Jul 1997
 sequence, F_n , for which statement (8) holds \wedge sequence $\langle F_n \rangle$ for which statement (8) holds,
- **Page 83** line 8 _____ 15 Oct 2007
 quadratic equation \wedge quadratic polynomial
- Page 83** lines 2–4 after (14) _____ 22 Sep 1999
 by A. De Moivre ... essentially \wedge early in the eighteenth century. (See D. Bernoulli, *Comment. Acad. Sci. Petrop.* **3** (1728), 85–100, §7; see also A. de Moivre, *Philos. Trans.* **32** (1722), 162–178, who showed how to solve general linear recurrences in essentially
- Page 86* bottom line _____ 07 Feb 2004
 (The k 's \wedge (See exercise 34. The k 's
- Page 87* line 15 _____ 17 Mar 1998
 A. De Moivre \wedge A. de Moivre
- **Page 87** line 22 _____ 27 Jul 1997
 125–169) \wedge 125–169
- **Page 88** line -3 _____ 27 Feb 1998
 $(a_0 + b_0)$ \wedge $(a_0 b_0)$
- Page 89* on the left side of (13) _____ 15 Oct 2007
 $n \bmod m = r$ \wedge $n \geq 0, n \bmod m = r$
- **Page 90** in Eq. (15) _____ 07 Jan 1998

$$\sum_{k \geq 0} \wedge \sum_{k \geq 1}$$
- Page 94* exercise 10 _____ 18 Aug 2002
 [Change the notation from ' a_m ' to the more modern ' e_m ', throughout this exercise.]
- **Page 95** line 6 of exercise 19 _____ 17 Jul 1997

$$\left(\frac{1}{n} - \frac{1}{n+x}\right) \wedge \left(\frac{1}{n} - \frac{1}{n+x}\right)$$

- Page 98 line -9 _____ 22 Aug 2002
 Paris **37** (1853) \rightsquigarrow **37** (Paris, 1853)
- Page 98 line -2 _____ 12 Sep 1997
 $(n-1)P_{(n-k)k} \rightsquigarrow (n-1)P_{(n-1)k}$
- Page 103 line -12 _____ 04 Oct 2005
 by Thiele in 1903. \rightsquigarrow by T. N. Thiele in 1889 [see A. Hald, *International Statistical Review* **68** (2000), 137–153].
- Page 103 in (23) _____ 29 Jul 1997
 $\kappa_1 t^2 \rightsquigarrow \kappa_2 t^2$
- Page 104 line 2 before the exercises _____ 30 Oct 2007
 for all $t \geq r$, \rightsquigarrow for all $t \geq r$, and if $f(t) \geq 0$ for all t in the domain of the random variable X ,
- Page 105 line -7 _____ 01 Jan 2006
 probability \rightsquigarrow Pr
- Page 106 line 1 of exercise 14 _____ 17 Mar 1998
 De Moivre \rightsquigarrow de Moivre
- Page 106 line 1 of exercise 18 _____ 01 Jan 2006
 distinct values \rightsquigarrow values
- Page 106 replacement for line 6 of exercise 18 _____ 15 Oct 2007

$$\left(\frac{k_n z}{k_n}\right) \left(\frac{k_{n-1} z + k_n}{k_{n-1} + k_n}\right) \left(\frac{k_{n-2} z + k_{n-1} + k_n}{k_{n-2} + k_{n-1} + k_n}\right) \cdots \left(\frac{k_1 z + k_2 + \cdots + k_n}{k_1 + k_2 + \cdots + k_n}\right) / z,$$
- Page 106 lines 3 and 4 of exercise 20 _____ 24 Dec 2001
 $a_1, a_2 \dots a_n \rightsquigarrow a_1 a_2 \dots a_n$
- Page 107 the line after (1) _____ 07 Sep 2001
 Oiler’s constant \rightsquigarrow Euler’s constant [pronounced ‘Oiler’s constant’]
- Page 108 line 3 _____ 31 Jul 1997
 $a_1 \rightsquigarrow a_1 n$
- Page 109 line -7 _____ 22 May 1998
 all value \rightsquigarrow all values
- Page 110 line -2 _____ 08 Nov 2005
 Big Theta notation \rightsquigarrow Big Theta
- Page 112 line -10 _____ 20 Mar 2006
 introduced by Jacques Bernoulli in his \rightsquigarrow introduced to European mathematicians in James Bernoulli’s
- Page 113 line 13 _____ 04 Nov 2001
 (9) \rightsquigarrow (9)

Page 114 line 10 _____ 10 Nov 2002

$$f^{(m)} \rightsquigarrow f^{(m)}(x)$$

Page 114 line 15 _____ 28 Nov 2002

$$\pm \int_1^\infty \rightsquigarrow - \int_1^\infty$$

►Page 115 line 1 of exercise 3 _____ 01 Jan 2006

$$((-1)^m (B_m/m!) \rightsquigarrow (B_m/m!)$$

Page 116 first line of exercise 8 _____ 01 Jan 2006

$$\ln(an^2 + bn)! \rightsquigarrow \ln(an^2 + bn)!$$

►Page 116 last line of exercise 8 _____ 03 May 1998

$$(1 + \epsilon) \rightsquigarrow (1 + \epsilon).$$

►Page 121 line 5 _____ 08 Feb 2000

$$\frac{1}{8505} \rightsquigarrow \frac{16}{8505}$$

Page 123 the displayed equation of exercise 17 _____ 27 Mar 1998

$$\sum_{0 \leq k} \rightsquigarrow \sum_{k \geq 0}$$


Page 125 line 1 _____ 17 Sep 2004

basic unit of information \rightsquigarrow basic unit of MIX data

Page 125 footnote _____ 03 Dec 1997

binary bits \rightsquigarrow binary digits

►Page 126 replacement for bottom of Fig. 13 _____ 15 Jul 1998



►Page 130 line 6 _____ 14 Feb 1999

right-hand of \rightsquigarrow right-hand portion of

►Page 130 line -15 _____ 14 Feb 1999

(store X), C \rightsquigarrow (store X). C

Page 131 line 13 _____ 03 Aug 1997

left of A \rightsquigarrow left of rA

Page 131 line -6 _____ 28 Sep 1997

10-byte number \rightsquigarrow 10-byte number rAX

►Page 131 line -2 _____ 28 Sep 1997

the quotient $\pm \lfloor |rA/V| \rfloor \rightsquigarrow$ the quotient $\pm \lfloor |rAX/V| \rfloor$

the remainder $\pm(|rA| \bmod |V|) \rightsquigarrow$ the remainder $\pm(|rAX| \bmod |V|)$

- Page 133 line -3 _____ 22 Aug 2005
 $C = 48 + i \rightsquigarrow C = 48 + i$
- Page 134 line 14 _____ 03 Aug 1997
 field of A \rightsquigarrow field of rA
- Page 135 line 14 _____ 18 Aug 2010
 These are \rightsquigarrow These 42 instructions are
- Page 135 line -7 _____ 24 Jan 2003
 $F = \text{number.} \rightsquigarrow F = \text{number, normally 1.}$
- Page 135 line -2 _____ 03 Sep 1997
 when the groups of locations involved overlap \rightsquigarrow when there's overlap between the locations involved
- Page 138 line 17 _____ 04 Apr 2003
 $40, \dots \text{convert} \rightsquigarrow 40, \dots \text{convert}$
- Page 138 line 23 _____ 03 Aug 1997
 register A and X \rightsquigarrow registers A and X
- Page 138 line -5 _____ 11 Oct 1998
 unspecified. \rightsquigarrow specified in Section 4.2.1.
- Page 139 line -4 in 1st or 2nd printing _____ 17 Apr 1998
 appears on page 133 \rightsquigarrow on page 133
- Page 139 line -4 in 3rd, 4th, 5th printings _____ 10 Jan 1999
 appears on page ?? \rightsquigarrow on page 133
- Page 142 line 3 of exercise 13 _____ 27 Feb 2000
 $\text{"JNOV 1001,} \rightsquigarrow \text{"JNOV 1001",}$
- Page 143 line 6 of exercise 23 _____ 19 Jul 1997
 to load \rightsquigarrow ability to load
- Page 144 last line of the exercise _____ 03 Sep 1997
 cannot be used; \rightsquigarrow instructions cannot be used;
- Page 145 line -24 _____ 03 Aug 1997
 $\text{"LOC OP ADDRESS"} \rightsquigarrow \text{"LOC", "OP", and "ADDRESS"}$
- Page 145 line -8 _____ 07 Sep 1998
 (page 1) \rightsquigarrow (page 96)
- Page 147 lines 16 and 19 _____ 18 Jul 2004
 program \rightsquigarrow algorithm (twice)
- Page 149 line -6 _____ 13 Jul 2008
 $25 \text{ and } 35 \rightsquigarrow 09, 25, \text{ and } 35$

- Page 150 line 15* _____ 05 May 2010
assembly procedure \curvearrowright assembly procedure (possibly with con2 first)
- Page 151 line 23* _____ 18 Apr 2010
“ALF” operation \curvearrowright pseudo-operation ALF
- Page 151 line -8* _____ 21 Sep 2007
42, and 48. \curvearrowright 42, 48, and 50.
- **Page 154 line 5** _____ 03 Aug 1997
that is., \curvearrowright that is,
- **Page 155 line 4 of point 10** _____ 19 Jul 1997
previous defined. \curvearrowright previously defined.
- Page 156** the line before the exercises _____ 14 Oct 2008
PL/MIX will be described in Chapter 10. \curvearrowright
- Page 156 lines 2–3 of exercise 1* _____ 13 Oct 2008
you wish to set \curvearrowright the algorithm is supposed to set
- **Page 157 line 1 of exercise 6** _____ 15 May 1999
 $1 \leq d \leq \sqrt{n}$ \curvearrowright $1 < d \leq \sqrt{n}$
- Page 159 line 1 of exercise 12** _____ 24 Apr 1997
[M47] \curvearrowright [HM42]
- Page 160 lines 11 and 12* _____ 10 Mar 2002
[March $((-D) \bmod 7)$ actually will be a Sunday.] \curvearrowright (March $((-D) \bmod 7)$ will actually be a Sunday.)
- Page 162 line -4 of exercise 20* _____ 21 May 1999
this traffic light \curvearrowright these lights
- **Page 162 line 4 of exercise 21** _____ 12 Jul 1997
junt below \curvearrowright just below
- Page 162 lines 10 and 11 of exercise 21** _____ 28 Mar 2000
Manuel Moschopoulos, who lived in Constantinople about 1300 \curvearrowright Ibn al-Haytham, who was born in Basra about 965 and died in Cairo about 1040
- **Page 163 line 3 after Fig. 18** _____ 10 Jul 1997
is what \curvearrowright is white
- Page 163** in the midst of the bottom paragraph _____ 18 Jan 2008
card corresponding to the matrix \curvearrowright card corresponding to the top row of the matrix
- Page 166 line 3* _____ 02 Nov 2005
[the spacing in this formula should be more consistent]
- Page 166 line 2 of step A1* _____ 30 Mar 1998
copy of the element \curvearrowright copy of the input symbol

- **Page 166** line -4 _____ 19 Jul 1997
 [See CURRENT.] $\wedge \rightarrow$ [Set CURRENT.]
- Page 167 first line of Table 1* _____ 27 May 1998
 The formula on this line should repeat the input formula (6).
- Page 168** line 8 _____ 24 Oct 2005
 cycles. $\wedge \rightarrow$ cycles. But it doesn't catch errors in the input.
- Page 168 line 30 of Program A* _____ 30 Mar 1998
 nonblank element $\wedge \rightarrow$ nonblank input symbol
- Page 170 lines 11 and 12* _____ 17 Sep 2005
 and it has led to some horribly inefficient uses of computing machinery. $\wedge \rightarrow$ and some
 horribly inefficient uses of computing machinery have arisen as a result.
- Page 170 replacement for lines -11 to -4* _____ 30 Nov 2009
- | | |
|------------|-------------------------|
| 26, 38, 39 | $A = 1 + (C - 1)$ |
| 33, 28, 29 | $C = B + (A - B)$ |
| 41, 84, 86 | $E = 1 + R$ |
| 42, 46, 48 | $F = E + (G - 1)$ |
| 64, 43, 44 | $H = F - G$ |
| 67, 70, 76 | $J = H + (K - (L - J))$ |
| 75, 79, 80 | $K = Q + (L - P)$ |
| 82, 72, 81 | $R = P - Q$ |
- **Page 172** line 5 _____ 31 May 2004
 (it involves ...), $\wedge \rightarrow$ (it involves $\dots + 3F + 4G + \dots + 3K + 4L + \dots = \dots + 7G + 7L + \dots$),
- Page 172 line -5* _____ 18 Aug 1997
homo sapiens $\wedge \rightarrow$ *Homo sapiens*
- **Page 174** line 26 of Program B _____ 13 Jul 2000
 ENT3 1 $\wedge \rightarrow$ ENT3 0
- **Page 176** step I5 of Algorithm I _____ 22 Feb 1998
 $X[i]$ was $\wedge \rightarrow X[-i]$ was
- **Page 178** line -5 _____ 24 Dec 2001
 permutation(1 6 3)(4 5) $\wedge \rightarrow$ permutation (1 6 3)(4 5)
- Page 182 line 1* _____ 17 Mar 1998
 A. De Moivre $\wedge \rightarrow$ A. de Moivre
- Page 182** line 4 _____ 18 Sep 2002
 W. A. Whitworth $\wedge \rightarrow$ I. Todhunter in his *Algebra* (second edition, 1860), §762,
 and by W. A. Whitworth
- **Page 182** the rating of exercise 6 _____ 24 Oct 2005
M23 $\wedge \rightarrow$ *M28*

Page 182 line 1 of exercise 11 _____ 02 Oct 1997

$\pi^{-1} \rightsquigarrow \pi^{-}$

►Page 182 line 4 of exercise 12 _____ 04 Aug 1997

$n \times n$ matrix $(b_{ij}) \rightsquigarrow n \times m$ matrix (b_{ij})

►Page 183 line 12 of exercise 22 _____ 09 Sep 1998

$P(n; k_1, k_2, k_3, \dots) \rightsquigarrow P(n; k_1, k_2, k_3, \dots)$

Page 184 in exercise 29 _____ 14 Oct 2008

lines 2–3: the “doubling” $\dots \bmod 2n + 1 \rightsquigarrow$ the “perfect shuffle” permutation of $\{1, 2, \dots, 2n\}$, which takes $(1, 2, \dots, 2n)$ into $(2, 4, \dots, 2n, 1, 3, \dots, 2n-1)$

line 4: the doubling permutation \rightsquigarrow the perfect shuffle

►Page 184 line 2 of exercise 30 _____ 01 Jan 2006

$(2^{d-1} - 1)(2n + 1)/(2^d - 1) \rightsquigarrow (2^d - 1)(2n + 1)/(2^{d+1} - 1)$

►Page 184 in exercise 31 _____ 14 Oct 2008

line 1: $HM33 \rightsquigarrow HM38$

line 4: fixed $m \rightsquigarrow$ fixed $m > 1$

line 5: $(m^{k+1} - (m-1)^k) \rightsquigarrow (m^{k+1} - (m-1)^k)$

Page 185 line 3 _____ 06 Aug 2001

this is the permutation on $\{0, 1, \dots, m+n-1\}$ that takes k into $(k+m) \bmod (m+n)$

\rightsquigarrow each element x_k should be replaced by $x_{p(k)}$ for $0 \leq k < m+n$, where $p(k) = (k+m) \bmod (m+n)$.

Page 185 new exercise _____ 29 Jan 2010

37. [M26] (*Even permutations.*) Let π be a permutation of $\{1, \dots, n\}$. Prove that π can be written as the product of an even number of 2-cycles if and only if π can be written as the product of exactly two n -cycles.

Page 186 line 14 _____ 17 Sep 2005

by having less space \rightsquigarrow by occupying less space

►Page 190 in (10) _____ 07 Jan 1999

EXIT C \rightsquigarrow EXITC

Page 193 lines 4 and 5 of exercise 6 _____ 21 Aug 2001

the time to execute the subroutine will be somewhat longer. \rightsquigarrow a subroutine naturally consumes more time and space than a hardware instruction does.

►Page 194 line –11 _____ 19 Dec 2001

stores J \rightsquigarrow stores rJ


Page 195 in (3) _____ 13 Jun 1999

The spaces between three-character groups should be wider.

Page 196 line –6 _____ 14 Jun 1999

three-digit groups \rightsquigarrow three-character groups

- **Page 198** line 17 from the bottom _____ 09 Jul 1997
 UNIX^{Exit conditions (when} users \rightsquigarrow UNIX[©] users
- **Page 198** line 15 from the bottom _____ 22 Jul 1997
 A, B, and C \rightsquigarrow B, C, and D
- Page 202 line 22 _____ 07 Aug 1997
 TeX: \rightsquigarrow T_EX:
- Page 202 line 25 _____ 11 Jul 1997
 PostScript[©] [Adobe \rightsquigarrow PostScript[©] [Adobe
- Page 202 line -3 _____ 08 Nov 2005
 simulator (or sometimes an *emulator*) \rightsquigarrow *simulator* (or sometimes an *emulator*)
- **Page 204** line 009 of the program _____ 14 Feb 1999
 see line 033 \rightsquigarrow (see line 033)
- **Page 205** line 042 of the program _____ 03 Sep 2008
 SPEC(1) \rightsquigarrow SPEC(10)
- Page 208 lines 161 and 164 of the program _____ 19 Dec 2001
 Let rI1 indicate the A register. \rightsquigarrow rI1 \leftarrow index of simulated rA.
- **Page 211** line 11 _____ 30 Dec 1997
 as does ENTA -5,1” \rightsquigarrow as does “ENTA -5,1”
- **Page 211** line -5 in 3rd, 4th, 5th printings _____ 16 Dec 1998
 1.3.1-?? \rightsquigarrow 1.3.1-26
- **Page 213** line 28 of the program _____ 24 Dec 2001
 JXL \rightsquigarrow JXNP
- **Page 216** line 7 _____ 02 Aug 1997
 52-53. \rightsquigarrow 52-53).
- Page 223 replacement for Fig. 25 _____ 08 Aug 1997



- Page 225 exercise 1 _____ 19 Dec 2001
 Would \rightsquigarrow (a) Would
 What if \rightsquigarrow (b) What if

- **Page 225** line −5 _____ 14 Feb 1999
 necessary \rightsquigarrow necessary.
- Page 230** line 12 _____ 27 Apr 2007
 Illiac I \rightsquigarrow ILLIAC I
- Page 230** line 16 _____ 07 Dec 2002
Nat. Conf. (1952) \rightsquigarrow (Toronto: 1952)
- Page 230** line −4 _____ 05 Feb 2002
 difficult. \rightsquigarrow difficult. See also the trace routine for IBM’s System/360 architecture, presented in *A Compiler Generator* by W. M. McKeeman, J. J. Horning, and D. B. Wortman (Prentice–Hall, 1970), 305–363.
- Page 231** line −21 _____ 19 Jun 2003
Nat. Conf. \rightsquigarrow *Nat. Meeting* **13**
- Page 231** lines −12 through −9 of the final paragraph _____ 29 Nov 2009
 X-1 computer . . . paper tape \rightsquigarrow X1 computer [see *Comp. J.* **2** (1959), 39–43]. Dijkstra’s doctoral thesis, “Communication with an Automatic Computer” (1959), discussed primitive synchronization operations by which users could create long chains of buffers with respect to paper tape
- **Page 231** quotation at bottom _____ 15 Jul 1997
 line −3: About 1000 \rightsquigarrow About 1,000
 line −2: the problems new envisioned. \rightsquigarrow problems now envisioned.
- **Page 233** line 19 _____ 28 Dec 1997
 JAVA \rightsquigarrow Java
- **Page 234** replacement for (3) re the 10 of clubs _____ 10 Aug 1997
- TOP \longrightarrow

+	0	2	2	•
+	□	□	2	□ D

\longrightarrow

+	0	4	3	•
+	□	□	3	□ S

\longrightarrow

+	1	1	10	•
+	□	1	0	□ C

\equiv (3)
- **Page 242** line 4 of exercise 2 _____ 21 Jul 1997
 require cards \rightsquigarrow require cars
- **Page 242** exercise 3 _____ 05 Jan 1998
 (a) through (h) \rightsquigarrow (i) through (viii)
 [In the first and second printings, make similar changes to exercise 2.]
- **Page 245** line 14 _____ 17 Jun 1998
 (6)and (7) \rightsquigarrow (6) and (7)
- Page 247 line 19* _____ 29 Oct 2006
 to be manipulated \rightsquigarrow to be treated
- **Page 247** line −5 _____ 06 Jan 2011
 $\leq \text{TOP}[i]$ \rightsquigarrow $< \text{TOP}[i]$

Page 248 line -6 _____ 20 Feb 2009
 $\text{NODE}(\text{TOP}[i]) \rightsquigarrow \text{CONTENTS}(\text{TOP}[i])$

►Page 249 line -2 _____ 06 Mar 1998
 $\text{NEWBASE}[j] > \text{NEWBASE}[j] \rightsquigarrow \text{NEWBASE}[j] > \text{BASE}[j]$

►Page 252 last line of exercise 5 _____ 13 Aug 1997
 exercise 4(iii) \rightsquigarrow exercise 4(c)

Page 253 lines 7 and 9 of exercise 12 _____ 18 Dec 1999
 $\max(k_1, k_2) \rightsquigarrow \max(k_1, k_2)$

►Page 256 replacement for line 26 _____ 08 Nov 1997

INFO	LINK
------	------

. (3)

Page 258 line -5 _____ 13 Aug 1997
 17 cycles, compared to 12 cycles \rightsquigarrow 17 units of time, compared to 12 units

Page 260 replacement for line -7 _____ 03 May 2004
 an empty queue is represented by $F = \Lambda$ and $R = \text{LOC}(F)$.

►Page 260 line 4 _____ 11 Aug 1997
 handle empty lists \rightsquigarrow to handle empty lists

Page 264 lines 1, 3, 6, 12 _____ 15 Aug 1997
 pairs \rightsquigarrow relations

Page 265 line 1 _____ 15 Aug 1997
 inputs pairs of relations \rightsquigarrow inputs a sequence of relations

Page 265 formerly line 3, now line 4 _____ 15 Aug 1997
 in linear order \rightsquigarrow in a linear order

Page 266 line 22 _____ 03 May 2004
 directly to Algorithm T. \rightsquigarrow directly to Algorithm T but with slight changes for efficiency.

►Page 266 line -20 _____ 25 May 1999
 $\text{COUNT}[j] \rightsquigarrow \text{COUNT}[k]$

►Page 268 line -2 _____ 10 Aug 1997
 $\text{UNDERFLOW} \rightsquigarrow \text{UNDERFLOW};$

►Page 270 line 2 of exercise 12 _____ 09 Mar 1998
 Given two \rightsquigarrow Give two

Page 270 line 2 of exercise 15 _____ 15 Aug 1997
 irredundant pairs of relations \rightsquigarrow irredundant relations

Page 273 line -3 of exercise 28 _____ 26 Jul 1997
 E. Lucas \rightsquigarrow É. Lucas

- Page 273** line -1 of exercise 28 _____ 08 Jul 2005
Winning Ways 2 (Academic Press, 1982) \rightsquigarrow *Winning Ways 3* (A. K. Peters, 2003)
- **Page 274** line 12 _____ 17 Nov 1997
 $\text{INFO}(P) \leftarrow Y'' \rightsquigarrow \text{INFO}(P) \leftarrow Y$
- **Page 275** second line after (4) _____ 12 Aug 1997
 don't have a \rightsquigarrow there is no
- Page 277 lines 2 and 3 of step M2* _____ 06 Dec 1999
 $\text{"if } ABC(P) < 0 \text{ then } -1, \text{ otherwise } ABC(P) + ABC(M)\text{"} \rightsquigarrow \text{"(if } ABC(P) < 0 \text{ then } -1, \text{ otherwise } ABC(P) + ABC(M)\text{"}$
- **Page 279** exercise 12 _____ 19 Dec 2001
 Algorithm A \rightsquigarrow Program A
- Page 281 lines 9 and 10* _____ 03 May 2004
 $NODE(X) \rightsquigarrow NODE(X)$ and $X \rightsquigarrow X$
- Page 285 step E3* _____ 20 Aug 1997
 $[Open\ door.] \rightsquigarrow [Open\ doors.]$
- Page 285 step E5* _____ 20 Aug 1997
 $[Close\ door.] \rightsquigarrow [Close\ doors.]$
- Page 287 step D2* _____ 20 Aug 1997
 $[Should\ door\ open?] \rightsquigarrow [Should\ doors\ open?]$
- Page 290 lines 047 and 049* _____ 20 Aug 1997
 Indicates door open \rightsquigarrow Indicates doors open
- Page 292 program line 098* _____ 25 Jan 2003
 Compute IN, OUT, \rightsquigarrow Set INFLOOR, OUTFLOOR,
- **Page 292** program lines 103 and 105 _____ 08 Jul 1998
 $POOLMAX \rightsquigarrow POOLMAX(0:2)$
- Page 293 line -14* _____ 24 Dec 2001
 coroutine E \rightsquigarrow Coroutine E
- Page 294 program line 201* _____ 20 Aug 1997
E3. Open door. \rightsquigarrow E3. Open doors.
- Page 295 program line 237* _____ 20 Aug 1997
E5. Close door. \rightsquigarrow E5. Close doors.
- **Page 296** line 7 _____ 08 Jul 1998
 $CON \rightsquigarrow NOP$
- **Page 296** line 20 _____ 14 Feb 1999
 by performed \rightsquigarrow be performed

Page 296 line 22 _____ 08 Oct 2005
quasi-parallel \rightsquigarrow *quasiparallel*

►Page 296 line -4 _____ 31 Aug 1998
 216–218 \rightsquigarrow 217–219

Page 298 last line of exercise 11 _____ 14 Feb 1999
 on the same step \rightsquigarrow in the same step

Page 299 line 6 _____ 09 Jun 2004
 an array \rightsquigarrow an array like (1)

►Page 301 replacement for Fig. 13 _____ 21 Jul 1997

	MALE	FEMALE	A21	A22	A23	A24	A28	BLUE	BROWN	GREEN	HAZEL	BLOND	RED	DARK	
PERSON [6]		•		•					•					•	Female, age 21, brown eyes, dark hair
PERSON [5]	•					•			•					•	Male, age 24, brown eyes, dark hair
PERSON [4]		•		•						•		•			Female, age 22, green eyes, blonde hair
PERSON [3]	•						•				•		•		Male, age 28, hazel eyes, blond hair
PERSON [2]		•		•				•					•		Female, age 22, blue eyes, red hair
PERSON [1]		•	•					•				•			Female, age 21, blue eyes, blonde hair
	⊥	⊥	⊥	⊥	⊥	⊥	⊥	⊥	⊥	⊥	⊥	⊥	⊥	⊥	

Page 305 last line before the exercises _____ 06 Jan 2011
 in Chapter 7 \rightsquigarrow in Chapter 7 (for example, Algorithm 7B)

►Page 305 line 1 of exercise 2 _____ 01 Feb 2004
 Formula (6) has \rightsquigarrow Formulas (5) and (6) have

►Page 305 line 2 of exercise 2 _____ 24 Dec 2001
 $l_r \leq l \leq u_r \rightsquigarrow l_r \leq l_r \leq u_r$

►Page 310 among the sons of Japheth _____ 30 Jul 1997
 Meschech \rightsquigarrow Meshech

Page 311 line 3 _____ 14 Mar 2004
 was being \rightsquigarrow was first being

Page 311 lines 26 and 27 _____ 05 Dec 2000
 represents, not a person ... so-and-so." \rightsquigarrow represents “a person in the role of mother or father of so-and-so,” not simply a person as an individual.

Page 311 near the bottom _____ 29 Sep 2010
 the ancestors of G are E , C , and A . \rightsquigarrow the ancestors of G are E , C , and A . Sometimes, especially when talking about “nearest common ancestors,” we consider a node to be an ancestor of itself (and a descendant of itself); the *inclusive ancestors* of G are G , E , C , and A , while its *proper ancestors* are just E , C , and A .

► **Page 313** line 1 after Fig. 21 _____ 29 Nov 1997

looks very much the \searrow looks very much like the

► **Page 313** line -3 _____ 19 Dec 2001

parent($F[a, b, c, d]$) \searrow the parent of $F[a, b, c, d]$

Page 315 line 8 before (3) _____ 28 Sep 2002

list structure \searrow list structure

Page 316 line 9 _____ 16 Feb 2003

lower case letters \searrow lowercase letters

Page 317 exercise 17 _____ 13 Feb 2001

If F is \searrow If Z stands for

$F[1, 2, 2]$ \searrow $Z[1, 2, 2]$

Page 317 line 2 of exercise 19 _____ 20 Aug 1997


this list? \searrow this List?

Page 318 exercise 22 _____ 13 Oct 1997

line 1: A_2, \dots \searrow A_2, \dots, A_n, \dots

line 2: $\sqrt{2}$ to 1. \searrow $\sqrt{2}$ to 1 and whose areas are 2^{-n} square meters.

► **Page 322** replacement for (7), has arrow $E \rightarrow G$ _____ 15 Jul 1997



► **Page 328** line -17 _____ 31 Aug 1998

u'_1, u'_2, \dots, u'_n \searrow u'_1, u'_2, \dots, u'_n

► **Page 328** line -9 _____ 24 Aug 1997

contents of \searrow complements of

► **Page 329** equation (15) _____ 07 Jul 1998


$f(u_{l+1})$ \searrow $f(u_{n_l+1})$

Page 331 new wording for exercises 7 and 8 _____ 04 Dec 2005

7. [22] Show that if we are given the preorder and the inorder of the nodes of a binary tree, the binary tree structure may be constructed. (Assume that the nodes are distinct.) Does the same result hold true if we are given the preorder and postorder, instead of preorder and inorder? Or if we are given the inorder and postorder?

8. [20] Find all binary trees whose nodes appear in exactly the same sequence in both (a) preorder and inorder; (b) preorder and postorder; (c) inorder and postorder. (As in the previous exercise, we assume that the nodes have distinct labels.)

- **Page 333** line 2 of exercise 31 _____ 21 Jun 2006
the tree node $\wedge \rightarrow$ the tree nodes
- Page 335 line 1* _____ 29 Nov 1999
 $45^\circ \wedge \rightarrow 45^\circ$ clockwise
- Page 337** line 9 _____ 12 Mar 2004
tree, $\wedge \rightarrow$ tree or forest,
- **Page 342** line 5 _____ 26 Dec 2002
 $rI1 \equiv P \wedge rI2 \equiv P$
- **Page 345** line 180 of the program _____ 01 Feb 2004
COPY2 \wedge COPYP2
- **Page 346** line 1 of exercise 7 _____ 21 Jul 1997
as a , \wedge as a partial ordering,
- Page 348** line 1 of exercise 20 _____ 29 Sep 2010
an ancestor \wedge a proper ancestor
- **Page 348** line 15 from the bottom _____ 15 Jul 1997
LLINK in tree There are \wedge There are
- **Page 349** replacement for (1) _____ 08 Aug 1997
 $(A(B, C(K)), D(E(H), F(J), G))$ (1)
- Page 351 line -2* _____ 29 Aug 1997
 $P \leftarrow P + 1 \wedge P \leftarrow P + c$
- **Page 352** lower middle _____ 26 Dec 2002
line -14: from 6 bytes to 3 \wedge from 5 bytes to 3
line -12: the latter has 60 \wedge the latter has 50
- **Page 353** replacement for the main part of Fig. 26 _____ 21 Aug 1997



- Page 354 line 1* _____ 29 Aug 1997
pairs of equivalences \wedge pairs of equivalent elements

► **Page 355** line 7 _____ 29 Aug 1997

the roots to two trees \rightsquigarrow the roots of two trees

Page 360 line 2 of exercise 8 _____ 29 Aug 1997

equivalences, \rightsquigarrow equivalent elements,

Page 360 line 2 of exercise 9 _____ 29 Aug 1997

pairs of equivalences \rightsquigarrow equivalences

Page 360 line 3 of exercise 11 _____ 29 Aug 1997

pairs of relations \rightsquigarrow a set of relations

► **Page 360** line 10 of exercise 11 _____ 25 Jul 1997

ARRAY X[0:10] \rightsquigarrow ARRAY X[0:10]

Page 363 updated terminology for lines 5–9 of §2.3.4.1 _____ 25 Jul 2008

(V_0, V_1, \dots, V_n) is a *path* ... vertex to itself. \rightsquigarrow (V_0, V_1, \dots, V_n) is a *walk* of length n from V to V' if $V = V_0$, V_k is adjacent to V_{k+1} for $0 \leq k < n$, and $V_n = V'$. The walk is a *path* if vertices V_0, V_1, \dots, V_n are distinct; it is a *cycle* if V_0 through V_{n-1} are distinct, $V_n = V_0$, and $n \geq 3$. Sometimes we are less precise, and refer to a cycle as “a path from a vertex to itself.” We often speak of a “simple path” to emphasize the fact that we’re talking about a path instead of an arbitrary walk. A graph is *connected* if there is a path between any two vertices of the graph.

Page 368 lines –21, –7, –2 _____ 25 Jul 2008

path \rightsquigarrow walk

Page 369 lines 9, –14, –12 _____ 25 Jul 2008

path \rightsquigarrow walk

Page 364 line 6 _____ 02 Sep 1997

edge $V_{k-1}V_k$ \rightsquigarrow edge $V_{k-1} - V_k$

Page 364 line 14 _____ 02 Sep 1997

For we take an arbitrary \rightsquigarrow This follows because we can find some

Page 364 lines 18 and 20 _____ 01 Jun 2006

a tree \rightsquigarrow a free tree


Page 366 line 1 below the caption to Fig. 32 _____ 01 Jun 2006

reach a subtree \rightsquigarrow reach a free subtree

► **Page 367** line –2 _____ 14 May 1999

for which e_j \rightsquigarrow for which E_j

Page 370 replacement for line –16 _____ 22 Jun 2004



► **Page 371** line 4 of exercise 10 _____ 02 Dec 1998
to that there \curvearrowright so that there

Page 372 lines 19–23 _____ 25 Jul 2008
is an *oriented path* ... vertex to itself. \curvearrowright is an *oriented walk* of length n from V to V' if $V = \text{init}(e_1)$, $V' = \text{fin}(e_n)$, and $\text{fin}(e_k) = \text{init}(e_{k+1})$ for $1 \leq k < n$. An oriented walk (e_1, e_2, \dots, e_n) is called *simple* if $\text{init}(e_1), \dots, \text{init}(e_n)$ are distinct and $\text{fin}(e_1), \dots, \text{fin}(e_n)$ are distinct; such a walk is an *oriented cycle* if $\text{fin}(e_n) = \text{init}(e_1)$, otherwise it's an *oriented path*.

Page 373 lines 18–25 _____ 02 Sep 1997
Furthermore, if ... an oriented \curvearrowright Furthermore, it has no cycles. For if (V_0, V_1, \dots, V_n) is an undirected cycle ... to the cycle

$$(V_1, V_0, V_{n-1}, \dots, V_1),$$


because $V_n = V_0$. Therefore an *oriented*

Page 374 and the next few pages too _____ 15 Jul 2004
[Starting with the 17th printing, the term ‘Eulerian circuit’ was changed to ‘Eulerian trail’, and ‘Hamiltonian circuit’ was changed to ‘Hamiltonian cycle’, in accordance with what has now become established usage. About thirty such changes were made, affecting pages 374, 375, 376, 379, 380, 581, and 584.]

Page 374 and subsequent pages too _____ 25 Jul 2008
[Starting with the 25th printing, the term ‘walk’ is often used formally instead of ‘path’, according to the definition now given on page 372, in order to conform more carefully to trends in the graph theory literature. In particular, ‘path’ should become ‘walk’ on page 374, line –24; page 375, lines –26, –24, –19, –18; page 376, lines 5, 17, –7; page 380, lines 18, –3; page 381, lines 2, 4, –24, –19, –15, –13, –11, –9; page 578, lines 4, 5; page 580, lines 1 and 2 of answer 1; page 584, lines 13 and 29.]

Page 374 lines –7 and –6 _____ 25 Jul 2008
oriented path, simple oriented path \curvearrowright oriented walk, oriented path

Page 375 replacement for Fig. 36 _____ 31 Dec 1997



Page 376 lines 4, 8, 10, and 11 _____ 01 Jan 2006
 $\text{init } \curvearrowright \text{ init}$ (thrice) and $\text{fin } \curvearrowright \text{ fin}$ (once)

Page 376 line 3 of exercise 4 _____ 22 Sep 1999
all edges $e \curvearrowright$ all arcs e

► **Page 377** line -2 of exercise 10 _____ 14 May 1999
the F table \rightsquigarrow the P table

Page 377 line 2 of exercise 14 _____ 25 Jul 2008
all paths \rightsquigarrow all oriented walks

Page 379 in exercise 21 _____ 06 Sep 2001
line 4: $n + 1$ vertices V_0, V_1, \dots, V_n \rightsquigarrow n vertices V_0, V_1, \dots, V_{n-1}
line 5: $(n + 1)m$ arcs \rightsquigarrow mn arcs
line 6: $(n + 1)m$ vertices \rightsquigarrow mn vertices
line 10: $m^{(n+1)(m-1)}$ \rightsquigarrow $m^{(m-1)n}$

Page 379 line 5 of exercise 21 _____ 26 Dec 2002
the graph \rightsquigarrow the digraph

► **Page 381** replacement for the bottom line _____ 01 Jan 2006

$$x_j = \sum_{\substack{\text{init}(e)=V_i \\ \text{fin}(e)=V_j}} p(e) x_i, \quad \text{for } 1 \leq j \leq n.$$

Page 387 lines -11, -10, -9 _____ 25 Jan 2003

The weight of $Y_1 \dots$ we have \rightsquigarrow If Y is any node in the Y_1 subtree, its weight must be at least $n - s_1 = 1 + s_2 + \dots + s_k$, since when Y is the assumed root there are at least $n - s_1$ vertices in its subtree containing X . Thus if Y is a centroid we have

Page 387 line -8 _____ 16 May 1999
weight(X) \rightsquigarrow weight(X)

Page 395 line 2 after (30) _____ 26 May 2000
some of this function's \rightsquigarrow this function's history and some of its

► **Page 395** line 11 after (30) _____ 19 Mar 1998
1997 \rightsquigarrow 1998

Page 395 line 2 of exercise 1 _____ 14 Aug 1997
 $\frac{1}{2}A(z^2)$ \rightsquigarrow $\frac{1}{2}A(z^2)$

Page 396 line 9 of exercise 4 _____ 14 Aug 1997
 $\frac{1}{2}A(z^2)$ \rightsquigarrow $\frac{1}{2}A(z^2)$

Page 397 line 5 _____ 14 Feb 1999
prove there \rightsquigarrow prove that there

► **Page 398** line 2 of exercise 25 _____ 27 Nov 1999
 $r(n, n - 1) = n^{n-1}$ \rightsquigarrow $r(n, n - 1) = n^{n-2}$

► **Page 399** line 3 _____ 27 Nov 1999
 $(x + \epsilon_1 z_1 + \dots + \epsilon_n z_n)^{\epsilon_1 + \dots + \epsilon_n - 1}$ \rightsquigarrow $(x + \epsilon_1 z_1 + \dots + \epsilon_n z_n)^{\epsilon_1 + \dots + \epsilon_n - 1}$

► **Page 399** line 1 of exercise 32 _____ 11 Dec 2002
 (1940) \rightsquigarrow (1941)

Page 399 line 2 of exercise 32 _____ 08 Nov 2005
 7–12). \rightsquigarrow 7–12.)

► **Page 402** line –4 _____ 11 Sep 1998
 (1951) \rightsquigarrow (1952)

Page 403 replacement for (10) _____ 14 Sep 1997



Page 404 line 4 and 5 _____ 22 Oct 2008
 of such a tree is minimized if and only if the tree \rightsquigarrow of a tree for the weights w_1 ,
 \dots , w_m that contains (10) as a subtree is minimized if and only if that tree

► **Page 406** line 9 _____ 19 Aug 2002
Messenger of Math. **58** (1939) \rightsquigarrow *Messenger of Math.* **58** (1929)

Page 406 line –9 _____ 24 Mar 2008
 König \rightsquigarrow König [to agree with the German spelling in the works cited]

Page 407 lines 2–4 _____ 29 Jan 2005
 by J. von Segner ... it was the subject \rightsquigarrow by L. Euler, who mentioned his
 results in a letter to C. Goldbach on 4 September 1751 [see J. von Segner and L.
 Euler, *Novi Commentarii Academiæ Scientiarum Petropolitanæ* **7** (1758–1759),
 summary 13–15, 203–210]. Euler’s problem was the subject

Page 407 lines 14 and 15 _____ 29 Jan 1999
 to appear \rightsquigarrow 1999

► **Page 411** line –6 _____ 28 Aug 1997
 contain a TYPE field \rightsquigarrow contains a TYPE field


► **Page 415** line 3 of step A2 _____ 17 Sep 1997
 not at atom \rightsquigarrow not an atom

► **Page 415** line 5 of step A2 _____ 17 Sep 1997
 $K \leftarrow \min(K1, \text{BLINK}(K)) \rightsquigarrow K1 \leftarrow \min(K1, \text{BLINK}(K))$

Page 415 line 22 _____ 03 May 2004
 “NODE(Λ)” \rightsquigarrow “NODE(Λ)”

Page 418 line 3 of step E4 _____ 17 Sep 1997
 the list \rightsquigarrow the List

► **Page 421** line 11 _____ 17 Sep 1997
 garbage is \rightsquigarrow total memory is

- **Page 421** line −15 _____ 01 Jan 2002
 499–506 \rightsquigarrow 499–507
- **Page 423** line −4 of exercise 11 _____ 08 Sep 1997
 $(a: (b: D), b, a: E)) \rightsquigarrow (a: (b: D), b, a: E)$
- Page 426* line 8 _____ 26 Oct 1998
 MONTH, DAY and YEAR \rightsquigarrow MONTH, DAY, and YEAR
- **Page 426** line 9 _____ 26 Dec 2002
 DAY, MONTH, YEAR \rightsquigarrow MONTH, DAY, and YEAR
- **Page 428** line −9 _____ 20 Sep 1997
 memory location \rightsquigarrow memory allocation
- **Page 430** line 6 _____ 08 May 2000
 set P = PREV(P) \rightsquigarrow set P \leftarrow PREV(P)
- **Page 430** line −9 _____ 15 Oct 2007
 $0 \leq j < n; \rightsquigarrow 0 \leq j < n - 1$ and that α and β are parents of A_{n-1} ;
- Page 431* line 16 _____ 31 Jan 2008
 qualifications \rightsquigarrow complete qualifications
- **Page 432** line −11 _____ 20 Sep 1997
 “NAME(S) = P_k ”, NAME(P) = NAME(Q)” \rightsquigarrow “NAME(S) = P_k ” and “NAME(P) = NAME(Q)”
- **Page 433** bottom line _____ 22 Jan 1999
 $A_{i+1} \rightsquigarrow A_{j+1}$
- Page 435* line −11 _____ 21 Aug 2007
 for a while \rightsquigarrow for awhile
- Page 436* subtle change in Fig. 42 at location 27198 _____ 20 Sep 1997
- 
- Page 443* line 1 _____ 10 Nov 2002
 present in all blocks, \rightsquigarrow present in all blocks, and which must not be tampered with by the users who reserve blocks,

Page 443 step R2 _____ 06 Nov 2006

$L \leftarrow \text{AVAILF}[j], P \leftarrow \text{LINKF}(L), \text{AVAILF}[j] \leftarrow P, \text{LINKB}(P) \leftarrow \text{LOC}(\text{AVAIL}[j]) \rightsquigarrow$
 $L \leftarrow \text{AVAILB}[j], P \leftarrow \text{LINKB}(L), \text{AVAILB}[j] \leftarrow P, \text{LINKF}(P) \leftarrow \text{LOC}(\text{AVAIL}[j])$

Page 444 line 16 _____ 20 Sep 1997

deleted \rightsquigarrow freed

Page 444 line 22 _____ 03 May 2004

$K \rightsquigarrow K$

►Page 444 bottom line _____ 15 Oct 2007

$1 - p \rightsquigarrow -1 + p.$

►Page 445 line 15 _____ 25 Mar 2001

$\text{BIT } 20 \rightsquigarrow \text{BIT } 20$

Page 445 lines -15 through -9 _____ 06 Jan 2011

[Reformat steps P1-P4 to make them like steps of a normal algorithm.]


Page 447 line after the Figure _____ 19 Dec 2001

time distribution chosen randomly between 1 and 100 \rightsquigarrow times distributed uniformly in $\{1, \dots, 100\}$

Page 447 line -7 _____ 20 Sep 1997

about four thirds \rightsquigarrow more than four thirds

►Page 448 replacement for Fig. 44 _____ 22 Jul 1997



►Page 450 line -15 _____ 28 Sep 2002

$38, \rightsquigarrow 38$

- Page 450 line -15 _____ 11 Nov 1998
 “distributed fit” method \rightsquigarrow “distributed-fit method”
- Page 455 refinements to exercise 32 _____ 09 Nov 2002
 line 1: $HM47 \rightsquigarrow HM46$
 line 2: $1 \leq k < n \rightsquigarrow 0 \leq k < n$
 line 4: $f(t_1 - (n - 1)) \rightsquigarrow f(t_0 - n)$
- Page 456 line 2 of exercise 39 _____ 10 Oct 1997
 $\lim_{m \rightarrow \infty} \rightsquigarrow \lim_{n \rightarrow \infty}$
- Page 456 line 1 of exercise 40 _____ 06 Nov 1997
 $\lim_{n \rightarrow \infty} \rightsquigarrow \lim_{m \rightarrow \infty}$
- Page 457 line -7 _____ 04 Dec 2000
 insertion and deletions \rightsquigarrow insertion and deletion
- Page 466 line 8 _____ 08 Oct 2010
Annals of Mathematics \rightsquigarrow *Annals of Mathematics* (2)
- Page 467 changes to answer 9 _____ 01 Jan 2002
 line 2: taking Ω_2 into Ω_1 \rightsquigarrow
 replacement for lines 4 and 5: a) If x is in I_1 then $h(g(x)) = x$.
 new line to follow line 7: c) If q is in Q_2 then $h(q)$ is in Ω_1 if and only if q is in Ω_2 .
- Page 468 line 2 of answer 2 _____ 12 Oct 2008
 $a^{-1} \rightsquigarrow a^{(n-1)-1} = a^{-1}$
- Page 468 opening lines of answer 8 _____ 28 Mar 2005
 (a) We will ... an inductive proof \rightsquigarrow (a) We must show that $(n^2 - n + 1) + (n^2 - n + 3) + \dots + (n^2 + n - 1)$ equals n^3 . And indeed, the sum is $n(n^2 - n) + (1 + 3 + \dots + (2n - 1)) = n^3 - n^2 + n^2$, by Eq. (2). But an inductive proof
- Page 468 line 1 of answer 10 _____ 06 Jan 1998
 $(1 + \frac{1}{10})^3 \rightsquigarrow (1 + 1/n)^3$
- Page 469 line 4 of answer 15(e) _____ 16 Dec 1997
 $(x_1, \dots, x_n) \prec (y_1, \dots, y_n) \rightsquigarrow (x_1, \dots, x_n) \prec (y_1, \dots, y_n)$
- Page 470 line 1 of answer 25 _____ 10 Oct 2001
 $z = 2^{-p} \lfloor 2^{p-k} x \rfloor \rightsquigarrow z = 2^{-p} \lfloor 2^{p-k} x \rfloor > 0$
- Page 470 line 3 of answer 28 _____ 03 Feb 2002
 $x \leftarrow 1 - \epsilon \rightsquigarrow x \leftarrow 1 - \epsilon - x$
- Page 471 line 6 _____ 13 Aug 1997
METAFONT: The Program \rightsquigarrow *METAFONT: The Program*
- Page 471 new answer following line 8 _____ 28 Jun 2003
 30. n .

Page 471 replacements for answers 1–3 _____ 04 Jan 2010

1. $-a_1$; and $a_2 + \cdots + a_1 = 0$. In general, sums with ‘ \cdots ’ are defined so that $(a_p + \cdots + a_q) + (a_{q+1} + \cdots + a_r) = a_p + \cdots + a_r$ for arbitrary integers p , q , and r .

2. $a_1 + a_2 + a_3$.

3. $\frac{1}{1} + \frac{1}{3} + \frac{1}{5} + \frac{1}{7} + \frac{1}{9} + \frac{1}{11}$; $\frac{1}{9} + \frac{1}{3} + \frac{1}{1} + \frac{1}{3} + \frac{1}{9}$. The rule for $p(j)$ is violated: In the first case $n^2 = 3$ occurs for no n , and in the second case $n^2 = 4$ occurs for *two* n . [See Eq. (18).]

Page 471 answer 19 _____ 01 May 2007

$$a_n - a_{m-1} \rightsquigarrow (a_n - a_{m-1})[m \leq n]$$

Page 472 replacement for lines 7–8 of answer 33 _____ 06 Jan 2009

where $P(x_j)$ is a polynomial of degree $n - 2$ in x_j whose coefficients are symmetric functions of $\{x_1, \dots, x_n\}$ that don’t depend on j . (See exercise 1.2.9–10.) We obtain the desired answer from the solutions for $r = 0, 1, \dots, n - 1$.

Page 473 replacement for lines 7 and 8 _____ 06 Jan 2009

$$\sum_{\substack{j_1 + \cdots + j_n = r - n + 1 \\ j_1, \dots, j_n \geq 0}} x_1^{j_1} \cdots x_n^{j_n} = \sum_{1 \leq j_1 \leq \cdots \leq j_{r-n+1} \leq n} x_{j_1} \cdots x_{j_{r-n+1}};$$

see Eq. 1.2.9–(33). [J. J. Sylvester, *Quart. J. Math.* **1** (1857), 141–152.]

Page 474 new copy for the beginning of answer 40 _____ 08 Jan 2005

[A. de Moivre, *The Doctrine of Chances*, 2nd edition (London: 1738), 197–199.] We have

► **Page 475** lines 10 and 11 _____ 15 Oct 2007

are Cauchy matrices \rightsquigarrow are determinants of Cauchy matrices
not Hilbert matrices \rightsquigarrow not determinants of Hilbert matrices

Page 475 last line of answer 45 _____ 13 Jan 2003

22. \rightsquigarrow 22; Cauchy, *Œuvres* (2) **12**, 173–182. [Beginning with the 24th printing, the latter reference was changed to: A. Cauchy, *Exercices d’analyse et de physique mathématique* **2** (1841), 151–159.]

► **Page 475** line 10 of answer 46 _____ 15 Dec 2001

$$\sum_{1 \leq k_1, \dots, k_m \leq m} \rightsquigarrow \sum_{1 \leq k_1, \dots, k_m \leq n}$$

► **Page 475** bottom line _____ 15 Oct 2007

$$n - 1 \rightsquigarrow n - 2$$

Page 476 line 3 of answer 7 _____ 13 May 2005

> 1 . $\rightsquigarrow > 1$; that is, if and only if $(-x) \bmod 1 + (-y) \bmod 1 < 1$.

► **Page 477** line 7 of answer 34 _____ 16 Oct 1997

i satisfied \rightsquigarrow is satisfied

► **Page 478** line –2 of answer 37 _____ 08 Nov 2005

$$0 \leq k \leq n \rightsquigarrow 0 < k \leq n$$

►Page 478 bottom line _____ 08 Mar 2002

$$e^{2\pi y} \rightsquigarrow e^{2\pi i y}$$

Page 479 displayed formula in answer 42 _____ 11 Aug 2005

$$k+1 \text{ is a power of } b \rightsquigarrow k+1 \text{ is a power of } b$$

►Page 480 line 2 of answer 13 _____ 08 Nov 2005

$$\text{Law 1.2.4D} \rightsquigarrow \text{Law 1.2.4B}$$

►Page 481 line 9 of answer 20 _____ 02 Dec 1997

$$\frac{1}{m} \int_0^m t^{x+1} e^{-t} \rightsquigarrow \frac{1}{m} \int_0^m t^{x+1} e^{-t} dt$$

►Page 483 line 1 of answer 25 _____ 21 Jan 2001

$$x^{\overline{m+n}} = x^{\overline{m}}(x-m)^{\overline{n}} \rightsquigarrow x^{\overline{m+n}} = x^{\overline{m}}(x+m)^{\overline{n}}$$

Page 484 last line of answer 10 _____ 09 Jan 2003

$$N. J. Fine \rightsquigarrow L. E. Dickson, \textit{Quart. J. Math.} \textbf{33} (1902), 383-384; N. J. Fine$$

►Page 485 replacement for answer 21 _____ 07 Jun 2003

21. The left-hand side is a polynomial of degree $\leq n$; the right-hand side is a polynomial of degree $m+n+1$. The polynomials agree at $n+1$ points, but that isn't enough to prove them equal. [In fact, the correct formula in general is

$$\sum_{k=0}^r \binom{r-k}{m} \binom{s+k}{n} = \binom{r+s+1}{m+n+1} - \sum_{k=0}^m \binom{r+1}{k} \binom{s}{m+n+1-k}$$

when m , n , and r are nonnegative integers.]

►Page 485 line 1 of answer 22 _____ 24 May 2000

$$\text{The } k\text{th term is} \rightsquigarrow \text{The } k\text{th term is } r \text{ times}$$

►Page 486 line 2 _____ 24 May 2000

$$tf(r-t-1, s+t, t, n-1) \rightsquigarrow tf(r+t-1, s-t, t, n-1)$$

Page 486 line 1 of answer 33 _____ 18 Jan 2005

[*Giornale* ... may therefore be \rightsquigarrow [*Mém. Acad. Roy. Sci.* (Paris, 1772), part 1, 492; C. Kramp, *Éléments d'Arithmétique Universelle* (Cologne: 1808), 359; *Giornale di Mat. Battaglini* **33** (1895), 179-182.] Since $x^{\overline{n}} = n! \binom{x+n-1}{n}$, the equation may be

Page 486 line -3 _____ 14 Jul 1997

$\sum_k \binom{n}{k} x(x-kz-1)^{\overline{k-1}} (y+kz)^{\overline{n-k}} \rightsquigarrow \sum_k \binom{n}{k} x(x-kz-1)^{\overline{k-1}} (y+kz)^{\overline{n-k}}$, an equivalent formula of Rothe [*Formulae de Serierum Reversione* (Leipzig: 1793), 18].

Page 487 line 9 of answer 38 _____ 11 Oct 2001

$$\text{formula.} \rightsquigarrow \text{formula. [See } \textit{Crelle} \textbf{11} (1834), 353-355.]$$

►Page 489 answer 52 _____ 26 Jan 1998

$$\text{line 4: } f(x, y, r) \rightsquigarrow f(x, y, n)$$

$$\text{line 5: } rf(x, y, n-1) \rightsquigarrow nf(x, y, n-1)$$

Page 489 replacement for answer 56 _____ 31 May 2002

56. 210 310 320 321 410 420 421 430 431 432 510 520 521 530 531 532 540 541 542 543 610. With a fixed, b and c run through the combinations of a things two at a time; with a and b fixed, c runs through the combinations of b things one at a time.

Similarly, we could express all numbers in the form $n = \binom{a}{4} + \binom{b}{3} + \binom{c}{2} + \binom{d}{1}$ with $a > b > c > d \geq 0$; the sequence begins 3210 4210 4310 4320 4321 5210 5310 5320 We can find the combinatorial representation by a “greedy” method, first choosing the largest possible a , then the largest possible b for $n - \binom{a}{4}$, etc. [Section 7.2.1.3 discusses further properties of this representation.]

Page 489 replacement for first three lines of answer 58 _____ 12 Jun 2003

58. [*Systematisches Lehrbuch der Arithmetik* (Leipzig: 1811), xxix.] Use induction and

$$\binom{n}{k}_q = \binom{n-1}{k}_q + \binom{n-1}{k-1}_q q^{n-k} = \binom{n-1}{k}_q q^k + \binom{n-1}{k-1}_q.$$

Therefore [F. Schweins, *Analysis* (Heidelberg: 1820), §151] the q -generalization of (21) is

Page 490 replacement for lines 5–7 _____ 12 Jun 2003

For further information, see G. Gasper and M. Rahman, *Basic Hypergeometric Series* (Cambridge Univ. Press, 1990). The q -nomial coefficients were introduced by Gauss in *Commentationes societatis regiae scientiarum Gottingensis recentiores* **1** (1808), 147–186; see also Cauchy [*Comptes Rendus Acad. Sci.* **17** (Paris, 1843), 523–531], Jacobi [*Crelle* **32** (1846), 197–204], and Heine [*Crelle* **34** (1847), 285–328].

► **Page 491** line 2 _____ 28 Jan 1998

$$q^{m-r+s-k)(n-k)} \searrow q^{(m-r+s-k)(n-k)}$$

Page 491 replacement for answer 66 _____ 08 Nov 2005

66. Let $X = \binom{x}{n}$, $\underline{X} = \binom{x}{n-1} = \frac{n}{x-n+1}X$, $\overline{X} = \binom{x}{n+1} = \frac{x-n}{n+1}X$, with similar notations for Y and Z . We may assume that $y > n-1$ is fixed, so that x is a function of z .

Let $F(z) = \overline{X} - \overline{Y} - \overline{Z}$, and suppose that $F(z) = 0$ for some $z > n-2$. We will prove that $F'(z) < 0$; therefore $z = y$ must be the only root $> n-2$, proving the second inequality. Since $F(z) = \frac{x-n}{n+1}(Y+Z) - \frac{y-n}{n+1}Y - \frac{z-n+1}{n}Z = 0$ and $x > y$ and $Y, Z > 0$, we must have $\frac{x-n}{n+1} < \frac{z-n+1}{n}$. Setting $X' = dX/dx$ and $Z' = dZ/dz = dX/dz$, we have

$$\frac{X'}{X} = \frac{1}{x} + \frac{1}{x-1} + \cdots + \frac{1}{x-n+1} > \frac{n}{n+1} \left(\frac{1}{z} + \cdots + \frac{1}{z-n+2} \right) = \frac{n}{n+1} \frac{Z'}{Z},$$

since $\frac{x-n+1}{n+1} < \frac{z-n+2}{n}$, \dots , $\frac{x-1}{n+1} < \frac{z}{n}$. Thus $dx/dz = Z'/X' < \frac{n+1}{n}(Z/X)$, and

$$F'(z) = \frac{X}{n+1} \frac{dx}{dz} + \frac{x-n}{n+1} Z' - \frac{Z}{n} - \frac{z-n+1}{n} Z' < \left(\frac{x-n}{n+1} - \frac{z-n+1}{n} \right) Z' < 0.$$

To prove the first inequality, we may assume that $n > 2$. Then if $\underline{X} = \underline{Y} + \underline{Z}$ for some $z > n-2$, the second inequality tells us that $z = y$.

References: L. Lovász, *Combinatorial Problems and Exercises*, Problem 13.31(a); R. M. Redheffer, *AMM* **103** (1996), 62–64.

Page 491 line –2 of that answer _____ 03 Nov 2008

Exercises, \searrow *Exercises* (1993),

► **Page 493** line 2 of answer 12 _____ 13 Feb 1998

$$\mathcal{F}(z) \rightsquigarrow \mathcal{G}(z)$$

Page 493 answer 14 _____ 25 Jun 2006

$$F_{m+1}F_{n-1} + (F_{m+2} + 1)F_n \rightsquigarrow F_{m+n+1} + F_n$$

Page 493 last line of answer 19 _____ 21 Jun 2007

$$\sin 72^\circ = \frac{1}{2}5^{1/4}\phi^{1/2}, \rightsquigarrow \sin 72^\circ = \frac{1}{2}5^{1/4}\phi^{1/2}. \text{ Another interesting angle is } \alpha = \arctan \phi = \frac{\pi}{4} + \frac{1}{2}\arctan \frac{1}{2}, \text{ for which we have } \sin \alpha = 5^{-1/4}\phi^{1/2}, \cos \alpha = 5^{-1/4}\phi^{-1/2}.$$

Page 494 replacement for line 4 _____ 26 Jan 2004

$$(n+1-x^n F_{n+1})/(2x+1).$$

Page 495 line 6 of answer 34 _____ 05 Feb 2005

182]; generalizations \rightsquigarrow 182]; but Section 7.2.1.7 points out that it was implicitly known in 14th-century India. Generalizations

Page 495 bottom line of answer 34 _____ 12 Oct 2006

exercise 5.4.2–10 \rightsquigarrow exercise 5.4.2–10 and in Section 7.1.3.

► **Page 495** line 7 of answer 35 _____ 16 Nov 1997

$$m - \phi^{2n}, \rightsquigarrow m - \phi^{-2n},$$

Page 495 line 8 of answer 35 _____ 16 Nov 1997

$$\phi^{k-1} + \dots + \phi^{k-2j+1} \rightsquigarrow \phi^{k-1} + \phi^{k-3} + \dots + \phi^{k-2j+1}$$

Page 495 line 7 of answer 36 _____ 23 Aug 2002

Jean Bernoulli \rightsquigarrow John Bernoulli

► **Page 496** line 2 of answer 5 _____ 01 Jan 2006

$$(n-1)! \rightsquigarrow \frac{(n-1)!}{k!}$$

Page 496 answer 10 _____ 18 Aug 2002

[Change the notation from ‘ a_m ’ to the more modern ‘ e_m ’, throughout this exercise and also in the answer to exercise 11 on page 497.]

► **Page 498** line 3 _____ 01 Jan 2006

$$f(x) \rightsquigarrow \lim_{x \rightarrow 1^-} f(x) = f(1)$$

► **Page 498** replacement for line 8 _____ 01 Jan 2006

$$\frac{i}{2} + \frac{i}{(\omega^{-p} - 1)} = \frac{i}{2} \left(\frac{1 + \omega^p}{1 - \omega^p} \right) = -\frac{i}{2} \left(\frac{\omega^{p/2} + \omega^{-p/2}}{\omega^{p/2} - \omega^{-p/2}} \right) = -\frac{1}{2} \cot \frac{p}{q} \pi.$$

► **Page 498** line 2 of answer 21 _____ 15 Apr 2002

$$E_1(x) \rightsquigarrow E_1(z)$$

► **Page 498** line 1 of answer 23 _____ 16 Dec 2001

When $m = 1$ \rightsquigarrow (a) When $m = 1$

► **Page 500** replacement for line 2 of answer 6 _____ 01 Aug 1997

$$\ln(q + pe^t) = \ln\left(1 + pt + \frac{pt^2}{2} + \frac{pt^3}{6} + \dots\right) = pt + p(1-p)\frac{t^2}{2} + p(1-p)(1-2p)\frac{t^3}{6} + \dots$$

- **Page 500** line 6 of answer 6 _____ 11 Jan 2002
 $H'(z) = e^t/(e^t - 1) - 1/t \rightsquigarrow H'(t) = e^t/(e^t - 1) - 1/t$
- **Page 500** line 1 of answer 8 _____ 30 Jul 1999
 $M^{\mathbb{Z}}/((M - n)! M^n) \rightsquigarrow M^{\mathbb{Z}}/M^n$
- **Page 501** last line of answer 13 _____ 01 Aug 1997
 n and x .] \rightsquigarrow n and x .
- Page 501* line 1 of answer 14 _____ 01 Aug 1997
 $e^{-itpn/\sqrt{pqn}}(q + pe^{it/\sqrt{pqn}})^n = (qe^{-itp/\sqrt{pqn}} + pe^{itq/\sqrt{pqn}})^n \rightsquigarrow$
 $e^{-itpn/\sqrt{pqn}}(q + pe^{it/\sqrt{pqn}})^n = (qe^{-itp/\sqrt{pqn}} + pe^{itq/\sqrt{pqn}})^n$
- **Page 501** line 2 of answer 15 _____ 25 Mar 2001
 $+ -t^2/2np \rightsquigarrow -t^2/(2np)$
- Page 502* line 1 of answer 21 _____ 10 Sep 1997
[There's slightly too much space before the word “The”.]
- **Page 502** line 2 of answer 21 _____ 01 Aug 1997
 $\Pr(X \geq n(p + \epsilon)) \rightsquigarrow \Pr(X \geq n(p + \epsilon))$
- **Page 502** line 3 of answer 21 _____ 05 Jul 1999
 $-\frac{1}{2q}\epsilon^2 \rightsquigarrow \epsilon - \frac{1}{2q}\epsilon^2$
- **Page 502** line 2 of answer 22 _____ 22 Jan 1999
493–509 \rightsquigarrow 493–507
- **Page 502** line 4 of answer 22 _____ 05 Jul 1999
 $\leq e^{\delta^2/2}$ when $\delta \leq 0$ and $\leq e^{\delta^2/3} \rightsquigarrow \leq e^{-\delta^2/2}$ when $\delta \leq 0$ and $\leq e^{-\delta^2/3}$
- Page 502* line 1 of answer 23 _____ 10 Sep 1997
[There's slightly too much space before the word “Setting”.]
- **Page 502** line 4 of answer 23 _____ 27 Oct 1998
 $f(\epsilon) \leq -\epsilon^2/(6pq) \rightsquigarrow$ that $f(\epsilon) \leq -\epsilon^2/(6pq)$ if $0 \leq \epsilon \leq p$.
- Page 503* answer 8 _____ 26 May 1998
the previous \rightsquigarrow the method of the previous
- **Page 503** line 1 of answer 9 _____ 05 May 2003
 $f(x) \rightsquigarrow f(z)$
- **Page 503** line 1 of answer 9 _____ 26 Feb 2008
 $e^{f(z)} \rightsquigarrow |e^{f(z)}|$
- **Page 503** line 2 of answer 12 _____ 30 Jul 1999
 $O(n^k/(n-1)!) \rightsquigarrow O(n^k/(k-1)!)$
- **Page 503** answer 1 _____ 01 Aug 1997
 $B_0 + B_1z + B_2z^2/2! + \dots) e^z \rightsquigarrow (B_0 + B_1z + B_2z^2/2! + \dots) e^z$

Page 504 lines 2 and 3 _____ 01 Jan 2006

$\ln(n-1)! \rightsquigarrow \ln(n-1)!$

►Page 504 line 5 of answer 7 _____ 20 May 2005

The number A is “Glaisher’s constant” 1.2824271... [Messenger \rightsquigarrow In these formulas, A is the “Kinkelin–Glaisher constant” 1.2824271... [Crelle **57** (1860), 122–158; Messenger

►Page 504 line 3 of answer 8 _____ 01 Jan 2006

$+ \sigma \rightsquigarrow + \frac{1}{2} \ln a + \sigma$

Page 504 line 4 of answer 8 _____ 01 Jan 2006

$\ln(cn^2)! - \ln(cn^2 - n)! - n \ln c - \ln n^2! + \ln(n^2 - n)! \rightsquigarrow$
 $\ln(cn^2)! - \ln(cn^2 - n)! - n \ln c - \ln n^2! + \ln(n^2 - n)!$

Page 504 line 1 of answer 9 _____ 01 Jan 2006

$\ln(2n)! \rightsquigarrow \ln(2n)! \quad \text{and} \quad \ln(n!)^2 \rightsquigarrow \ln(n!)^2$

►Page 504 line 3 of answer 9 _____ 24 Mar 1998

$O(n^{-3}) = \rightsquigarrow O(n^{-3}) =$

►Page 505 second display in answer 9 _____ 17 Oct 2007

$x^z \rightsquigarrow x^x$

►Page 506 lines 3 and 4 _____ 10 Aug 2008

$\sum_{k,m} \rightsquigarrow \sum_{k,m \geq 0}$ (twice)

Page 506 line −1 of answer 11 _____ 10 Aug 2008

$(-1)^n n! / e^n n^n \rightsquigarrow (-1)^n n! / (e^n n^n)$

►Page 507 line 4 of answer 20 _____ 01 Jan 2006

$\sum_{k=1}^{m-1} c_k (2u)^{k/2-1} \rightsquigarrow \sum_{k=1}^{m-1} k c_k (2u)^{k/2-1}$

Page 507 line 2 of answer 6 _____ 20 Apr 1998

into an index \rightsquigarrow into an index

►Page 507 line 2 of answer 7 _____ 28 Sep 2002

$|V| \rightsquigarrow |V|$

Page 508 line 4 of answer 14 _____ 25 Mar 2003

JSJ \rightsquigarrow STJ $\ast(0:0)$, STZ $\ast(0:0)$, and STZ $\ast(3:3)$; JSJ

Page 508 line 2 of answer 21 _____ 20 Apr 1998

only by jumping \rightsquigarrow only by jumping

►Page 509 line 3 _____ 03 May 1999

a sign \rightsquigarrow the sign

Page 509 line 18 _____ 28 Sep 1997

time = 54u. \rightsquigarrow time = 54u, not counting the HLT.

Page 509 lines -12 through -9 _____ 17 Nov 2007

(3495) $(-5)^{13}$ [This line needed only when $b > 65$.]

(3496) $(-4)^{13}$

\vdots

(3505) $(+5)^{13}$ [This line needed only when $b > 65$.] ■

►Page 510 line 03 of the program _____ 21 Nov 1998

$\sqcup C \sqcup 04 \searrow \sqcup C \sqcup 04$

►Page 510 line -2 _____ 30 Dec 1997

$\sqcup \sqcup \sqcup B. \searrow \sqcup J \sqcup B.$

►Page 512 right column, line 12 of the program _____ 25 Aug 2002

GOOD \searrow MEMORY

Page 512 right column, program lines 18-19 and 24-25 _____ 06 Dec 2008

CMPA =5=; JG BAD \searrow DECA 5; JAP BAD

CMPX =3999=; JLE GOOD \searrow DECX 3999; JXNP GOOD

Page 513 changes for consistency _____ 29 Oct 1997

line 1: *Solution 1*: \searrow *Solution 1*:

line -12: *Solution 2*: \searrow *Solution 2*:

►Page 513 line -8 _____ 22 Mar 1999

$R(j) \leq a_{ij} \searrow R(i) \leq a_{ij}$

Page 514 end of the program _____ 18 May 1999

program lines -3 and -2: interchange 'ENT1 0' with 'J3P 3B'

line 6 after the program: $540u \searrow 530u$

Page 514 line -6 _____ 25 Aug 2002

we can solve \searrow and $m \geq n$, we can solve

►Page 515 line 14 _____ 31 Jul 1997

in 924744796234036231 our case, \searrow in our case,

Page 515 new answer _____ 24 Apr 1997

12. M. Hofri and P. Jacquet [*Algorithmica* **22** (1998), 516-528] have analyzed the case when the $m \times n$ matrix entries are distinct and in random order. The running times of the two MIX programs are then respectively $(6mn + 5mH_n + 8m + 6 + 5(m+1)/(n-1))u + O((m+n)^2/\binom{m+n}{m})$ and $(5mn + 2nH_m + 7m + 7n + 9H_n)u + O(1/n) + O((\log n)^2/m)$, as $m \rightarrow \infty$ and $n \rightarrow \infty$, assuming that $(\log n)/m \rightarrow 0$.

Page 516 answer 14 _____ 31 Aug 1998

[Udo Wermuth has indeed squeezed out an additional $9u$, $10u$, or $12u$, depending on the year. But details are omitted here because MIX will soon be replaced.]

Page 518 lines 18 and 19 _____ 10 Jan 2003

Calendrical Calculations ... 1997) \searrow *Calendrical Calculations* by E. M. Reingold and N. Dershowitz (Cambridge Univ. Press, 2001)

- **Page 519** line 2 of answer 17 _____ 20 May 1999
 $H_n \rightsquigarrow H_N$ and $H_{2m} \rightsquigarrow 2H_{2m}$
- **Page 520** line -3 of answer 19 _____ 25 Nov 1997
Société \rightsquigarrow *Société*
- **Page 521** replacement for line 11 _____ 20 Jan 2008
 LDA * 4 Waste $2u$ of time.
- **Page 522** lines 1 and 2 of answer 6 _____ 24 Oct 2005
 The tital time is increased ... name. \rightsquigarrow The total time decreases by $8u$ for every blank word following a “(”, because lines 30–32 cost $4u$ while lines 26–28, 33–34, 36–38 cost $12u$. It decreases by $2u$ for every blank word following a name, because lines 68–71 cost $5u$ while 42–46 or 75–79 cost $7u$.
- Page 524 line 1 of answer 17 _____ 26 Feb 2008
 The probability \rightsquigarrow The probability p_m
- Page 524 line 2 of answer 17 _____ 24 Dec 2001
 $(m/mH_n) \rightsquigarrow m/(mH_n)$
- **Page 524** answer 19 _____ 15 Sep 1999
 $1/ \rightsquigarrow n!/$ (in three places)
- **Page 524** lines 1, 3, and 9 of answer 22 _____ 01 Jan 2006
 $j \geq 0 \rightsquigarrow j > 0$ (four times)
- Page 525** lines 4 and 5 of answer 23 _____ 04 Nov 2003
 [to appear] \rightsquigarrow [Ph.D. thesis, École Polytechnique (Paris, 1996)]
- **Page 525** answer 24 _____ 24 Dec 2001
Proc. IFIP Congress (1971) \rightsquigarrow *Proc. IFIP Congress (1971)*
- Page 526** new answer _____ 14 Oct 2008
 28. See I. N. Herstein and I. Kaplansky, *Matters Mathematical* (1974), §3.5.
- Page 526** new sentence for answer 29 _____ 07 Apr 2003
 Incidentally, the original number of the k th man executed is $2n + 1 - (2n + 1 - 2k)2^{\lfloor \lg(2n/(2n+1-2k)) \rfloor}$. [Armin Shams, *Proc. Nat. Computer Conf. 2002*, English papers section, 2 (Mashhad, Iran: Ferdowsi University, 2002), 29–33.]
- Page 526** new answer 31 _____ 14 Oct 2008
 See *CMath*, Section 3.3. Let $x_0 = jm$ and $x_{i+1} = (m(x_i - n) - d_i)/(m - 1)$, where $1 \leq d_i < m$. Then $x_k = j$ if and only if $a_k j = b_k n + t_k$, where $a_k = m^{k+1} - (m-1)^k$, $b_k = m(m^k - (m-1)^k)$, and $t_k = \sum_{i=0}^{k-1} m^{k-1-i}(m-1)^i d_i$. Since $a_k \perp b_k$ and the $(m-1)^k$ possibilities for t_k are distinct, the average number of k -step fixed elements is $(m-1)^k/a_k$.
- Page 527** last line of answer 34 _____ 26 Oct 1997
 efficient. \rightsquigarrow efficient. [W. Fletcher and R. Silver, *CACM* 9 (1966), 326.]
- **Page 527** lines -3 and -2 of answer 35 _____ 01 Jan 2006
 interchange γ'' with $\beta\gamma'$ \rightsquigarrow interchange $\gamma''\beta$ with γ'

► **Page 527** new answer _____ 29 Jan 2010

37. The result is clear when $n \leq 2$. Otherwise we can find $a, b < n$ such that π takes a to b . Then $(na)\pi(bn) = (\alpha a)(b\beta)$ for $(n-1)$ -cycles (αa) and $(b\beta)$ if and only if $\pi = (n\alpha a)(b\beta n)$. [See A. Jacques, C. Lenormand, A. Lentin, and J.-F. Perrot, *Comptes Rendus Acad. Sci.* **266** (Paris, 1968), A446–A448.]

► **Page 529** lines 2–7 of answer 3 _____ 17 Jul 2006

But if ... program.) \rightsquigarrow But the comparison indicator isn't initialized; and if the final period is preceded by a replication digit, it won't be noticed. [Note: The most nitpickingly efficient program would probably remove lines 40, 44, and 48, and would insert "CMPA PERIOD" between lines 26 and 27, "CMPX PERIOD" between lines 59 and 60. The state of the comparison indicator should then become part of the coroutine characteristics in the program documentation.]

► **Page 529** line -4 _____ 28 Dec 1997

is not to invent \rightsquigarrow is not hard to invent

► **Page 531** in the right-hand column of answer 2 _____ 03 Sep 2008

.....lines 14–31		LD1 I1REG
LD1 I1REG	\rightsquigarrowlines 14–35
.....lines 32–35		

► **Page 532** lines 3–8 of answer 3, right column _____ 04 Sep 2008

BUFMAX CON ENDBUF1		* BUFFER AREAS
* BUFFER AREAS		OUTBUF1 ORIG **100
OUTBUF1 ORIG **100	\rightsquigarrow	ENDBUF1 CON **101 (ENDBUF2)
ENDBUF1 CON ENDBUF2		OUTBUF2 ORIG **100
OUTBUF2 ORIG **100		ENDBUF2 CON ENDBUF1
ENDBUF2 CON ENDBUF1 █		BUFMAX CON ENDBUF1 █

► **Page 532** line 1 of answer 7 _____ 16 Jul 1998

One way \rightsquigarrow One way is

► **Page 534** new instruction to follow 'J1P 1B' in answer 15 _____ 05 Sep 2008

JBUS *(U)

► **Page 534** line 7 of answer 18 _____ 11 Mar 1998

goint \rightsquigarrow going

Page 534 in answer 18 _____ 06 Jan 2011

[The revised algorithms are now called A' , R' , and B' , to distinguish them from Algorithms A, R, and B of the text.]

► **Page 534** line 10 of answer 19 _____ 24 Dec 2001

1339–1342 \rightsquigarrow 1338–1342

► **Page 536** line 1 of answer 4 _____ 01 Jan 2009

D. André (1878): \rightsquigarrow J. Aebly and D. Mirimanoff [*L'Enseignement Math.* **23** (1923), 185–189]:

Page 536 line -13 _____ 17 Mar 1998

Abraham De Moivre \rightsquigarrow Abraham de Moivre

►Page 537 line -4 _____ 01 Jan 2006

$$A_m(n, -2) \rightsquigarrow A_m(n+1, -2)$$

►Page 539 answer 12 _____ 17 Nov 1999

line 3: We have \rightsquigarrow We have, for sufficiently small α ,
replacement for line 7:

$$(-1)^n \binom{k+1/2}{n} = \binom{n-k-3/2}{n} = \frac{\Gamma(n-k-1/2)}{\Gamma(n+1)\Gamma(-k-1/2)} = \frac{(-1/2)^{k+1}}{\sqrt{\pi n}} n^{-k-1/2},$$

replacement for line 11:

$$c_j = \sqrt{\frac{1-\alpha}{\pi}} \sum_{k=0}^j \binom{1/2}{k} (-1/2)^{k+1} \left\{ \begin{matrix} j+1/2 \\ k+1/2 \end{matrix} \right\} \frac{\alpha^k}{(1-\alpha)^k}.$$

►Page 540 line 4 of answer 3 _____ 18 Jun 1998

$$\text{LDA } 2, 7:1 \rightsquigarrow \text{LDA } X2, 7:1$$

►Page 540 answer 4 _____ 13 Aug 1997

(i) (ii) (iii) (iv) (v) \rightsquigarrow (a) (b) (c) (d) (e)

Page 541 line 11 _____ 05 Oct 1999

$$\mathbb{N} \leftarrow 0 \rightsquigarrow \mathbb{N} \leftarrow 0$$

►Page 542 line 2 of answer 10 _____ 10 Sep 2008

$$\sum_{1 \leq j \leq k \leq n} \rightsquigarrow \sum_{1 \leq j < k \leq n}$$

►Page 542 new answer for exercise 11 _____ 04 Oct 2005

11. Counting as before, we find that the expected number is

$$E_{mnt} = \frac{1}{n^m} \binom{n}{2} \sum_{k=1}^m \sum_{r \geq t} (k-1) \binom{k-2}{r} (n-1)^{k-2-r} n^{m-k};$$

here r is the number of entries in a_1, a_2, \dots, a_{k-1} that equal a_k . This quantity can also be expressed in the simpler form

$$E_{mnt} = \frac{1}{n^m} \binom{n}{2} \sum_{k > t} \binom{m}{k} (n-1)^{m-k} \left(\binom{k}{2} - \binom{t+1}{2} \right), \quad \text{for } t \geq 0.$$

Is there a simpler way yet to give the answer? Apparently not, since the generating function for given n and t is

$$\sum_m E_{mnt} z^m = \frac{n-1}{2n} \frac{z}{(1-z)^3} \left(\frac{z}{n-(n-1)z} \right)^{t+1} (z + (1-z)n(t+1)).$$

►Page 543 line 1 of answer 14 _____ 18 Nov 1999

$$n/m + \sqrt{n} x_j \rightsquigarrow m/n + \sqrt{m} x_j$$

►Page 543 line 5 of answer 14 _____ 11 Sep 2008

$$(\sqrt{m})^{1-m} \rightsquigarrow (\sqrt{m})^{1-n}$$

►Page 543 line -2 _____ 18 Nov 1999

$$I_1 = I_2/n \rightsquigarrow I_1 = I_2/n^2$$

- **Page 544** line 6 _____ 18 Nov 1999
 $3 \leq j \leq m \rightsquigarrow 3 \leq j \leq n$
- **Page 544** line 3 of answer 17 _____ 20 Feb 2009
less \rightsquigarrow more
- **Page 547** line 3 of answer 13 _____ 13 May 2001
 $2^{2^n(n+O(1))}$ by a factor of $2^{2^n+O(n)} \rightsquigarrow 2^{2^n(n+O(\log n))}$ by a factor of $e^{2^n+O(n)}$
- Page 549* line 12 _____ 28 Sep 2002
alpha. \rightsquigarrow alphameric.
- **Page 550** answer 27 _____ 26 Feb 2008
line 10: J2Z B1 \rightsquigarrow J2Z 2F
line 18: LD1 N \rightsquigarrow 2H LD1 N
- **Page 550** line -14 _____ 26 Jul 1997
those move \rightsquigarrow those moves
- Page 555* line 04 of answer 9 _____ 20 Aug 1997
D2. Should door open? \rightsquigarrow D2. Should doors open?
- Page 555* line 14 of answer 9 _____ 04 Jul 2001
are zero \rightsquigarrow are zero.
- Page 555* last line of answer 9 _____ 03 Aug 1998
subroutine. \rightsquigarrow subroutine. **I**
- **Page 556** line 1 of answer 5 _____ 28 Sep 2002
A0 \rightsquigarrow A0
- **Page 557** replacement for lines 2 and 3 _____ 25 Oct 1997
in $X[I_{a_1} + B_1, I_{a_2} + B_2, \dots, I_{a_m} + B_m]$, where $B_1 B_2 \dots B_m$ is an inversion table for $a_1 a_2 \dots a_m$ as defined in exercise 5.1.1-7.)
- **Page 557** line 1 of answer 15 _____ 24 Dec 2001
rI1 \equiv PIV0T, J, rI2 \equiv P0, rI3 \equiv Q0, rI4 \equiv P, rI5 \equiv P1, X; \rightsquigarrow rI1 \equiv PIV0T, J; rI2 \equiv P0;
rI3 \equiv Q0; rI4 \equiv P; rI5 \equiv P1, X;
- **Page 560** line 4 of answer 21 _____ 01 Jan 2006
 $L_{r-1} \rightsquigarrow L_{r-1} + (M_r - 1)^r$
- Page 561* line 1 of answer 6 _____ 30 Aug 1997
[There's slightly too much space before the word "Let".]
- **Page 561** line -2 _____ 19 Dec 2001
 $Z_1 \cap Z_2 \neq 0 \rightsquigarrow Z_1 \cap Z_2 \neq \emptyset$
- Page 562** line 1 _____ 29 Sep 2010
descendant of X , if \rightsquigarrow descendant of X (that is, a *proper* ancestor or descendant), if
- **Page 562** new answer 13 _____ 15 Oct 2007
13. $a_1, a_2, \dots, a_k, a_1, a_2, \dots, a_{k-1}, \dots, a_1, a_2, a_1.$

- Page 563 answer 17 _____ 13 Feb 2001
 $F \rightsquigarrow Z$ (three times)
- Page 564 line 6 _____ 03 Feb 1998
 2-D trees \rightsquigarrow 2-d trees
- Page 564 line 8 of answer 5 _____ 03 Sep 1997
 level order; \rightsquigarrow level order;
- Page 565 line 9 of answer 11 _____ 13 Nov 1999
 $+ \frac{11}{24} \sqrt{\frac{\pi}{n}} \rightsquigarrow - \frac{13}{24} \sqrt{\frac{\pi}{n}} + \frac{1}{2n}$
- Page 565 line 1 of answer 12 _____ 14 Jan 2000
 between step T2 and T3 \rightsquigarrow between steps T2 and T3
 between step T4 and T2 \rightsquigarrow in step T5
- Page 565 line 6 of answer 13 _____ 20 Apr 2001
 “Visit” \rightsquigarrow Visit
- Page 565 in answer 13 _____ 06 Jan 2011
 [The revised algorithm is now called T', to distinguish it from Algorithm T of the text.]
- Page 568 line 6 _____ 01 Jan 2006
 will triple traverse the tree \rightsquigarrow will traverse the tree in triple order
- Page 568 line 12 _____ 21 Mar 2004
 7.2.1 \rightsquigarrow 7.2.1.6
- Page 568 line -4 of answer 22 _____ 30 Jul 1998
 [insert ‘**I**’ in the rightmost column, to end the program.]
- Page 568 line -3 _____ 15 Sep 1999
 $\text{RLINK}(P) \leftarrow Q \rightsquigarrow \text{RLINKT}(Q) \leftarrow \text{RLINKT}(P), \text{RLINK}(P) \leftarrow Q$
- Page 569 line 5 of answer 26 _____ 26 Dec 2002
 $1 \leq j \leq n \rightsquigarrow 1 \leq j \leq 2n.$
- Page 570 line 5 of answer 31 _____ 30 Jul 1998
 $\Lambda. \rightsquigarrow \Lambda. \quad \mathbf{I}$
- Page 571 lines 2 and 4 of answer 37 _____ 28 Sep 2002
 $\text{NODE}(Q+1) \rightsquigarrow \text{NODE}(Q+1)$
- Page 572 line 2 of answer 12 _____ 15 Oct 2007
 $R \leftarrow \text{TREE}(\text{“}\uparrow\text{”}, \text{TREE} \rightsquigarrow R \leftarrow \text{TREE}(\text{“}\uparrow\text{”}, R, \text{TREE}$
- Page 575 line 6 of answer 2 _____ 29 Aug 1997
 $P \leftarrow P - 1 \rightsquigarrow P \leftarrow P - c$
- Page 577 line 5 of answer 17 _____ 29 Aug 1997
 $P \leftarrow P - 1 \rightsquigarrow P \leftarrow P - c$

- **Page 577** line 9 of answer 18 _____ 24 Dec 2001
 $\text{INFO2}[k] \leftarrow \text{INFO}[i] \rightsquigarrow \text{INFO2}[k] \leftarrow \text{INFO1}[i]$
- Page 579** line -2 of answer 8 _____ 21 Aug 2001
 D. E. Knuth \rightsquigarrow A. Nahapetian, *Acta Informatica* **3** (1973), 37–41; D. E. Knuth
- Page 579* line 4 of answer 11 _____ 14 Feb 1999
 “common”, \rightsquigarrow “common,”
- Page 580* lines 3 and 6 of answer 13 _____ 23 Oct 2006
 a tree \rightsquigarrow a free tree
- **Page 581** replacement for answer 15 _____ 01 Feb 2004
 15. True for finite graphs: If it is connected and balanced and has more than one vertex, it has an Eulerian trail that touches all the vertices. (But false in general.)
- **Page 581** lines 3 and 4 of answer 16 _____ 21 Aug 2004
 tracing out ... (this graph is balanced). \rightsquigarrow tracing out an Eulerian trail in G , because the game ends when the fourth arc to V_{13} is encountered (namely, when the fourth king turns up).
- Page 581* line 3 of answer 17 _____ 20 Apr 1998
 methods of Section \rightsquigarrow methods of Section
- Page 581* bottom line and beginning of page 582 _____ 05 Sep 2006
 2.3.4.4; it also ... direct proof: \rightsquigarrow 2.3.4.4. But such a simple answer deserves a simple, direct proof, and indeed there is one [see Tor B. Staver, *Norsk Matematisk Tidsskrift* **28** (1946), 88–89].
- **Page 582** line 13 _____ 10 Dec 1998
 See Section \rightsquigarrow See
- **Page 583** line 3 _____ 28 Dec 1997
 oriented trees is \rightsquigarrow oriented trees
- Page 583** better answer for exercise 21 _____ 06 Sep 2001
 Replace the last several lines on page 583, beginning with ‘Add the indeterminate $\lambda \dots$ ’, by the following new copy:
 Add the indeterminate λ to every diagonal element of A and A^* . If $t(G)$ and $t(G^*)$ are the numbers of oriented subtrees of G and G^* , we then have $\det A = \lambda t(G) + O(\lambda^2)$, $\det A^* = \lambda t(G^*) + O(\lambda^2)$. (The number of oriented subtrees of a balanced graph is the same for any given root, by exercise 22, but we do not need that fact.)
 If we group vertices V_{jk} for equal k the matrix A^* can be partitioned as shown above. Let $B_{kk'}$ be the submatrix of A^* consisting of the rows for V_{jk} and the columns for $V_{j'k'}$, for all j and j' such that V_{jk} and $V_{j'k'}$ are in G^* . By adding the 2nd, ..., m th columns of each submatrix to the first column and then subtracting the first row of each submatrix from the 2nd, ..., m th rows, the matrix A^* is transformed so that

$$B_{kk'} = \begin{pmatrix} a_{kk'} & * & \dots & * \\ 0 & 0 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & 0 \end{pmatrix} \quad \text{for } k \neq k', \quad B_{kk} = \begin{pmatrix} \lambda + a_{kk} & * & \dots & * \\ 0 & \lambda + m & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & \lambda + m \end{pmatrix}.$$

The asterisks in the top rows of the transformed submatrices turn out to be irrelevant, because the determinant of A^* is now seen to be $(\lambda + m)^{(m-1)n}$ times


$$\det \begin{pmatrix} \lambda + a_{00} & a_{01} & \dots & a_{0(n-1)} \\ a_{10} & \lambda + a_{11} & \dots & a_{1(n-1)} \\ \vdots & \vdots & \ddots & \vdots \\ a_{(n-1)0} & a_{(n-1)1} & \dots & \lambda + a_{(n-1)(n-1)} \end{pmatrix} = \lambda t(G) + O(\lambda^2).$$

Page 584 replacement for the top four lines _____ 06 Sep 2001

Notice that when $n = 1$ and there are m arcs from V_0 to itself, we find in particular that exactly m^{m-1} oriented trees are possible on m nodes labeled. This result will be obtained by quite different methods in Section 2.3.4.4.

► **Page 584** line 8 _____ 30 Jul 1997
314.) $\wedge \rightarrow$ 314.

Page 584 replacement for figure at bottom left _____ 31 Dec 1997



► **Page 585** last line of answer 27 _____ 14 Feb 1999
is combination $\wedge \rightarrow$ is a combination

► **Page 585** in answer 28 _____ 01 Jan 2006
line 9: is the determinant $\wedge \rightarrow$ is (*) times the determinant
line 17: column j_0 of the left section $\wedge \rightarrow$ row j_0 of the bottom section

Page 586 last lines of answer 4 _____ 08 Oct 2005

(But the stated assumption ... set of types.) $\wedge \rightarrow$
[But the stated assumption ... set of types. On the other hand, if such a tiling does exist, there is always a tiling that is *quasitoroidal*, in the sense that each of its $n \times n$ blocks occurs at least once in every $f(n) \times f(n)$ block, for some function f . See B. Durand, *Theoretical Computer Science* **221** (1999), 61–75.]

► **Page 587** line 7 _____ 22 Jul 2005
 $\gamma ST \wedge \rightarrow \gamma SJ$

Page 589 line 3 of answer 5 _____ 30 Jun 2008
 $\frac{1}{2} \wedge \rightarrow \frac{1}{2}$

Page 589 last line of answer 6 _____ 08 Oct 2010
Annals of Math. $\wedge \rightarrow$ *Annals of Math.* (2)

► **Page 590** line 1 of answer 16 _____ 08 Nov 2005
 $1 \leq j \leq n \wedge \rightarrow 1 \leq j < n$

- **Page 592** line 4 of answer 29 _____ 07 Jan 1999

$$\sum_{j,k} \frac{z^k}{k!} \sum_j \rightsquigarrow \sum_k \frac{z^k}{k!} \sum_j$$
- **Page 592** line 5 of answer 29 _____ 08 Nov 2005

$$\sum_k \frac{z^k}{k!} \binom{k-1}{j} \rightsquigarrow \sum_k \frac{z^k}{k!} \sum_j \binom{k-1}{j}$$
- Page 593** line 3 _____ 03 Dec 2004
 344–350. \rightsquigarrow 344–350; F. Acerbi, *Archive for History of Exact Sciences* **57** (2003), 465–502.
- **Page 593** line –6 _____ 08 Nov 2005

$$\sum_{j=1}^k (1 - d_j) \rightsquigarrow \sum_{j=1}^{k-1} (1 - d_j)$$
- **Page 594** denominator of display on line 12 _____ 10 Sep 1997

$$n_1! n_2! \rightsquigarrow n_1! n_2! \dots$$
- **Page 596** line 8 of answer 15 _____ 21 Sep 1997
 at each step. \rightsquigarrow at each step as in part (b).
- **Page 596** line 2 of answer 16 _____ 13 Jan 1999

$$(l'_j - l'_{j+1}) \rightsquigarrow (l'_k - l'_{k+1}) \quad (\text{twice})$$
- Page 596** line –2 _____ 31 Oct 1997
 5.2.3–20 and 21. \rightsquigarrow 5.2.3–20 and 5.2.3–21.
- Page 597** append a sentence to answer 1 _____ 01 Mar 2004
 [See H. G. Forder, *Mathematical Gazette* **45** (1961), 199–201.]
- Page 597** lines 1 and 2 of answer 3 _____ 28 Nov 1997
 from j to k if they are consecutive elements of the same part and $j < k$; \rightsquigarrow from i to j if i and j are consecutive elements of the same part and $i < j$;
- **Page 598** line 12 _____ 14 Jan 1999

$$(n-1, n+1, 1) \rightsquigarrow (n-1, n+1, k)$$
- Page 598** line 15 _____ 12 Mar 2004
 $(k-1)!$. \rightsquigarrow $(k-1)!$, a so-called *Narayana number* [T. V. Narayana, *Comptes Rendus Acad. Sci.* **240** (Paris, 1955), 1188–1189].
- **Page 598** line 18 _____ 17 Sep 1997
 different from the discussed \rightsquigarrow different from the one discussed
- **Page 598** line 19 _____ 01 Dec 1997
 (1970) \rightsquigarrow (1971)
- Page 598** line 21 _____ 07 Apr 2006
 171–179. \rightsquigarrow 171–179; N. Dershowitz and S. Zaks, *Discrete Math.* **64** (1986), 215–218.
- **Page 598** line –2 _____ 14 Feb 1999
 these the new \rightsquigarrow these new

► **Page 600** line 3 following (***) _____ 08 Nov 2005

$(t, w)(v, u) \rightsquigarrow (t, w)(u, v)$

Page 603 changes for consistency _____ 29 Oct 1997

line -20: Solution 1: \rightsquigarrow *Solution 1:*

line -9: Solution 2: \rightsquigarrow *Solution 2:*

Page 606 in answer 12 _____ 06 Jan 2011

[The revised algorithm is now called B', to distinguish it from Algorithm B of the text.]

Page 606 in answer 14, continuing to page 607 _____ 06 Jan 2011

[The revised algorithm is now called C', to distinguish it from Algorithm C of the text.]

► **Page 607** line 3 of answer 6 _____ 19 Oct 1998

ROVER \leftarrow LINK(Q) \rightsquigarrow ROVER \leftarrow LINK(Q) if LINK(Q) \neq Λ , otherwise set ROVER \leftarrow LOC(AVAIL).

Page 608 in answer 8 _____ 06 Jan 2011

[The revised algorithm is now called A'', to distinguish it from Algorithms A and A' of the text.]

► **Page 608** replacement for lines 2–4 of answer 10 _____ 15 Oct 2003

At the beginning of step B3, insert “If $P0 + N > P$ and $P \neq \Lambda$, set $N \leftarrow \max(N, P + \text{SIZE}(P) - P0)$, $P \leftarrow \text{LINK}(P)$, and repeat step B3.”

Step B4, for “ $Q + \text{SIZE}(Q) = P0$ ”, read “ $Q + \text{SIZE}(Q) \geq P0$ ”; and for “ $\text{SIZE}(Q) \leftarrow \text{SIZE}(Q) + N$ ” read “ $\text{SIZE}(Q) \leftarrow \max(\text{SIZE}(Q), P0 + N - Q)$ ”.

Page 608 in answer 12 _____ 06 Jan 2011

[The revised algorithm is now called A*, to distinguish it from Algorithm A and A' of the text and Algorithm A'' of answer 10. This change extends to page 609, and also affects answer 19 on pages 610–611.]

Page 610 in answers 15, 16 _____ 06 Jan 2011

D1 \rightsquigarrow C1, D2 \rightsquigarrow C2, ..., D8 \rightsquigarrow C8

Page 610 line -3 _____ 02 Feb 1999

LINK(P2+1) \rightsquigarrow LINK(P2 + 1)

Page 612 program lines 17, 18, 20 in answer 27 _____ 06 Nov 2006

LINKF \rightsquigarrow LINKB, LINKB \rightsquigarrow LINKF, AVAILF \rightsquigarrow AVAILB (six changes altogether)

► **Page 612** program line 04 in answer 28 _____ 20 Aug 1997

buddy_kL \rightsquigarrow buddy_k(L)

► **Page 612** lines 12 and 13 of answer 28 _____ 04 Sep 2003

11	S2	LD2	0,5(LINKB)	S	<u>S2. Combine with buddy.</u>
12		LD3	0,5(LINKF)	S	

Page 613 new answer _____ 10 Nov 2002

MALLAT

32. Steven Crain points out that the method always frees all blocks and starts afresh before 16667 units of time have elapsed; hence the stated limit certainly exists. *Proof:* Let $u_n = n + t_n$, so that $g_n = \lfloor \frac{5}{4} \min(10000, f(u_{n-1} - n), f(u_{n-2} - n), \dots, f(u_0 - n)) \rfloor$. Let $x_0 = 0$ and $x_1 = u_0$, and $x_{k+1} = \max(u_0, \dots, u_{x_k-1})$ for $k \geq 1$. If $x_k > x_{k-1}$ then

$$u_n \leq n + \frac{5}{4}f(x_k - n) = \frac{5}{4}x_k - \frac{1}{4}n \leq \frac{5}{4}x_k - \frac{1}{4}x_{k-1} \quad \text{for } x_{k-1} \leq n < x_k;$$

therefore $x_{k+1} - x_k \leq \frac{1}{4}(x_k - x_{k-1})$, and we must have $x_k = x_{k-1}$ before reaching time $12500 + \lfloor 12500/4 \rfloor + \lfloor 12500/4^2 \rfloor + \dots$.

Page 620 Table 2 06 May 1999

In the next edition I plan to give these constants to 36 hexadecimal places, instead of 45 octal places.

► **Page 624** entry for Kronecker delta _____ 14 Aug 1998

1.2.6 $\searrow \rightarrow$ 1.2.3

► **Page 625** new entry (just above $[a..b]$) _____ 20 Mar 2006

$$a_1 + a_2 + \dots + a_n \mid n\text{-fold sum: } \sum_{j=1}^n a_j \quad \mid 1.2.3$$

► **Page 626** entry for $\exp x$ _____ 09 Aug 1999

1.2.2 $\searrow \rightarrow$ 1.2.9

Page 628 a new appendix _____ 06 Jan 2011

Beginning with the 27th printing, Volume 1 now has an “Appendix C” analogous to Appendix C of Volume 4A, which lists the page numbers on which named algorithms, theorems, programs, definitions, ..., can be found. A copy of this two-page appendix can be found online in the author’s website. The main index therefore begins now on page 630.

Page 629 quotation for bottom of the page _____ 07 Jan 2011

Numerical experimentations are necessary to fully understand the algorithms and theorems in this book.

— STÉPHANE MALLAT, *A Wavelet Tour of Signal Processing* (1998)

Page 628 and following 24 Apr 1997

Miscellaneous changes to the existing index of Volume 1 are collected here, including corrections and amendments to the old entries as well as new entries that are occasioned by the new material. Thus, the lines of the full index that have changed serve also as an index to the present document. However, when a correction or amendment has caused an old index entry to be deleted, the deletion is usually not indicated.

- 2-d trees, 564.
 Aarons, Roger Michael, 528.
 Abel, binomial theorem, 58, 71–73, 398.
 Abel, limit theorem, 95, 588.
 Acerbi, Fabio, 593.
 Aebly, Jakob, 536.
 ALF (alphabetic data), 151, 152, 155.
 ALGOL language, viii, 202, 229.
 Alhazen, *see* Ibn al-Haytham.
 Ancestor, in a tree structure, 311, 348, 562.
 Arabic mathematics, 1, 162.
 Assertions, inductive, 15–20.
 Barrington (= Mix Barrington), David Arno, 526.
 Berners-Lee, Mary Lee née Woods, 230.
 Bernoulli, Daniel, 83.
 Bhāskara II, Ācārya, son of Maheśvara (भास्कराचार्य, महेश्वरपुत्र), 54.
 Binary operator: A function of two variables, 337.
 Binary trees, 312, 317, 318–337, 363, 459.
 Binet, Jacques Philippe Marie, 36, 39, 407, 475, 582.
 Binomial coefficients, inequalities involving, 74.
 Binomial number system, *see* Combinatorial number system.
 Bolzano, Bernard Placidus Johann Nepomuk, 382.
 Boolean functions, 526–527.
 Bracket notation for logical statements, 32–33, 471, *see* Iverson's convention.
 C language, 233, 556.
 Capelli, Alfredo, 50.
 Carlitz, Leonard, 96, 499, 506.
 Case Institute of Technology, i.
 Cassini, Gian (= Giovanni = Jean) Domenico (= Dominique), 81.
 Cauchy, Augustin Louis, 39, 92, 475, 490, 506, 520, 582.
 Chakravarti, Gurugovinda (গুরুগোবিন্দ চক্রবর্তী), 53.
 Charles Philip Arthur George of Edinburgh, Prince of Wales, 310.
 Chia Hsien (賈宪), 53.
 Chinese mathematics, 53, 59, 75, 407.
 Chowla, Paromita (পারমিতা চাওলা or पारोमिता चावला), 307.
 Christie Mallowan, Agatha Mary Clarissa Miller, xvii.
 COBOL language, 424–434, 457, 458.
 Combinations of n objects taken k at a time, 52–53, 69, 94.
 with repetitions permitted, 73, 92, 95, 386, 388.
 Composition of permutations, *see* Multiplication of permutations.
 Connected directed graph, 372.
 Continued fractions, 498, 565.
 Convex function, 404, 596.
 Conway, Melvin Edward, 151, 229.
 Conway, John Horton, 19, 80, 273, 408, 600.
 Cook, Theodore Andrea, 81.
 Coxeter, Harold Scott MacDonald, 80, 162, 407, 408.
 Crain, Steven Paul, 613.
 Cullen, Christopher, 75.
 Cycle, Hamiltonian, 374, 379.
 Cycle lemma, 593.
 Cyclic shift, 185.
 Dahl, Ole-Johan, 229, 230, 461, 462.
 de Moivre, Abraham, 74, 83, 87, 106, 182, 474, 536.
 Dershowitz, Nachum (נחום דרשוביץ), 518, 588, 598.
 Descendant, in a tree structure, 311, 348, 562.
 Dewey, Melville (= Melvil) Louis Kossuth, notation for binary trees (due to Galton), 317, 331, 346, 405.
 Dickson, Leonard Eugene, 81, 484.
 Dijkstra, Edsger Wybe, 17, 191, 230, 231, 240, 459, 462, 545, 580, 605.
 d'Imperio, Mary Evelyn, 462.
 Doyle, Arthur Ignatius Conan, 465.
 Durand, Bruno, 586.
 Dvoretzky, Aryeh (אריה דבורצקי), 593.
 e , 23, 619–620, 626.
 Ellen [Fich] Whyne, Faith, 523.
 Ellipses (\cdots or \dots), 27, 34, 46.
 Erdélyi, Artúr (= Arthur), 399.
 Euler, Leonhard (Ейлеръ, Леонардъ = Эйлер, Леонард), 49, 50, 52, 57, 75, 76, 87, 111, 374, 407, 472, 496, 536, 600.
 constant γ , 75, 107, 114, 619–620.

- Eulerian trails in a directed graph, 374–376, 379, 584.
enumeration of, 380.
Even permutations, 185.
Fibonacci, Leonardo, of Pisa (= Leonardo filio Bonacii Pisano), 79–80, 84.
Fibonomial coefficients, 85, 484, 499.
Fich, Faith Ellen Whyne, 523.
Figurate numbers, 54.
Flajolet, Philippe Patrick Michel, 501, 506, 543, 565.
Fletcher, William, 527.
Floyd, Robert W, x, 17, 20, 422, 467, 509.
Forder, Henry George, 597.
Förstemann, Wilhelm August, 490.
FORTRAN language, viii, 231, 233, 296, 360, 458, 460.
Fraenkel, Aviezri S (אביעזרי פרנקל), 251.
Gauß (= Gauss), Johann Friderich Carl (= Carl Friedrich), 49, 58, 95, 490.
Generating functions, 82–84, 87–96, 243, 386–389, 391–392, 394–399, 539, 542.
Goldbach, Christian, 49, 407, 472.
Goldstine, Herman Heine, 18, 229, 231.
Greek mathematics, 2, 19, 54, 593.
Gustavson, Fred Gehrung, 305.
Halāyudha (हलायुध), 53.
Hald, Anders, 103.
Harmonic numbers, table, 621–622.
Haros, Charles, 520.
Heap, 435, *see* Pool of available nodes.
Hemacandra, Ācārya (आचार्य हेमचन्द्र), 80.
Herstein, Israel Nathan, 526.
Hindu mathematics, 53–54, 80, 495.
Hipparchus of Nicæa (Ἱππαρχος ὁ ἐκ Νικαίας), 592.
Hofri, Micha (מיכה חפרי), 515.
Horning, James Jay, 230.
Hwang, Frank Kwangming (黃光明), 405, 595.
Hwang, Hsien-Kuei (黃顯貴), 66.
Hyperfactorial, 116.
Ibn al-Haytham, Abū ‘Alī al-Ḥasan (= Alhazen, أبو علي الحسن بن الهيثم), 162.
ILLIAC I computer, 230.
in situ permutation, 9, 165, 185, 523.
Inclusive ancestors and descendants, 311.
Indian mathematics, 53–54, 80, 495.
Inductive assertions, 15–20.
Inorder for a binary tree, 319–323, 330–332, 346.
Irrational radix, 86.
Islamic mathematics, 1, 162.
Iverson, convention, 32–33, 61, 103, 471.
Jacobi, Carl Gustav Jacob, 490.
Jacques, Alain, 527.
Jacquet, Philippe Pierre, 515.
Japanese mathematics, 112, 115.
Java language, 233.
Jenkins, David Philip, 460.
Josephus, Flavius, son of Matthias (יוסף בן מתתיהו = Φλάβιος Ἰωσήπος Μαθθίου), problem, 162, 184.
Kahn, Arthur Bertram, 268.
Kaplansky, Irving, 184, 526.
Kepler, Johannes, 80, 81.
Kinkelin, Hermann, 504.
Kramp, Christian, 49, 486.
Kruskal, Joseph Bernard, Jr., 386, 588.
Left-child/right-sibling links, 335, 348.
Left-sibling/right-child links, 346.
Legendre (= Le Gendre), Adrien Marie, 48, 49, 51.
Lenormande, Claude, 527.
Lentin, André, 527.
Lexicographic order, 20, 299–300, 306, 564.
Lushbaugh, Warren Arthur, 19.
Mahāvīra, Ācārya (महावीराचार्य), 54.
Mallat, Stéphane Georges, 629.
Mark I computer (Manchester/Ferranti), 18.
Markov (= Markoff), Andrei Andreevich (Марков, Андрей Андреевич), the elder, 495.
McKeeman, William Marshall, 230.
Meek, Homer Vergil, 230.
Meton of Athens (Μέτων ὁ Ἀθηναῖος), cycle, 160.
Mirimanoff, Dmitri (Мириманов, Дмитрий Семёнович), 536.
Mix Barrington, David Arno, 526.
Moivre, Abraham de, 74, 83, 87, 106, 182, 474, 536.
Monte Carlo method: Experiments with random data, 254, 445–447.
Moore School of Electrical Engineering, 230.
Motzkin, Theodor (= Theodore) Samuel (תיאודור שמואל מוצקין), 85, 593.
Multilist representation, 301.
Multiple-precision constants, 525, 619–621.
Narayana, Tadepalli Venkata, 598.
Nearest common ancestors, 311.
Needham, Noel Joseph Terence Montgomery (李約瑟), 59.
Nicomachus of Gerasa (Νικόμαχος ὁ ἐκ Γεράσων), 19.
NOP (no operation), 136, 142.
Nygaard, Kristen, 229, 461.
Oriented walk in a directed graph, 372.
Overlapping subtrees, 326, 603.
Paper sizes, 318.
Paper tape, 136–137, 229, 231.
Parsing Machine, 201.
Patterns in permutations, 243.
Peck, John Edward Lancelot, 461.
Pedigree, 310–312.
Perfect shuffle, 184.
Perrot, Jean-François, 527.
Persian mathematics, 1.

- Pi (π), as “random” example, 397.
 Piṅgala, Ācārya (आचार्य पिङ्गल), 53.
 Prinz, Dietrich Günter, 230.
 Profile of a program: The number of times each instruction is performed, 145, 170, 214, 296, 528.
 Proper ancestors and descendants, 311, 562.
 q -nomial coefficients, 65, 73, 484, 491.
 Quasitoroidal tiling, 586.
 Rabbits, 79, 84.
 Rahman, Mizan (মিজান রহমান), 490.
 Recursion induction, 321, 565, 569.
 Reingold, Edward Martin (ריינוולד), 23, 518.
 Remainder, 40, 160.
 Redheffer, Raymond Moos, 491.
 Reversal of a string, 185.
 Ribenboim, Paulo, 466.
 Riemann, Georg Friedrich Bernhard, zeta function $\zeta(s)$, 76, 504.
 Right-child/left-sibling links, 346.
 Right-sibling/left-child links, 335, 348.
 Root of a number, 22, 25, 110.
 Root of a tree, 308, 309, 317.
 changing, in an oriented tree, 377.
 Rothe, Heinrich August, 63, 71, 486.
 Schlesvig-Holstein-Sønderborg-Glücksborg, Christian von, *see* Christian IX.
 Schweins, Franz Ferdinand, 489.
 Seki, Takakazu (關孝和), 112, 115.
 Shams Baragh, Armin (آرمین شمس براق), 526.
 Shared subtrees, 326, 603.
 Sibling link, 334, 336, 427–433; *see also* **RLINK** in trees.
 Sideways addition, 131, 480.
 Silver, Roland Lazarus, 527.
 Singh, Parmanand (परमानन्द सिंह), 80.
 Solitaire (patience), 377–378.
 Spanning subtrees, 365–370, 378–379.
 Staver, Tor B., 582.
 Stirling, James, approximation, 47, 51, 115–116.
 Stirling numbers, 66–69, 71–74, 78, 99–100, 506, 582, 591.
 Symmetric functions, 92–94, 472–473, 497.
 System/360 computers, 124, 189, 230.
 Szekeres, György (= George), 595.
 Tape, paper, 136–137, 229, 231.
 Tamari, Dov (דוב תמרי), born Bernhard Teitler, 577.
 Taylor, Brook, series, 83, 102.
 TeX, iv, xi, 193, 202, 611, 650.
 Three-dots notation (\dots or \dots), 27, 34, 46.
 Trails, Eulerian, in a directed graph, 374–376, 379–380, 584.
 Todhunter, Isaac, 182.
 Toscano, Letterio, 50.
 Tricks versus techniques, 574–575.
 Unary operator: A function of one variable, 337.
 Vandermonde, Alexandre Théophile, 59, 70, 71.
 von Neumann, John (= Margittai Neumann János), 18, 229, 231, 457.
 Walk, in a graph, 363.
 oriented, 372.
 random, 380–381.
 Warren, Don Wyman, 359.
 Wegner, Peter (= Weiden, Puttilo Leonovich = Вейден, Путтило Леонович), 306.
 Weierstrass (= Weierstraß), Karl Theodor Wilhelm, 382.
 Wilf, Herbert Saul, 65, 66, 93, 94, 484.
 Wirth, Niklaus Emil, 191, 461.
 Woods Berners-Lee, Mary Lee, 230.
 Wortman, David Barkley, 230.
 X1 computer, 231.
 XOR (bitwise exclusive-or), 442, 455, 553.
 Yang Hui (楊輝), 53.
 Zaks, Shmuel (שמואל זקס), 598.
 Zeckendorf, Edouard, 495.
 Zeta function $\zeta(s, x)$, 44, 76, 504.