

Asymmetric Cryptography: A Deep Dive

Eli Holderness — @eli@hachyderm.io — they/them/theirs

Asymmetric Cryptography: A Deep Dive

Eli Holderness — @eli@hachyderm.io — they/them/theirs

Eli (pronounced /'i:lai/) is a freelance developer advocate, recovering mathematician, and audience participator.

They like people, the web, and learning weird facts about computers.

They can be found on Mastodon at [@eli@hachyderm.io](mailto:eli@hachyderm.io), and — for now — on Twitter at [@eliholderness](https://twitter.com/eliholderness).



Agenda

Agenda

1. Brief history

Agenda

1. Brief history
2. How RSA works

Agenda

1. Brief history
2. How RSA works
3. How ECC works

Agenda

1. Brief history
2. How RSA works
3. How ECC works
4. QC & Shor's Algorithms

Agenda

1. Brief history
2. How RSA works
3. How ECC works
4. QC & Shor's Algorithms
5. What next?

1

A brief history of cryptography

KCDC is great!

A	B	C	D	E	F	G	H	I	J	K
G	H	I	J	K	L	M	N	O	P	Q

QIJI oy mxkgz!

KCDC is great!

+6	1	2	3	4	5	6	7	8	9	10	11
	7	8	9	10	11	12	13	14	15	16	17

QIJI oy mxkgz!

MESSAGE

13 5 19 19 1 7 5

MESSAGE

+

CIPHERT

13 5 19 19 1 7 5

+

3 9 16 8 5 18 20

MESSAGE

+

CIPHERT

13 5 19 19 1 7 5

+

3 9 16 8 5 18 20

=

16 14 35 27 6 25 25

MESSAGE

+

CIPHERT

13 5 19 19 1 7 5

+

3 9 16 8 5 18 20

=

16 14 9 2 6 25 25

MESSAGE

+

CIPHERTEXT

=

PNIBFY

13 5 19 19 1 7 5

+

3 9 16 8 5 18 20

=

16 14 9 2 6 25 25

symmetric cryptography
requires both parties to know
a specific secret

2

RSA & group theory

RSA cryptosystem

RSA cryptosystem

published 'officially' in 1977 by Rivest, Shamir
and Adleman

RSA cryptosystem

published 'officially' in 1977 by Rivest, Shamir
and Adleman

also developed independently in 1973 by
Clifford Cocks at GCHQ

RSA cryptosystem

published 'officially' in 1977 by Rivest, Shamir
and Adleman

also developed independently in 1973 by
Clifford Cocks at GCHQ

security based on the difficulty of factoring
large numbers $N = pq$ where p, q prime

worked example with $N = 323 = 17 * 19$

worked example with $N = 323 = 17 * 19$

$$a \equiv b \pmod{N}$$

when

$$a = b + kN \text{ for some integer } k$$

worked example with $N = 323 = 17 * 19$

We need to know $\lambda(N)$, the smallest number where
 $a^{\lambda(N)} \equiv 1 \pmod N$ for every a coprime to N

worked example with $N = 323 = 17 * 19$

$$\lambda(N) = 144$$

We need to know $\lambda(N)$, the smallest number where
 $a^{\lambda(N)} \equiv 1 \pmod{N}$ for every a coprime to N

$$\lambda(N) = \text{lcm}(\lambda(p), \lambda(q)) = \text{lcm}(p-1, q-1) = \text{lcm}(16, 18) = 144$$

worked example with $N = 323 = 17 * 19$

$$\lambda(N) = 144$$

We need to know $\lambda(N)$, the smallest number where
 $a^{\lambda(N)} \equiv 1 \pmod{N}$ for every a coprime to N

$$\lambda(N) = \text{lcm}(\lambda(p), \lambda(q)) = \text{lcm}(p-1, q-1) = \text{lcm}(16, 18) = 144$$

Choose e between 2 and N coprime to N ; let's pick 5

worked example with $N = 323 = 17 * 19$

$$\lambda(N) = 144 \quad e = 5; d = 29$$

We need to know $\lambda(N)$, the smallest number where
 $a^{\lambda(N)} \equiv 1 \pmod{N}$ for every a coprime to N

$$\lambda(N) = \text{lcm}(\lambda(p), \lambda(q)) = \text{lcm}(p-1, q-1) = \text{lcm}(16, 18) = 144$$

Choose e between 2 and N coprime to N ; let's pick 5

Find d such that $d * e \equiv 1 \pmod{\lambda(N)}$; this is 29

worked example with $N = 323 = 17 * 19$

$$\lambda(N) = 144 \quad e = 5; d = 29$$

Our public key is $(N, e) = (323, 5)$ and our private key is $d = 29$

worked example with $N = 323 = 17 * 19$

$$\lambda(N) = 144 \quad e = 5; d = 29$$

Our public key is $(N, e) = (323, 5)$ and our private key is $d = 29$

Someone wants to send us the message 14, 4, 3

worked example with $N = 323 = 17 * 19$

$$\lambda(N) = 144 \quad e = 5; d = 29$$

Our public key is $(N, e) = (323, 5)$ and our private key is $d = 29$

Someone wants to send us the message 14, 4, 3

To encrypt a number, they raise it to the power of $e = 5$:

$$14^5, 4^5, 3^5 = 537824, 1024, 243$$

worked example with $N = 323 = 17 * 19$

$$\lambda(N) = 144 \quad e = 5; d = 29 \quad m = (29, 55, 243)$$

Our public key is $(N, e) = (323, 5)$ and our private key is $d = 29$

Someone wants to send us the message 14, 4, 3

To encrypt a number, they raise it to the power of $e = 5$:

$$14^5, 4^5, 3^5 = 537824, 1024, 243$$

Then take the modulus of N :

$$14^5, 4^5, 3^5 \equiv 29, 55, 243 \pmod{N}$$

worked example with $N = 323 = 17 * 19$

$\lambda(N) = 144$ $e = 5; d = 29$ $m = (29, 55, 243)$

We received the message (29, 55, 243)

worked example with $N = 323 = 17 * 19$

$$\lambda(N) = 144 \quad e = 5; d = 29 \quad m = (29, 55, 243)$$

We received the message (29, 55, 243)

Decode by raising each number to the power of $d = 29$,
then taking the modulus of N

worked example with $N = 323 = 17 * 19$

$$\lambda(N) = 144 \quad e = 5; d = 29 \quad m = (29, 55, 243)$$

We received the message (29, 55, 243)

Decode by raising each number to the power of $d = 29$,
then taking the modulus of N

$$29^{29}, 55^{29}, 243^{29} \equiv 14, 4, 3 \pmod{N}$$

worked example with $N = 323 = 17 * 19$

$$\lambda(N) = 144 \quad e = 5; d = 29 \quad m = (29, 55, 243)$$

worked example with $N = 323 = 17 * 19$

$$\lambda(N) = 144 \quad e = 5; d = 29 \quad m = (29, 55, 243)$$

This works because $a^{\lambda(N)} \equiv 1 \pmod{N}$ for every a coprime to N

worked example with $N = 323 = 17 * 19$

$$\lambda(N) = 144 \quad e = 5; d = 29 \quad m = (29, 55, 243)$$

This works because $a^{\lambda(N)} \equiv 1 \pmod{N}$ for every a coprime to N

so $a^{\lambda(N)+1} \equiv a \pmod{N}$ for every a coprime to N

worked example with $N = 323 = 17 * 19$

$$\lambda(N) = 144 \quad e = 5; d = 29 \quad m = (29, 55, 243)$$

This works because $a^{\lambda(N)} \equiv 1 \pmod{N}$ for every a coprime to N

so $a^{\lambda(N)+1} \equiv a \pmod{N}$ for every a coprime to N

$$a^{\lambda(N)+1} = a^{145} = a^5 \times 29 = (a^5)^{29}$$

worked example with $N = 323 = 17 * 19$

$$\lambda(N) = 144 \quad e = 5; d = 29 \quad m = (29, 55, 243)$$

This works because $a^{\lambda(N)} \equiv 1 \pmod{N}$ for every a coprime to N

so $a^{\lambda(N)+1} \equiv a \pmod{N}$ for every a coprime to N

$$a^{\lambda(N)+1} = a^{145} = a^5 \times^{29} = (a^5)^{29}$$

So $(a^5)^{29} \equiv a \pmod{N}$ and we can recover the original message from the encrypted intermediate

limitations & considerations

limitations & considerations

requires large prime numbers, which are expensive to find

limitations & considerations

requires large prime numbers, which are expensive to find

if e is small enough that $M = m^e < N$, an attacker
can simply do $\sqrt[e]{M}$ to recover m


limitations & considerations


requires large prime numbers, which are expensive to find


if e is small enough that $M = m^e < N$, an attacker
can simply do $\sqrt[e]{M}$ to recover m

without padding, messages can be vulnerable to
chosen plaintext attacks

TURKEY TROTS TO WATER GG
FROM CINCPAC ACTION COM
THIRD FLEET INFO COMINCH
CTF SEVENTY-SEVEN X WHERE
IS RPT WHERE IS TASK FORCE
THIRTY FOUR **RR THE WORLD**
WONDERS








identity element

adding 0 doesn't change an element




identity element

adding 0 doesn't change an element

inverses

for every a in the group, there's
a b that makes $a + b = 0$ true



identity element


adding 0 doesn't change an element

inverses

for every a in the group, there's
a b that makes $a + b = 0$ true

associativity

$$1 + (4 + 2) = (1 + 4) + 2$$



identity element

adding 0 doesn't change an element

inverses


for every a in the group, there's a b that makes $a + b = 0$ true


associativity

$$1 + (4 + 2) = (1 + 4) + 2$$


closure

If a and b are in the group and $a + b = c$, then c is in the group






$4 \times 13 = 52$



$$\begin{aligned} & \textcircled{4} \times 13 = 52 \\ & = (4 \times 11) + 8 \\ & = \textcircled{8} \end{aligned}$$




$$\textcircled{4} \times 13 = 52$$

$$= (4 \times 11) + 8$$

$$= \textcircled{8}$$

you can multiply an
element of the group by
something that is NOT in
the group




$$\text{★} \times 13 = 52$$

$$= (4 \times 11) + 8$$

$$= \text{♣}$$

you can multiply an
element of the group by
something that is NOT in
the group



$\{a, b, c, \dots\}$ & '+'

identity element

there is an element 0 such that
 $0 + n = n$ for every n in the group

inverses

for every a in the group, there's
a b that makes $a + b = 0$ true

associativity

$$a + (b + c) = (a + b) + c$$


closure

If a and b are in the group and
 $a + b = c$, then c is in the group

3


Elliptic Curve Cryptography


$$y^2 \equiv x^3 + ax + b$$



$$y^2 \equiv x^3 + ax + b$$

where x and y
are in \mathbb{Z}_p








$$y^2 \equiv x^3 + ax + b$$

where x and y
are in \mathbb{Z}_p


and three collinear
points 'sum' to \mathbf{O}










$$a + b + c = \mathbf{O}$$








$$a + b + c = \text{O}$$




?????




$$a + b + c = \text{blue circle}$$



$$a + b + \text{blue circle} = \text{blue circle}$$




$$a + b + \mathbf{O} = \mathbf{O}$$



$$a + b + \text{blue circle} = \text{blue circle}$$

\Downarrow

$$a + b = \text{blue circle}$$




$$a + b + \text{blue circle} = \text{blue circle}$$

$$\Downarrow$$

$$a + b = \text{blue circle}$$

$$\Downarrow$$

$$a = -b$$



$$y^2 \equiv x^3 + x + 5$$

elliptic curve domain parameters over F_p

$$T = (p, a, b, G, n, h)$$

elliptic curve domain parameters over F_p

$$T = (p, a, b, G, n, h)$$

an integer defining
the field F_p

elliptic curve domain parameters over F_p

$$T = (p, a, b, G, n, h)$$

two elements of F_p defining

$$E: y^2 \equiv x^3 + ax + b$$

elliptic curve domain parameters over F_p

$$T = (p, a, b, G, n, h)$$

a point on $E(F_p)$ written as

$$G = (x_G, y_G)$$

elliptic curve domain parameters over F_p

$$T = (p, a, b, G, n, h)$$

the order of G in $E(F_p)$ – i.e.,

$$n \times G = \mathbf{O}$$

elliptic curve domain parameters over F_p

$$T = (p, a, b, G, n, h)$$


the cofactor of G in $E(F_p)$, which is
 $|E(F_p)| / n$

elliptic curve domain parameters over F_p


$$T = (p, a, b, G, n, h)$$

or more properly, $\text{orb}(G)$

the cofactor of G in $E(F_p)$, which is
 $|E(F_p)| / n$




$$y^2 \equiv x^3 + x + 5$$




$$y^2 \equiv x^3 + x + 5$$

$$T = (p, a, b, G, n, h)$$




$$y^2 \equiv x^3 + x + 5$$


$$T = (11, a, b, G, n, h)$$



$$y^2 \equiv x^3 + x + 5$$

$$T = (11, 1, 5, G, n, h)$$







$$y^2 \equiv x^3 + x + 5$$

$$T = (11, 1, 5, (0,4), 11, 1)$$


< worked example at the end >




$$y^2 \equiv x^3 + x + 5$$



$$y^2 \equiv x^3 + x + 5$$




$$y^2 \equiv x^3 + x + 5$$



$$y^2 \equiv x^3 + x + 5$$

$$y^2 \equiv x^3 + x + 5$$



$$1 \times G = (0, 4)$$

$$2 \times G = (5, 5)$$

$$3 \times G = (10, 5)$$

$$4 \times G = (2, -2)$$

$$5 \times G = (7, -5)$$

$$6 \times G = (7, 5)$$

$$7 \times G = (2, 2)$$

$$8 \times G = (10, -5)$$

$$9 \times G = (5, -5)$$

$$10 \times G = (0, -4)$$

$$11 \times G = (\infty, \infty)$$

$$1 \times G = (0, 4)$$

$$2 \times G = (5, 5)$$

$$3 \times G = (10, 5)$$

$$4 \times G = (2, -2)$$

$$5 \times G = (7, -5)$$

$$6 \times G = (7, 5)$$


$$7 \times G = (2, 2)$$

$$8 \times G = (10, -5)$$

$$9 \times G = (5, -5)$$

$$10 \times G = (0, -4)$$

$$11 \times G = (\infty, \infty)$$



comparison with RSA

comparison with RSA

smaller key size per security

comparison with RSA

smaller key size per security

smaller payload size

comparison with RSA

smaller key size per security

smaller payload size

faster computation





4

Quantum Computing & Shor's Algorithms

the Integer Factorisation problem

if $pq = N$ with p & q prime, find p and q given only N

the Integer Factorisation problem

if $pq = N$ with p & q prime, find p and q given only N

the Discrete Logarithm problem

if g generates a subgroup of a finite field F , and y is another member of F , find x such that $g^x = y$

the Integer Factorisation problem

if $pq = N$ with p & q prime, find p and q given only N

the Discrete Logarithm problem

if g generates a subgroup of a finite field F , and y is another member of F , find x such that $g^x = y$

the Elliptic Curve Discrete Logarithm problem

if G generates a subgroup of an elliptic curve over a field F , and P is another member of that elliptic curve, find k such that $P = kG$

Shor's order-finding algorithm

for a given number N , and any number a between 1 and N , we can find the smallest r such that $a^r \equiv 1 \pmod{N}$, in polynomial time

Shor's order-finding algorithm

Shor's order-finding algorithm

let $N = 323$. Choose $a = 11$.

Shor's algorithm gives us that $11^{48} \equiv 1 \pmod{323}$

Shor's order-finding algorithm

let $N = 323$. Choose $a = 11$.

Shor's algorithm gives us that $11^{48} \equiv 1 \pmod{323}$

$11^{48} - 1 \equiv 0 \pmod{323}$, so $(11^{24} - 1)(11^{24} + 1) \equiv 0 \pmod{323}$,
which is equivalent to $323 \mid (11^{24} - 1)(11^{24} + 1)$

Shor's order-finding algorithm

let $N = 323$. Choose $a = 11$.

Shor's algorithm gives us that $11^{48} \equiv 1 \pmod{323}$

$11^{48} - 1 \equiv 0 \pmod{323}$, so $(11^{24} - 1)(11^{24} + 1) \equiv 0 \pmod{323}$,
which is equivalent to $323 \mid (11^{24} - 1)(11^{24} + 1)$

we know 323 doesn't divide $11^{24} - 1$, or else we'd have

$$11^{24} \equiv 1 \pmod{323}$$

Shor's order-finding algorithm

let $N = 323$. Choose $a = 11$.

Shor's algorithm gives us that $11^{48} \equiv 1 \pmod{323}$

$11^{48} - 1 \equiv 0 \pmod{323}$, so $(11^{24} - 1)(11^{24} + 1) \equiv 0 \pmod{323}$,
which is equivalent to $323 \mid (11^{24} - 1)(11^{24} + 1)$

we know 323 doesn't divide $11^{24} - 1$, or else we'd have

$$11^{24} \equiv 1 \pmod{323}$$

so at least some of the factors of 323 must also
divide $11^{24} + 1$

Shor's order-finding algorithm

given that at least some of the factors of 323
must also divide $11^{24} + 1$

Shor's order-finding algorithm

given that at least some of the factors of 323
must also divide $11^{24} + 1$

calculate $\gcd(323, 11^{24} + 1) = 17$,
which is computationally efficient on classical computers

Shor's order-finding algorithm

given that at least some of the factors of 323
must also divide $11^{24} + 1$

calculate $\gcd(323, 11^{24} + 1) = 17$,
which is computationally efficient on classical computers

find that $17 \mid 323$ and $323 = 17 * 19$.

Shor's order-finding algorithm

given that at least some of the factors of 323
must also divide $11^{24} + 1$

calculate $\gcd(323, 11^{24} + 1) = 17$,
which is computationally efficient on classical computers

find that $17 \mid 323$ and $323 = 17 * 19$.

this breaks RSA!

the Integer Factorisation problem

if $pq = N$ with p & q prime, find p and q given only N

the Discrete Logarithm problem

if g generates a subgroup of a finite field F , and y is another member of F , find x such that $g^x = y$

the Elliptic Curve Discrete Logarithm problem

if G generates a subgroup of an elliptic curve over a field F , and P is another member of that elliptic curve, find k such that $P = kG$

~~the Integer Factorisation problem~~

~~if $pq = N$ with p & q prime, find p and q given only N~~

the Discrete Logarithm problem

if g generates a subgroup of a finite field F , and y is another member of F , find x such that $g^x = y$

the Elliptic Curve Discrete Logarithm problem

if G generates a subgroup of an elliptic curve over a field F , and P is another member of that elliptic curve, find k such that $P = kG$

~~the Integer Factorisation problem~~

~~if $pq = N$ with p & q prime, find p and q given only N~~

~~the Discrete Logarithm problem~~

~~if g generates a subgroup of a finite field F , and y is another member of F , find x such that $g^x = y$~~

~~the Elliptic Curve Discrete Logarithm problem~~

~~if G generates a subgroup of an elliptic curve over a field F , and P is another member of that elliptic curve, find k such that $P = kG$~~

~~the Integer Factorisation problem~~


~~if $pq = N$ with $p \leq \sqrt{N}$ and q given only N~~

~~the Discrete Logarithm problem~~

~~if g generates a subgroup of F and F , and y is another member of F , then that $g^x = y$~~

~~the Elliptic Curve Discrete Logarithm problem~~

~~if G generates a subgroup of an elliptic curve over a field F , and P is another member of that elliptic curve, find k such that $P = kG$~~



Post-quantum Cryptography

the isogeny-finding problem

given two elliptic curves between which we know there exists an isogeny, find the mapping that describes it

the isogeny-finding problem

given two elliptic curves between which we know there exists an isogeny, find the mapping that describes it

SIKE and SIDH, which are considered insecure

the isogeny-finding problem

given two elliptic curves between which we know there exists an isogeny, find the mapping that describes it

SIKE and SIDH, which are considered insecure

CSIDH

Quantum Security Analysis of CSIDH and Ordinary Isogeny-based Schemes

Xavier Bonnetain^{1,2} and André Schrottenloher²

¹ Sorbonne Université, Collège Doctoral, F-75005 Paris, France

² Inria, France

Abstract. CSIDH is a recent proposal by Castryck, Lange, Martindale, Panny and Renes for post-quantum non-interactive key-exchange. It is similar in design to a scheme by Couveignes, Rostovtsev and Stolbunov.

<https://who.rocq.inria.fr/Xavier.Bonnetain/pdfs/csidh-attack.pdf>

@eli@hachyderm.io

Quantum Security Analysis of CSIDH and Ordinary Isogeny-based Schemes

Xavier Bonnetain^{1,2} and André Schrottenloher²

¹ Sorbonne Université, Collège Doctoral, F-75005 Paris, France

² Inria, France

Abstract. CSIDH is a recent proposal by Castryck, Lange, Martindale, Panny and Renes for post-quantum non-interactive key-exchange. It is similar in design to a scheme by Couveignes, Rostovtsev and Stolbunov.

7 Conclusion

We presented a comprehensive quantum security assessment of CSIDH. In particular, when compared to the cost of a classical key-exchange, we showed that the parameters set in [6] actually seem to provide only around half of the expected security, as summarized in Table 7.

<https://who.rocq.inria.fr/Xavier.Bonnetain/pdfs/csidh-attack.pdf>

@eli@hachyderm.io

the isogeny-finding problem

given two elliptic curves between which we know there exists an isogeny, find the mapping that describes it

SIKE and SIDH, which are considered insecure

CSIDH, which should also be considered insecure


the Learning With Errors problem

introducing noise to encodings and using probability to decode

the Learning With Errors problem

introducing noise to encodings and using probability to decode

CRYSTALS-Kyber (key encapsulation) and
CRYSTALS-Dilithium (signatures)



<https://security.googleblog.com/2023/08/toward-quantum-resilient-security-keys.html>

In Chrome, you can now enable
'X25519Kyber768' for key exchange during TLS



OPEN QUANTUM SAFE

*software for prototyping
quantum-resistant cryptography*

<https://openquantumsafe.org/>

@eli@hachyderm.io

what I hope to see

what I hope to see

more diverse quantum-resilient cryptosystems

what I hope to see

more diverse quantum-resilient cryptosystems

quantum-resilient hardware tokens

what I hope to see

more diverse quantum-resilient cryptosystems

quantum-resilient hardware tokens

wider accessibility & rollout

wrapping up

how we got here

how we got here

RSA & ECDSA

how we got here

RSA & ECDSA

...and how quantum breaks them

how we got here

RSA & ECDSA

...and how quantum breaks them

what's next



Asymmetric Cryptography: A Deep Dive

Eli Holderness
@eli@hachyderm.io
[they/them/theirs](#)

sources: history

<https://www.redhat.com/en/blog/brief-history-cryptography>

sources: RSA + group theory

<https://ee.stanford.edu/~hellman/publications/24.pdf>

<https://weakdh.org/imperfect-forward-secrecy-ccs15.pdf>

[https://en.wikipedia.org/wiki/Padding_\(cryptography\)](https://en.wikipedia.org/wiki/Padding_(cryptography))

sources: ECC

<https://scholar.rose-hulman.edu/cgi/viewcontent.cgi?article=1389&context=rhumj>

<http://koclab.cs.ucsb.edu/teaching/ecc/eccPapers/Washington-ch04.pdf>

<http://www.secg.org/sec2-v2.pdf>

sources: QC & Shor

<https://research.kudelskisecurity.com/2021/08/24/quantum-attack-resource-estimate-using-shors-algorithm-to-break-rsa-vs-dh-dsa-vs-ecc/>

<https://arxiv.org/pdf/quant-ph/9508027.pdf>

<https://www.omnicalculator.com/math/power-modulo>

sources: PQC

<https://security.googleblog.com/2023/08/toward-quantum-resilient-security-keys.html>

<https://csidh.isogeny.org/>

<https://sike.org/>

<https://eprint.iacr.org/2019/725>

<https://blog.chromium.org/2023/08/protecting-chrome-traffic-with-hybrid.html>

<https://www.ietf.org/archive/id/draft-tls-westerbaan-xyber768d00-02.html>

<https://openquantumsafe.org/>

<https://eprint.iacr.org/2022/1225.pdf>

<https://github.com/signalapp/libsignal/commit/ff09619432e19e96231ebed913fe4433f26ee0d2>

worked example with $T = (11, 1, 5, (0,4), 11, 1)$

worked example with $T = (11, 1, 5, (0,4), 11, 1)$

$$d_{PK} = 3$$

Pick a random number d_{PK} from $[1, \dots, n-1] = [1, \dots, 10]$.

Let's pick 3. This is our private key.

worked example with $T = (11, 1, 5, (0,4), 11, 1)$

$$d_{PK} = 3 \quad Q_{PK} = (10, 5)$$

Pick a random number d_{PK} from $[1, \dots, n-1] = [1, \dots, 10]$.

Let's pick 3. This is our private key.

Calculate $Q_{PK} = d_{PK} \times G$, which in our case is
 $3 \times (0,4) = (10, 5)$. This is our public curve point.

worked example with $T = (11, 1, 5, (0,4), 11, 1)$
 $d_{PK} = 3$ $Q_{PK} = (10, 5)$

We have some binary message, e , to sign. Let's say we want to sign the message 01001110 01000100 01000011.

worked example with $T = (11, 1, 5, (0,4), 11, 1)$

$$z = 3 \quad d_{PK} = 3 \quad Q_{PK} = (10, 5)$$

We have some binary message, e , to sign. Let's say we want to sign the message 01001110 01000100 01000011.

The size of our group is 11, or 1101 in binary - 4 bits long.

Take the last 4 bits of our message: 0011. Call it z .

worked example with $T = (11, 1, 5, (0,4), 11, 1)$

$$z = 3 \quad d_{PK} = 3 \quad Q_{PK} = (10, 5)$$

Pick another random number k from $[1, \dots, n-1]$. This time let's choose 5. This must be random per signature.

worked example with $T = (11, 1, 5, (0,4), 11, 1)$

$$k^{-1} = 9 \quad z = 3 \quad d_{\text{PK}} = 3 \quad Q_{\text{PK}} = (10, 5)$$

Pick another random number k from $[1, \dots, n-1]$. This time let's choose 5. This must be random per signature.

Find its inverse k^{-1} in \mathbf{F}_{11} , which is 9.

worked example with $T = (11, 1, 5, (0,4), 11, 1)$

$$x_k = 7, y_k = -5 \quad k^{-1} = 9 \quad z = 3 \quad d_{PK} = 3 \quad Q_{PK} = (10, 5)$$

Pick another random number k from $[1, \dots, n-1]$. This time let's choose 5. This must be random per signature.

Find its inverse k^{-1} in \mathbf{F}_{11} , which is 9.

Calculate $k \times G = 5 \times (0,4) = (7, -5)$. Take its coordinates, so we have $x_k = 7, y_k = -5$

worked example with $T = (11, 1, 5, (0,4), 11, 1)$

$$x_k = 7, y_k = -5 \quad k^{-1} = 9 \quad z = 3 \quad d_{PK} = 3 \quad Q_{PK} = (10, 5)$$

Now calculate r and s , where

$$r \equiv x_k \bmod n \text{ and } s \equiv k^{-1}(z + r * d_{PK}) \bmod n$$

worked example with $T = (11, 1, 5, (0,4), 11, 1)$

$$x_k = 7, y_k = -5 \quad k^{-1} = 9 \quad z = 3 \quad d_{PK} = 3 \quad Q_{PK} = (10, 5)$$

Now calculate r and s , where

$$r \equiv x_k \bmod n \text{ and } s \equiv k^{-1}(z + r * d_{PK}) \bmod n$$

This gives us $r = 7$ and $s = 7$, and this is our signature:

$$(r,s) = (7, 7).$$

worked example with $T = (11, 1, 5, (0,4), 11, 1)$

$$x_k = 7, y_k = -5 \quad k^{-1} = 9 \quad z = 3 \quad d_{PK} = 3 \quad Q_{PK} = (10, 5)$$

Now calculate r and s , where

$$r \equiv x_k \bmod n \text{ and } s \equiv k^{-1}(z + r * d_{PK}) \bmod n$$

This gives us $r = 7$ and $s = 7$, and this is our signature:

$$(r,s) = (7, 7).$$

If either r or s are 0, we have to go back and pick a different k .

worked example with $T = (11, 1, 5, (0,4), 11, 1)$

$$x_k = 7, y_k = -5 \quad k^{-1} = 9 \quad z = 3 \quad d_{PK} = 3 \quad Q_{PK} = (10, 5)$$

We've now generated a signature $(r,s) = (7, 7)$ over the
binary message 01001110 01000100 01000011.

Let's verify it!

worked example with $T = (11, 1, 5, (0,4), 11, 1)$
 $r = 7, s = 7$ $Q_{PK} = (10, 5)$

worked example with $T = (11, 1, 5, (0,4), 11, 1)$
 $z = 3$ $r = 7, s = 7$ $Q_{PK} = (10, 5)$

We have the message, 01001110 01000100 01000011.
Take the last 4 bits as we did before to get $z = 3$.

worked example with $T = (11, 1, 5, (0,4), 11, 1)$

$$u_1 = 2, u_2 = 5 \quad z = 3 \quad r = 7, s = 7 \quad Q_{PK} = (10, 5)$$

We have the message, 01001110 01000100 01000011.

Take the last 4 bits as we did before to get $z = 3$.

Calculate $u_1 \equiv zs^{-1} \pmod{n}$: $u_1 \equiv 3*8 \equiv 2 \pmod{11}$

Calculate $u_2 \equiv rs^{-1} \pmod{n}$: $u_2 \equiv 7*7 \equiv 5 \pmod{11}$

worked example with $T = (11, 1, 5, (0,4), 11, 1)$

$$u_1 = 2, u_2 = 5 \quad z = 3 \quad r = 7, s = 7 \quad Q_{PK} = (10, 5)$$

Calculate a new point on the curve, $(x, y) = u_1 \times G + u_2 \times Q_{PK}$

worked example with $T = (11, 1, 5, (0,4), 11, 1)$

$$u_1 = 2, u_2 = 5 \quad z = 3 \quad r = 7, s = 7 \quad Q_{PK} = (10, 5)$$

Calculate a new point on the curve, $(x, y) = u_1 \times G + u_2 \times$

$$u_1 \times G = 2 \times (0,4)$$

$$u_2 \times Q_{PK} = 5 \times (10,5) = 5 \times (3 \times (0,4)) = 4 \times (0,4)$$

worked example with $T = (11, 1, 5, (0,4), 11, 1)$

$$u_1 = 2, u_2 = 5 \quad z = 3 \quad r = 7, s = 7 \quad Q_{PK} = (10, 5)$$

Calculate a new point on the curve, $(x, y) = u_1 \times G + u_2 \times$

$$u_1 \times G = 2 \times Q_{PK} \times (0,4)$$

$$u_2 \times Q_{PK} = 5 \times (10,5) = 5 \times (3 \times (0,4)) = 4 \times (0,4)$$

$$\text{so } (x, y) = 2 \times (0,4) + 4 \times (0,4) = 6 \times (0,4) = (7,5)$$

worked example with $T = (11, 1, 5, (0,4), 11, 1)$

$$u_1 = 2, u_2 = 5 \quad z = 3 \quad r = 7, s = 7 \quad Q_{PK} = (10, 5)$$

Calculate a new point on the curve, $(x, y) = u_1 \times G + u_2 \times$

$$u_1 \times G = 2 \times Q_{PK} \times (0,4)$$

$$u_2 \times Q_{PK} = 5 \times (10,5) = 5 \times (3 \times (0,4)) = 4 \times (0,4)$$

$$\text{so } (x, y) = 2 \times (0,4) + 4 \times (0,4) = 6 \times (0,4) = (7,5)$$

The signature is valid if $x = r \bmod n$, which it is!

