

Question	Answer	Marks	Guidance
(a)	Use chain rule at least once	M1	Needs $\frac{dy}{dt} = \frac{1}{\tan t} \frac{d}{dt}(\tan t)$ or $\frac{dx}{dt} = (-1)(\cos^{-2}t) \frac{d}{dt}(\cos t)$. BOD if + and $(-1)(-1)$ not seen. $\frac{dx}{dt} = \sec t \tan t$ (from List of Formulae MF19) M1 A1. If $\frac{dx}{dt} = -\sec t \tan t$ M1 A0.
	Obtain $\frac{dx}{dt} = \sec t \tan t$	A1	OE e.g. $\sin t (\cos t)^{-2}$. If e.g. $\frac{dx}{dt} = \sec x \tan x$ or $\sec \theta \tan \theta$ or $\sec t \tan x$, condone recovery on next line.
	Obtain $\frac{dy}{dt} = \frac{\sec^2 t}{\tan t}$	A1	OE e.g. $\frac{1}{\sin t \cos t}$. If e.g. $\frac{dy}{dt} = \frac{\sec^2 x}{\tan x}$ or $\frac{\sec^2 \theta}{\tan \theta}$, condone recovery on next line. Only penalise notation errors once in $\frac{dx}{dt}$ and $\frac{dy}{dt}$ if no recovery.
	Use $\frac{dy}{dx} = \frac{dy}{dt} \div \frac{dx}{dt}$	M1	Allow even if previous M0 scored, but must be using derivatives.
	Obtain given answer $\frac{\cos t}{\sin^2 t}$	A1	AG After $\frac{dy}{dx} = \frac{dy}{dt} \div \frac{dx}{dt}$ used, any notation error A0. Must cancel $\cos t$ correctly.
		5	

Question	Answer	Marks	Guidance
(b)	State or imply $t = \frac{1}{4}\pi$ when $y = 0$	B1	
	Form the equation of the tangent at $y = 0$ or find c	M1	$x = \sqrt{2}$, $\frac{dy}{dx} = \sqrt{2}$ and $y = 0$, <i>their</i> coordinates and gradient used in $y = mx + c$.
	Obtain answer $y = \sqrt{2}x - 2$	A1	OE e.g. $y = \sqrt{2}(x - \sqrt{2})$ ISW. Allow $y = 1.41x - 2[.00]$ or $1.41(x - 1.41)$.
		3	

