

# Quantitative Methods

## Human Sciences, 2020–2021

Elias Nosrati

Lecture 2: 22 October 2020

Today

# Today

- ▶ Recap on probability and counting: the birthday problem.

# Today

- ▶ Recap on probability and counting: the birthday problem.
- ▶ Introduction to conditional probability.

# Today

- ▶ Recap on probability and counting: the birthday problem.
- ▶ Introduction to conditional probability.
- ▶ Problem sheet 1 (tutorial).

# The birthday problem

# The birthday problem

- ▶ There are  $k$  people in a room.

# The birthday problem

- ▶ There are  $k$  people in a room.
- ▶ Assume each person's birthday is equally likely to be any of the 365 days of the year and assume people's birthdays are independent.



# The birthday problem

- ▶ There are  $k$  people in a room.
- ▶ Assume each person's birthday is equally likely to be any of the 365 days of the year and assume people's birthdays are independent.
- ▶ What is the probability that at least one pair of people in the group have the same birthday?

## The birthday problem

- ▶ There are  $k$  people in a room.
- ▶ Assume each person's birthday is equally likely to be any of the 365 days of the year and assume people's birthdays are independent.
- ▶ What is the probability that at least one pair of people in the group have the same birthday?
- ▶ Hint: Recall that  $\mathbb{P}(A) = 1 - \mathbb{P}(A^c)$ .

## The birthday problem: solution

## The birthday problem: solution

- ▶ Total number of ways of assigning birthdays to  $n$  people (sampling from 365 days *with* replacement):  $365^n$ .

## The birthday problem: solution

- ▶ Total number of ways of assigning birthdays to  $n$  people (sampling from 365 days *with* replacement):  $365^n$ .
- ▶ Complement: the number of ways to assign birthdays to  $n$  people such that no two people share the same birthday (sampling from 365 days *without* replacement):

$$365 \times 364 \times \cdots \times (365 - n + 1)$$

for  $n \leq 365$ .

## The birthday problem: solution

- ▶ Total number of ways of assigning birthdays to  $n$  people (sampling from 365 days *with* replacement):  $365^n$ .
- ▶ Complement: the number of ways to assign birthdays to  $n$  people such that no two people share the same birthday (sampling from 365 days *without* replacement):

$$365 \times 364 \times \cdots \times (365 - n + 1)$$

for  $n \leq 365$ .

- ▶  $\mathbb{P}(\text{no birthday match}) = \frac{365 \times 364 \times \cdots \times (365 - n + 1)}{365^n}$ .

## The birthday problem: solution

- ▶ Total number of ways of assigning birthdays to  $n$  people (sampling from 365 days *with* replacement):  $365^n$ .
- ▶ Complement: the number of ways to assign birthdays to  $n$  people such that no two people share the same birthday (sampling from 365 days *without* replacement):

$$365 \times 364 \times \cdots \times (365 - n + 1)$$

for  $n \leq 365$ .

- ▶  $\mathbb{P}(\text{no birthday match}) = \frac{365 \times 364 \times \cdots \times (365 - n + 1)}{365^n}$ .
- ▶  $\mathbb{P}(\text{at least one birthday match}) = 1 - \mathbb{P}(\text{no birthday match})$ .

## The birthday problem: solution

- ▶ Total number of ways of assigning birthdays to  $n$  people (sampling from 365 days *with* replacement):  $365^n$ .
- ▶ Complement: the number of ways to assign birthdays to  $n$  people such that no two people share the same birthday (sampling from 365 days *without* replacement):

$$365 \times 364 \times \cdots \times (365 - n + 1)$$

for  $n \leq 365$ .

- ▶  $\mathbb{P}(\text{no birthday match}) = \frac{365 \times 364 \times \cdots \times (365 - n + 1)}{365^n}$ .
- ▶  $\mathbb{P}(\text{at least one birthday match}) = 1 - \mathbb{P}(\text{no birthday match})$ .
- ▶ In this room,  $\mathbb{P}(\text{at least one birthday match}) \approx 4\%$ .



## The birthday problem in R

```
# Create a function  
pmatch <- function(n) {  
  1 - prod(365:(365 - n + 1)) / (365 ^ n)  
}
```

# The birthday problem in R

```
# Create a function
pmatch <- function(n) {
  1 - prod(365:(365 - n + 1)) / (365 ^ n)
}
```

```
# Placeholder
probs <- NULL

# For loop
for (i in 1:70) {
  probs[i] <- pmatch(i)
}
```

# The birthday problem in R

```
# Create a function  
pmatch <- function(n) {  
  1 - prod(365:(365 - n + 1)) / (365 ^ n)  
}
```

```
# Placeholder  
probs <- NULL  
  
# For loop  
for (i in 1:70) {  
  probs[i] <- pmatch(i)  
}
```

```
# Alternative method  
probs <- sapply(1:70, pmatch)
```

## The birthday problem in **R** (cont.)

```
save <- data.frame("n" = 1:70, "prob" = probs)
```

## The birthday problem in **R** (cont.)

```
save <- data.frame("n" = 1:70, "prob" = probs)
```


```
head(save)
```

##	n	prob
## 1	1	0.000000000
## 2	2	0.002739726
## 3	3	0.008204166
## 4	4	0.016355912
## 5	5	0.027135574
## 6	6	0.040462484

## The birthday problem in **R** (cont.)

```
ggplot(save, aes(n, prob)) +  
  labs( # Axis labels  
    x = "Number of people in the room",  
    y = "Probability of same birthday") +  
  scale_x_continuous( # Modify X-axis  
    breaks = seq(0, 70, 5)) +  
  scale_y_continuous( # Modify Y-axis  
    breaks = seq(0, 1, 0.1),  
    label = scales::percent) +  
  geom_point( # Modify points  
    size = 2,  
    colour = "darkred",  
    alpha = 0.7) +  
  geom_hline( # When is P(match) > 0.5 or 0.9?  
    yintercept = c(0.5, 0.9), linetype = "dashed") +  
  theme_minimal() # Remove redundant lines
```

Probability of same birthday



## Conditional probability: motivation



# Conditional probability: motivation

- ▶ Probability is a means of quantifying uncertainty about events.

## Conditional probability: motivation

- ▶ Probability is a means of quantifying uncertainty about events.
- ▶ Whenever new evidence is observed, we acquire information that may affect our uncertainties.

## Conditional probability: motivation

- ▶ Probability is a means of quantifying uncertainty about events.
- ▶ Whenever new evidence is observed, we acquire information that may affect our uncertainties.
- ▶ Conditional probability allows us to update our beliefs in light of new evidence.

## Conditional probability: motivation

- ▶ Probability is a means of quantifying uncertainty about events.
- ▶ Whenever new evidence is observed, we acquire information that may affect our uncertainties.
- ▶ Conditional probability allows us to update our beliefs in light of new evidence.
- ▶ “Conditioning is the soul of statistics” (Blitzstein and Hwang, 2019: 46).

# Defining conditional probability

## Definition

If  $A$  and  $B$  are events, with  $\mathbb{P}(B) > 0$ , then the *conditional probability* of  $A$  given  $B$  is

$$\mathbb{P}(A \mid B) = \frac{\mathbb{P}(A \cap B)}{\mathbb{P}(B)}.$$

# Defining conditional probability

## Definition

If  $A$  and  $B$  are events, with  $\mathbb{P}(B) > 0$ , then the *conditional probability* of  $A$  given  $B$  is

$$\mathbb{P}(A \mid B) = \frac{\mathbb{P}(A \cap B)}{\mathbb{P}(B)}.$$

- ▶  $A$  is the event whose uncertainty we want to update and  $B$  is the (new) evidence we observe.

# Defining conditional probability

## Definition

If  $A$  and  $B$  are events, with  $\mathbb{P}(B) > 0$ , then the *conditional probability* of  $A$  given  $B$  is

$$\mathbb{P}(A \mid B) = \frac{\mathbb{P}(A \cap B)}{\mathbb{P}(B)}.$$

- ▶  $A$  is the event whose uncertainty we want to update and  $B$  is the (new) evidence we observe.
- ▶  $\mathbb{P}(A)$  is called the *prior* probability of  $A$  (*before* updating beliefs based on new evidence).

# Defining conditional probability

## Definition

If  $A$  and  $B$  are events, with  $\mathbb{P}(B) > 0$ , then the *conditional probability* of  $A$  given  $B$  is

$$\mathbb{P}(A \mid B) = \frac{\mathbb{P}(A \cap B)}{\mathbb{P}(B)}.$$

- ▶  $A$  is the event whose uncertainty we want to update and  $B$  is the (new) evidence we observe.
- ▶  $\mathbb{P}(A)$  is called the *prior* probability of  $A$  (*before* updating beliefs based on new evidence).
- ▶  $\mathbb{P}(A \mid B)$  is called the *posterior* of  $A$  (*after* updating beliefs based on new evidence).



# Defining conditional probability

## Definition


If  $A$  and  $B$  are events, with  $\mathbb{P}(B) > 0$ , then the *conditional probability* of  $A$  given  $B$  is

$$\mathbb{P}(A \mid B) = \frac{\mathbb{P}(A \cap B)}{\mathbb{P}(B)}.$$

- ▶  $A$  is the event whose uncertainty we want to update and  $B$  is the (new) evidence we observe.
- ▶  $\mathbb{P}(A)$  is called the *prior* probability of  $A$  (*before* updating beliefs based on new evidence).
- ▶  $\mathbb{P}(A \mid B)$  is called the *posterior* of  $A$  (*after* updating beliefs based on new evidence).
- ▶ Note that  $\mathbb{P}(A \mid A) = 1$ .


## Visualising a conditional probability

Suppose we have a sample space  $S$  and two events  $A$  and  $B$ :



## Visualising a conditional probability


Suppose we have a sample space  $S$  and two events  $A$  and  $B$ :



- Suppose that  $B$  occurred: get rid of all elements of  $B^c$ .

## Visualising a conditional probability


Suppose we have a sample space  $S$  and two events  $A$  and  $B$ :



- ▶ Suppose that  $B$  occurred: get rid of all elements of  $B^c$ .
- ▶ The only probability measure remaining that we can assign to  $A$  is  $\mathbb{P}(A \cap B)$ .

## Visualising a conditional probability


Suppose we have a sample space  $S$  and two events  $A$  and  $B$ :



- ▶ Suppose that  $B$  occurred: get rid of all elements of  $B^c$ .
- ▶ The only probability measure remaining that we can assign to  $A$  is  $\mathbb{P}(A \cap B)$ .
- ▶ *Renormalise*: create a new probability space that assigns an updated probability measure to each possible event such that all the probabilities add up to 1.

## Visualising a conditional probability

Suppose we have a sample space  $S$  and two events  $A$  and  $B$ :




- ▶ Suppose that  $B$  occurred: get rid of all elements of  $B^c$ .
- ▶ The only probability measure remaining that we can assign to  $A$  is  $\mathbb{P}(A \cap B)$ .
- ▶ *Renormalise*: create a new probability space that assigns an updated probability measure to each possible event such that all the probabilities add up to 1.
- ▶ Divide by  $\mathbb{P}(B)$ , the total mass of the outcomes in  $B$ .

## Visualising a conditional probability (cont.)

The updated probability measure assigned to the event  $A$  is the conditional probability


$$\mathbb{P}(A \mid B) = \mathbb{P}(A \cap B) / \mathbb{P}(B).$$



## Visualising a conditional probability (cont.)

The updated probability measure assigned to the event  $A$  is the conditional probability

$$\mathbb{P}(A \mid B) = \mathbb{P}(A \cap B) / \mathbb{P}(B).$$




- We have updated our probabilities in accordance with observed evidence.



## Visualising a conditional probability (cont.)

The updated probability measure assigned to the event  $A$  is the conditional probability

$$\mathbb{P}(A \mid B) = \mathbb{P}(A \cap B) / \mathbb{P}(B).$$




- ▶ We have updated our probabilities in accordance with observed evidence.
- ▶ Outcomes that contradict the evidence are discarded.

## Visualising a conditional probability (cont.)

The updated probability measure assigned to the event  $A$  is the conditional probability

$$\mathbb{P}(A \mid B) = \mathbb{P}(A \cap B) / \mathbb{P}(B).$$



- ▶ We have updated our probabilities in accordance with observed evidence.
- ▶ Outcomes that contradict the evidence are discarded.
- ▶ Relative measures of uncertainty are redistributed amongst remaining possible outcomes.

# Frequentist interpretation of conditional probability

## Frequentist interpretation of conditional probability

- ▶ Frequentist interpretation of probability: relative frequency over a large number of repeated trials.

## Frequentist interpretation of conditional probability

- ▶ Frequentist interpretation of probability: relative frequency over a large number of repeated trials.
- ▶  $\mathbb{P}(A \mid B)$  is the fraction of times  $A$  occurs, restricting attention to trials where  $B$  occurs.

### Example

- ▶ We flip a coin several times, record the resulting strings of Heads and Tails, and repeat the experiment  $n$  times.

## Frequentist interpretation of conditional probability

- ▶ Frequentist interpretation of probability: relative frequency over a large number of repeated trials.
- ▶  $\mathbb{P}(A \mid B)$  is the fraction of times  $A$  occurs, restricting attention to trials where  $B$  occurs.

### Example

- ▶ We flip a coin several times, record the resulting strings of Heads and Tails, and repeat the experiment  $n$  times.
- ▶  $B$  is the event that the first flip is Heads,  $A$  is the event that the second flip is Tails.

## Frequentist interpretation of conditional probability

- ▶ Frequentist interpretation of probability: relative frequency over a large number of repeated trials.
- ▶  $\mathbb{P}(A \mid B)$  is the fraction of times  $A$  occurs, restricting attention to trials where  $B$  occurs.

### Example

- ▶ We flip a coin several times, record the resulting strings of Heads and Tails, and repeat the experiment  $n$  times.
- ▶  $B$  is the event that the first flip is Heads,  $A$  is the event that the second flip is Tails.
- ▶ We condition on  $B$  by isolating all the times where  $B$  occurs ( $n_B$ ) and look at the fraction of times  $A$  also occurs ( $n_{AB}$ ).

## Frequentist interpretation of conditional probability

- ▶ Frequentist interpretation of probability: relative frequency over a large number of repeated trials.
- ▶  $\mathbb{P}(A \mid B)$  is the fraction of times  $A$  occurs, restricting attention to trials where  $B$  occurs.

### Example

- ▶ We flip a coin several times, record the resulting strings of Heads and Tails, and repeat the experiment  $n$  times.
- ▶  $B$  is the event that the first flip is Heads,  $A$  is the event that the second flip is Tails.
- ▶ We condition on  $B$  by isolating all the times where  $B$  occurs ( $n_B$ ) and look at the fraction of times  $A$  also occurs ( $n_{AB}$ ).
- ▶ Thus

$$\mathbb{P}(A \mid B) = n_{AB}/n_B = (n_{AB}/n)/(n_B/n) = \mathbb{P}(A \cap B)/\mathbb{P}(B).$$



# Joint probability and conditional probability

## Theorem

For any events  $A$  and  $B$  with positive probabilities,

$$\mathbb{P}(A \cap B) = \mathbb{P}(B)\mathbb{P}(A \mid B) = \mathbb{P}(A)\mathbb{P}(B \mid A).$$

# Joint probability and conditional probability

## Theorem

For any events  $A$  and  $B$  with positive probabilities,

$$\mathbb{P}(A \cap B) = \mathbb{P}(B)\mathbb{P}(A \mid B) = \mathbb{P}(A)\mathbb{P}(B \mid A).$$

This can be generalised to the intersection of  $n$  events:

$$\mathbb{P}(A_1 \cap \cdots \cap A_n) = \mathbb{P}(A_1)\mathbb{P}(A_2|A_1)\mathbb{P}(A_3|A_1 \cap A_2) \cdots \mathbb{P}(A_n|A_1 \cap \cdots \cap A_{n-1}).$$

# Bayes' Rule

# Bayes' Rule

## Theorem

For any events  $A$  and  $B$  with positive probabilities,

$$\mathbb{P}(A \mid B) = \frac{\mathbb{P}(B \mid A)\mathbb{P}(A)}{\mathbb{P}(B)}.$$

# Bayes' Rule

## Theorem

For any events  $A$  and  $B$  with positive probabilities,

$$\mathbb{P}(A \mid B) = \frac{\mathbb{P}(B \mid A)\mathbb{P}(A)}{\mathbb{P}(B)}.$$



This follows immediately from the previous Theorem (which in turn follows immediately from the definition of conditional probability).

## The Law of Total Probability (revisited)

## The Law of Total Probability (revisited)

Recall that, for any two events  $A$  and  $B$ , the Law of Total Probability states that

$$\mathbb{P}(A) = \mathbb{P}(A \cap B) + \mathbb{P}(A \cap B^c).$$



## The Law of Total Probability (revisited)

Viewed differently, the Law of Total Probability relates conditional probability to unconditional probability:



## The Law of Total Probability (revisited)

Viewed differently, the Law of Total Probability relates conditional probability to unconditional probability:

### Theorem

Let  $A_1, \dots, A_n$  be a partition of the sample space  $S$  (quiz: what is a partition again?),

## The Law of Total Probability (revisited)

Viewed differently, the Law of Total Probability relates conditional probability to unconditional probability:

### Theorem

Let  $A_1, \dots, A_n$  be a partition of the sample space  $S$  (quiz: what is a partition again?), with  $\mathbb{P}(A_i) > 0$  for all  $i$ . Then, for any event  $B$ ,

$$\mathbb{P}(B) = \sum_{i=1}^n \mathbb{P}(B \mid A_i) \mathbb{P}(A_i).$$


# The Law of Total Probability (revisited)

Viewed differently, the Law of Total Probability relates conditional probability to unconditional probability:

## Theorem

Let  $A_1, \dots, A_n$  be a partition of the sample space  $S$  (quiz: what is a partition again?), with  $\mathbb{P}(A_i) > 0$  for all  $i$ . Then, for any event  $B$ ,

$$\mathbb{P}(B) = \sum_{i=1}^n \mathbb{P}(B \mid A_i) \mathbb{P}(A_i).$$



Example: testing for a rare disease

## Example: testing for a rare disease

- ▶ Consider a rare disease  $D$  that affects 1% of the population.

## Example: testing for a rare disease

- ▶ Consider a rare disease  $D$  that affects 1% of the population.
- ▶ Suppose there is a test  $T$  that is 95% accurate such that

$$\mathbb{P}(T \mid D) = 0.95 \quad \text{and} \quad \mathbb{P}(T^c \mid D^c) = 0.95.$$

## Example: testing for a rare disease

- ▶ Consider a rare disease  $D$  that affects 1% of the population.
- ▶ Suppose there is a test  $T$  that is 95% accurate such that

$$\mathbb{P}(T \mid D) = 0.95 \quad \text{and} \quad \mathbb{P}(T^c \mid D^c) = 0.95.$$

- ▶ A randomly selected person takes the test and tests positive. What is the probability that this person has the disease?

## Example: testing for a rare disease

- ▶ Consider a rare disease  $D$  that affects 1% of the population.
- ▶ Suppose there is a test  $T$  that is 95% accurate such that

$$\mathbb{P}(T \mid D) = 0.95 \quad \text{and} \quad \mathbb{P}(T^c \mid D^c) = 0.95.$$

- ▶ A randomly selected person takes the test and tests positive.  
What is the probability that this person has the disease?

*Solution:* Apply Bayes' Rule and the Law of Total Probability:

$$\mathbb{P}(D \mid T) = \frac{\mathbb{P}(T \mid D)\mathbb{P}(D)}{\mathbb{P}(T)}$$



## Example: testing for a rare disease

- ▶ Consider a rare disease  $D$  that affects 1% of the population.
- ▶ Suppose there is a test  $T$  that is 95% accurate such that

$$\mathbb{P}(T \mid D) = 0.95 \quad \text{and} \quad \mathbb{P}(T^c \mid D^c) = 0.95.$$

- ▶ A randomly selected person takes the test and tests positive.  
What is the probability that this person has the disease?

*Solution:* Apply Bayes' Rule and the Law of Total Probability:

$$\begin{aligned}\mathbb{P}(D \mid T) &= \frac{\mathbb{P}(T \mid D)\mathbb{P}(D)}{\mathbb{P}(T)} \\ &= \frac{\mathbb{P}(T \mid D)\mathbb{P}(D)}{\mathbb{P}(T \mid D)\mathbb{P}(D) + \mathbb{P}(T \mid D^c)\mathbb{P}(D^c)}\end{aligned}$$

## Example: testing for a rare disease

- ▶ Consider a rare disease  $D$  that affects 1% of the population.
- ▶ Suppose there is a test  $T$  that is 95% accurate such that

$$\mathbb{P}(T \mid D) = 0.95 \quad \text{and} \quad \mathbb{P}(T^c \mid D^c) = 0.95.$$

- ▶ A randomly selected person takes the test and tests positive.  
What is the probability that this person has the disease?

*Solution:* Apply Bayes' Rule and the Law of Total Probability:

$$\begin{aligned}\mathbb{P}(D \mid T) &= \frac{\mathbb{P}(T \mid D)\mathbb{P}(D)}{\mathbb{P}(T)} \\ &= \frac{\mathbb{P}(T \mid D)\mathbb{P}(D)}{\mathbb{P}(T \mid D)\mathbb{P}(D) + \mathbb{P}(T \mid D^c)\mathbb{P}(D^c)} \\ &= \frac{0.95 \times 0.01}{(0.95 \times 0.01) + (0.05 \times 0.99)}\end{aligned}$$

## Example: testing for a rare disease

- ▶ Consider a rare disease  $D$  that affects 1% of the population.
- ▶ Suppose there is a test  $T$  that is 95% accurate such that

$$\mathbb{P}(T \mid D) = 0.95 \quad \text{and} \quad \mathbb{P}(T^c \mid D^c) = 0.95.$$

- ▶ A randomly selected person takes the test and tests positive.  
What is the probability that this person has the disease?

*Solution:* Apply Bayes' Rule and the Law of Total Probability:

$$\begin{aligned}\mathbb{P}(D \mid T) &= \frac{\mathbb{P}(T \mid D)\mathbb{P}(D)}{\mathbb{P}(T)} \\&= \frac{\mathbb{P}(T \mid D)\mathbb{P}(D)}{\mathbb{P}(T \mid D)\mathbb{P}(D) + \mathbb{P}(T \mid D^c)\mathbb{P}(D^c)} \\&= \frac{0.95 \times 0.01}{(0.95 \times 0.01) + (0.05 \times 0.99)} \\&\approx 16\%.\end{aligned}$$

Conditional probabilities are probabilities

Conditional probabilities are probabilities

$$\mathbb{P}(\cdot \mid E) : S \rightarrow [0, 1].$$

## Conditional probabilities are probabilities

$$\mathbb{P}(\cdot \mid E) : S \rightarrow [0, 1].$$

By conditioning on some event  $E$ , we have

## Conditional probabilities are probabilities

$$\mathbb{P}(\cdot \mid E) : S \rightarrow [0, 1].$$

By conditioning on some event  $E$ , we have

- ▶  $\mathbb{P}(A \mid E) \geq 0$  for any event  $A$ .

## Conditional probabilities are probabilities

$$\mathbb{P}(\cdot \mid E) : S \rightarrow [0, 1].$$

By conditioning on some event  $E$ , we have

- ▶  $\mathbb{P}(A \mid E) \geq 0$  for any event  $A$ .
- ▶  $\mathbb{P}(\emptyset \mid E) = 0$ ,  $\mathbb{P}(S \mid E) = 1$ .



## Conditional probabilities are probabilities

$$\mathbb{P}(\cdot \mid E) : S \rightarrow [0, 1].$$

By conditioning on some event  $E$ , we have

- ▶  $\mathbb{P}(A \mid E) \geq 0$  for any event  $A$ .
- ▶  $\mathbb{P}(\emptyset \mid E) = 0$ ,  $\mathbb{P}(S \mid E) = 1$ .
- ▶ If  $A_1, \dots, A_n$  are disjoint, then  $\mathbb{P}(\cup_i A_i \mid E) = \sum_i \mathbb{P}(A_i \mid E)$ .

## Conditional probabilities are probabilities

$$\mathbb{P}(\cdot \mid E) : S \rightarrow [0, 1].$$

By conditioning on some event  $E$ , we have

- ▶  $\mathbb{P}(A \mid E) \geq 0$  for any event  $A$ .
- ▶  $\mathbb{P}(\emptyset \mid E) = 0$ ,  $\mathbb{P}(S \mid E) = 1$ .
- ▶ If  $A_1, \dots, A_n$  are disjoint, then  $\mathbb{P}(\cup_i A_i \mid E) = \sum_i \mathbb{P}(A_i \mid E)$ .

Conclusion:

- ▶ Conditional probabilities are probabilities.

## Conditional probabilities are probabilities

$$\mathbb{P}(\cdot \mid E) : S \rightarrow [0, 1].$$

By conditioning on some event  $E$ , we have

- ▶  $\mathbb{P}(A \mid E) \geq 0$  for any event  $A$ .
- ▶  $\mathbb{P}(\emptyset \mid E) = 0$ ,  $\mathbb{P}(S \mid E) = 1$ .
- ▶ If  $A_1, \dots, A_n$  are disjoint, then  $\mathbb{P}(\cup_i A_i \mid E) = \sum_i \mathbb{P}(A_i \mid E)$ .

Conclusion:

- ▶ Conditional probabilities are probabilities.
- ▶ *All probabilities are conditional probabilities.*

# Independence

## Definition

Events  $A$  and  $B$  are *independent* if  $\mathbb{P}(A \cap B) = \mathbb{P}(A)\mathbb{P}(B)$ .

# Independence

## Definition

Events  $A$  and  $B$  are *independent* if  $\mathbb{P}(A \cap B) = \mathbb{P}(A)\mathbb{P}(B)$ .

- ▶ If  $\mathbb{P}(A) > 0$  and  $\mathbb{P}(B) > 0$ , this is equivalent to

$$\mathbb{P}(A \mid B) = \mathbb{P}(A) \quad \text{and} \quad \mathbb{P}(B \mid A) = \mathbb{P}(B).$$

# Independence

## Definition

Events  $A$  and  $B$  are *independent* if  $\mathbb{P}(A \cap B) = \mathbb{P}(A)\mathbb{P}(B)$ .

- ▶ If  $\mathbb{P}(A) > 0$  and  $\mathbb{P}(B) > 0$ , this is equivalent to

$$\mathbb{P}(A \mid B) = \mathbb{P}(A) \quad \text{and} \quad \mathbb{P}(B \mid A) = \mathbb{P}(B).$$

- ▶ Independence is a symmetric relation: if  $A$  is independent of  $B$ ,  $B$  is independent of  $A$ .

# Independence

## Definition

Events  $A$  and  $B$  are *independent* if  $\mathbb{P}(A \cap B) = \mathbb{P}(A)\mathbb{P}(B)$ .

- ▶ If  $\mathbb{P}(A) > 0$  and  $\mathbb{P}(B) > 0$ , this is equivalent to

$$\mathbb{P}(A \mid B) = \mathbb{P}(A) \quad \text{and} \quad \mathbb{P}(B \mid A) = \mathbb{P}(B).$$

- ▶ Independence is a symmetric relation: if  $A$  is independent of  $B$ ,  $B$  is independent of  $A$ .
- ▶ Warning: independence  $\neq$  disjointness. In fact, disjoint events can only be independent if  $\mathbb{P}(A) = 0$  or  $\mathbb{P}(B) = 0$ . (Why?)

# Conditional independence

## Definition

Events  $A$  and  $B$  are said to be *conditionally independent* given a third event  $E$  if

$$\mathbb{P}(A \cap B \mid E) = \mathbb{P}(A \mid E)\mathbb{P}(B \mid E).$$



# Conditional independence

## Definition

Events  $A$  and  $B$  are said to be *conditionally independent* given a third event  $E$  if

$$\mathbb{P}(A \cap B \mid E) = \mathbb{P}(A \mid E)\mathbb{P}(B \mid E).$$

**Health warning:** independence does *not* imply conditional independence and vice versa.

# Conditional independence and complements

## Conditional independence and complements

- ▶ Suppose there are two types of teachers: those who give grades that reflect student effort ( $E$ ), and those who randomly assign grades, regardless of student effort ( $E^c$ ).

## Conditional independence and complements

- ▶ Suppose there are two types of teachers: those who give grades that reflect student effort ( $E$ ), and those who randomly assign grades, regardless of student effort ( $E^c$ ).
- ▶ Let  $W$  be the event that you work hard and let  $G$  be the event that you receive a good grade.

## Conditional independence and complements

- ▶ Suppose there are two types of teachers: those who give grades that reflect student effort ( $E$ ), and those who randomly assign grades, regardless of student effort ( $E^c$ ).
- ▶ Let  $W$  be the event that you work hard and let  $G$  be the event that you receive a good grade.
- ▶ Then  $W$  and  $G$  are conditionally independent given  $E^c$ , but they are not conditionally independent given  $E$ .

Conditional independence  $\not\Rightarrow$  independence

## Conditional independence $\nRightarrow$ independence

- ▶ You have one fair coin and one biased coin which lands Heads with probability  $3/4$ .

## Conditional independence $\nRightarrow$ independence

- ▶ You have one fair coin and one biased coin which lands Heads with probability  $3/4$ .
- ▶ You pick one of the coins at random, without knowing which one you've chosen, and flip it several times.



## Conditional independence $\nRightarrow$ independence

- ▶ You have one fair coin and one biased coin which lands Heads with probability  $3/4$ .
- ▶ You pick one of the coins at random, without knowing which one you've chosen, and flip it several times.
- ▶ Conditional on choosing either the fair or the biased coin, the coin flips are independent.

## Conditional independence $\nRightarrow$ independence

- ▶ You have one fair coin and one biased coin which lands Heads with probability  $3/4$ .
- ▶ You pick one of the coins at random, without knowing which one you've chosen, and flip it several times.
- ▶ Conditional on choosing either the fair or the biased coin, the coin flips are independent.
- ▶ However, the coin flips are not unconditionally independent: without knowing which coin we've chosen, each flip gives us new data from which we can predict outcomes of future tosses.

## Conditional independence $\nRightarrow$ independence

- ▶ You have one fair coin and one biased coin which lands Heads with probability  $3/4$ .
- ▶ You pick one of the coins at random, without knowing which one you've chosen, and flip it several times.
- ▶ Conditional on choosing either the fair or the biased coin, the coin flips are independent.
- ▶ However, the coin flips are not unconditionally independent: without knowing which coin we've chosen, each flip gives us new data from which we can predict outcomes of future tosses.
- ▶ (Think about the definition of independence.)

Independence  $\nRightarrow$  conditional independence

## Independence $\not\Rightarrow$ conditional independence

- ▶ Suppose my friends Alice and Ben are the only two people who call me on my mobile phone.

## Independence $\nRightarrow$ conditional independence

- ▶ Suppose my friends Alice and Ben are the only two people who call me on my mobile phone.
- ▶ Each day, they decide independently whether to call me that day.

## Independence $\nRightarrow$ conditional independence

- ▶ Suppose my friends Alice and Ben are the only two people who call me on my mobile phone.
- ▶ Each day, they decide independently whether to call me that day.
- ▶ Let  $A$  be the event that Alice calls me tomorrow and let  $B$  be the event that Ben calls me tomorrow.

## Independence $\nRightarrow$ conditional independence

- ▶ Suppose my friends Alice and Ben are the only two people who call me on my mobile phone.
- ▶ Each day, they decide independently whether to call me that day.
- ▶ Let  $A$  be the event that Alice calls me tomorrow and let  $B$  be the event that Ben calls me tomorrow.
- ▶ Then  $A$  and  $B$  are unconditionally independent, with  $\mathbb{P}(A) > 0$  and  $\mathbb{P}(B) > 0$ .



## Independence $\nRightarrow$ conditional independence

- ▶ Suppose my friends Alice and Ben are the only two people who call me on my mobile phone.
- ▶ Each day, they decide independently whether to call me that day.
- ▶ Let  $A$  be the event that Alice calls me tomorrow and let  $B$  be the event that Ben calls me tomorrow.
- ▶ Then  $A$  and  $B$  are unconditionally independent, with  $\mathbb{P}(A) > 0$  and  $\mathbb{P}(B) > 0$ .
- ▶ However, given that I receive exactly one call tomorrow ( $C$ ),  $A$  and  $B$  are no longer independent:

$$\mathbb{P}(A \mid C) > 0, \quad \text{but} \quad \mathbb{P}(A \mid C \cap B) = 0.$$

# The Monty Hall problem

# The Monty Hall problem

- ▶ Choose one of three doors: two with goats, one with car.

# The Monty Hall problem

- ▶ Choose one of three doors: two with goats, one with car.
- ▶ After your choice, Monty Hall opens one door with a goat and asks if you want to switch to the remaining door. Should you switch?

# The Monty Hall problem

- ▶ Choose one of three doors: two with goats, one with car.
- ▶ After your choice, Monty Hall opens one door with a goat and asks if you want to switch to the remaining door. Should you switch?
- ▶ Solve analytically or by simulation.

## The Monty Hall problem: solving by simulation

```
sims <- 10 ^ 5 # Simulations  
doors <- 1:3 # Label doors  
win_no_switch <- 0 # Win count without switch  
win_switch <- 0 # Win count with switch
```

# The Monty Hall problem: solving by simulation

```
sims <- 10 ^ 5 # Simulations
doors <- 1:3 # Label doors
win_no_switch <- 0 # Win count without switch
win_switch <- 0 # Win count with switch
```

```
for (i in 1:sims) {
  # Pick a door for the car
  car_door <- sample(doors, 1)
  # Pick a door for the contestant
  chosen_door <- sample(doors, 1)
  # If they match, add to "no switch" count
  if (car_door == chosen_door)
    win_no_switch <- win_no_switch + 1
  # Otherwise, add to switch count
  else
    win_switch <- win_switch + 1
}
```

## The Monty Hall problem: solving by simulation (cont.)

```
cat("P(car | no switch) =", win_no_switch / sims)  
## P(car | no switch) = 0.33504
```



## The Monty Hall problem: solving by simulation (cont.)

```
cat("P(car | no switch) =", win_no_switch / sims)  
## P(car | no switch) = 0.33504
```

```
cat("P(car | switch) =", win_switch / sims)  
## P(car | switch) = 0.66496
```

How is this possible?

# The Monty Hall problem: analytic solution

## The Monty Hall problem: analytic solution

- ▶ Conditioning as wishful thinking: condition on what you wish you knew.

## The Monty Hall problem: analytic solution

- ▶ Conditioning as wishful thinking: condition on what you wish you knew.
- ▶ Let  $C_i$  be the event that the car is behind door  $i$ , for  $1 \leq i \leq 3$ , and let  $W$  be the event that you have chosen the winning door.

## The Monty Hall problem: analytic solution

- ▶ Conditioning as wishful thinking: condition on what you wish you knew.
- ▶ Let  $C_i$  be the event that the car is behind door  $i$ , for  $1 \leq i \leq 3$ , and let  $W$  be the event that you have chosen the winning door.
- ▶ Since the  $C_i$  partition the sample space, we can use the Law of Total Probability:

$$\mathbb{P}(W) = \mathbb{P}(W \mid C_1) \times \frac{1}{3} + \mathbb{P}(W \mid C_2) \times \frac{1}{3} + \mathbb{P}(W \mid C_3) \times \frac{1}{3}.$$

## The Monty Hall problem: analytic solution

- ▶ Conditioning as wishful thinking: condition on what you wish you knew.
- ▶ Let  $C_i$  be the event that the car is behind door  $i$ , for  $1 \leq i \leq 3$ , and let  $W$  be the event that you have chosen the winning door.
- ▶ Since the  $C_i$  partition the sample space, we can use the Law of Total Probability:

$$\mathbb{P}(W) = \mathbb{P}(W \mid C_1) \times \frac{1}{3} + \mathbb{P}(W \mid C_2) \times \frac{1}{3} + \mathbb{P}(W \mid C_3) \times \frac{1}{3}.$$

- ▶ Suppose, without loss of generality, that you chose door 1.

## The Monty Hall problem: analytic solution

- ▶ Conditioning as wishful thinking: condition on what you wish you knew.
- ▶ Let  $C_i$  be the event that the car is behind door  $i$ , for  $1 \leq i \leq 3$ , and let  $W$  be the event that you have chosen the winning door.
- ▶ Since the  $C_i$  partition the sample space, we can use the Law of Total Probability:

$$\mathbb{P}(W) = \mathbb{P}(W \mid C_1) \times \frac{1}{3} + \mathbb{P}(W \mid C_2) \times \frac{1}{3} + \mathbb{P}(W \mid C_3) \times \frac{1}{3}.$$

- ▶ Suppose, without loss of generality, that you chose door 1.
- ▶ If  $C_1$ , the switching strategy will fail:

$$\mathbb{P}(W \mid C_1) = 0.$$

## The Monty Hall problem: analytic solution

- ▶ Conditioning as wishful thinking: condition on what you wish you knew.
- ▶ Let  $C_i$  be the event that the car is behind door  $i$ , for  $1 \leq i \leq 3$ , and let  $W$  be the event that you have chosen the winning door.
- ▶ Since the  $C_i$  partition the sample space, we can use the Law of Total Probability:

$$\mathbb{P}(W) = \mathbb{P}(W \mid C_1) \times \frac{1}{3} + \mathbb{P}(W \mid C_2) \times \frac{1}{3} + \mathbb{P}(W \mid C_3) \times \frac{1}{3}.$$

- ▶ Suppose, without loss of generality, that you chose door 1.
- ▶ If  $C_1$ , the switching strategy will fail:

$$\mathbb{P}(W \mid C_1) = 0.$$

- ▶ If the car is behind doors 2 or 3, switching will succeed:

$$\mathbb{P}(W \mid C_2) = \mathbb{P}(W \mid C_3) = 1.$$



## The Monty Hall problem: analytic solution

- ▶ Conditioning as wishful thinking: condition on what you wish you knew.
- ▶ Let  $C_i$  be the event that the car is behind door  $i$ , for  $1 \leq i \leq 3$ , and let  $W$  be the event that you have chosen the winning door.
- ▶ Since the  $C_i$  partition the sample space, we can use the Law of Total Probability:

$$\mathbb{P}(W) = \mathbb{P}(W \mid C_1) \times \frac{1}{3} + \mathbb{P}(W \mid C_2) \times \frac{1}{3} + \mathbb{P}(W \mid C_3) \times \frac{1}{3}.$$

- ▶ Suppose, without loss of generality, that you chose door 1.
- ▶ If  $C_1$ , the switching strategy will fail:

$$\mathbb{P}(W \mid C_1) = 0.$$

- ▶ If the car is behind doors 2 or 3, switching will succeed:

$$\mathbb{P}(W \mid C_2) = \mathbb{P}(W \mid C_3) = 1.$$

- ▶ Hence,  $\mathbb{P}(W) = 0 \times \frac{1}{3} + 1 \times \frac{1}{3} + 1 \times \frac{1}{3} = \frac{2}{3}$ .

# Problem sheet 1: preamble

```
# Clear environment, set working directory  
rm(list = ls())  
setwd("/Users/Elias/Documents/.../QM/Problem sheets")
```

```
# Load tidyverse packages  
library(tidyverse)
```

## Question A1

```
# Construct die and sample space  
die <- 1:6  
S <- expand.grid(die, die, die)
```

## Question A2

```
# Create new variable  
S <- S %>%  
  mutate(Value = Var1 + Var2 + Var3)
```

## Question A3

```
#  $P(A)$  = # elements in  $A$  / # elements in  $S$   
sum(S$Value == 12) / nrow(S)  
  
## [1] 0.1157407
```

## Question A4

```
# What if dice are biased?
```

```
Prob <- c("1" = 1/8, "2" = 1/8, "3" = 1/8,  
          "4" = 1/8, "5" = 1/8, "6" = 3/8)
```

## Question A4

```
# What if dice are biased?
```

```
Prob <- c("1" = 1/8, "2" = 1/8, "3" = 1/8,  
          "4" = 1/8, "5" = 1/8, "6" = 3/8)
```

```
# Assign individual and joint probabilities
```

```
S$Prob1 <- Prob[S$Var1]  
S$Prob2 <- Prob[S$Var2]  
S$Prob3 <- Prob[S$Var3]  
S$Prob_joint <- S$Prob1 * S$Prob2 * S$Prob3
```

## Question A5

```
# Extract event A and calculate probability  
A <- subset(S, Value == 12)  
sum(A$Prob_joint)  
  
## [1] 0.1074219
```



## Question B1

```
# Load data sets  
kenya <- read_csv("kenya.csv")  
sweden <- read_csv("sweden.csv")  
world <- read_csv("world.csv")
```

## Question B2

```
# Inspect
summary(kenya)
summary(sweden)
summary(world)

glimpse(kenya)
glimpse(sweden)
glimpse(world)

head(kenya)
head(sweden)
head(world)

print(kenya, n = 30)
print(sweden, n = 30)
print(world, n = 30)
```

## Question B3

```
# Calculate age-specific fertility rate
asfr <- function(data) {
  data %>%
    mutate(
      asfr = births / py.women) %>%
    select(period, age, asfr) %>%
    data.frame() # Convert tibble to data frame
}
```

## Question B3 (cont.)

```
asfr(kenya)[c(4:10, 19:25), ]
```

##	period	age	asfr
## 4	1950-1955	15-19	0.16884585
## 5	1950-1955	20-24	0.35596942
## 6	1950-1955	25-29	0.34657814
## 7	1950-1955	30-34	0.28946367
## 8	1950-1955	35-39	0.20644016
## 9	1950-1955	40-44	0.11193267
## 10	1950-1955	45-49	0.03905205
## 19	2005-2010	15-19	0.10057087
## 20	2005-2010	20-24	0.23583536
## 21	2005-2010	25-29	0.23294721
## 22	2005-2010	30-34	0.18087964
## 23	2005-2010	35-39	0.13126805
## 24	2005-2010	40-44	0.05626214
## 25	2005-2010	45-49	0.03815044

## Question B3 (cont.)

```
asfr(sweden)[c(4:10, 19:25), ]
```

##	period	age	asfr
## 4	1950-1955	15-19	0.0389089519
## 5	1950-1955	20-24	0.1277108826
## 6	1950-1955	25-29	0.1252436647
## 7	1950-1955	30-34	0.0873641591
## 8	1950-1955	35-39	0.0486037714
## 9	1950-1955	40-44	0.0162101857
## 10	1950-1955	45-49	0.0013418290
## 19	2005-2010	15-19	0.0059709097
## 20	2005-2010	20-24	0.0507320271
## 21	2005-2010	25-29	0.1162085625
## 22	2005-2010	30-34	0.1322744621
## 23	2005-2010	35-39	0.0625923991
## 24	2005-2010	40-44	0.0121600765
## 25	2005-2010	45-49	0.0006143942

## Question B3 (cont.)

```
asfr(world)[c(4:10, 19:25), ]
```

##		period	age	asfr
## 4	1950-1955	15-19	0.090295213	
## 5	1950-1955	20-24	0.237633702	
## 6	1950-1955	25-29	0.252452289	
## 7	1950-1955	30-34	0.204164096	
## 8	1950-1955	35-39	0.138105344	
## 9	1950-1955	40-44	0.063608319	
## 10	1950-1955	45-49	0.015190644	
## 19	2005-2010	15-19	0.048489719	
## 20	2005-2010	20-24	0.151971307	
## 21	2005-2010	25-29	0.146980966	
## 22	2005-2010	30-34	0.093813813	
## 23	2005-2010	35-39	0.046689639	
## 24	2005-2010	40-44	0.016268995	
## 25	2005-2010	45-49	0.004510245	

## Question B4

```
# Calculate total fertility rate
tfr <- function(data) {
  out <- asfr(data)
  out %>%
    group_by(period) %>%
    summarise(
      tfr = 5 * sum(asfr))
}
```

## Question B4 (cont.)

```
tfr(kenya)
```

```
## # A tibble: 2 x 2
##   period      tfr
##   <chr>      <dbl>
## 1 1950-1955  7.59
## 2 2005-2010  4.88
```

```
tfr(sweden)
```

```
## # A tibble: 2 x 2
##   period      tfr
##   <chr>      <dbl>
## 1 1950-1955  2.23
## 2 2005-2010  1.90
```



## Question B4 (cont.)

```
tfr(world)
```

```
## # A tibble: 2 x 2
```

```
##   period      tfr
```

```
##   <chr>      <dbl>
```

```
## 1 1950-1955  5.01
```

```
## 2 2005-2010  2.54
```

## Question B5

```
# Calculate age-specific death rate
asdr <- function(data) {
  data %>%
    mutate(
      # Convert rates to per 1000 population
      asdr = 1000 * deaths / (py.men + py.women)) %>%
    select(period, age, asdr) %>%
    data.frame() # Convert tibble to data frame
}
```

## Question B5 (cont.)

```
sample_n(asdr(kenya), 10)
```

##		period	age	asdr
## 1		1950-1955	15-19	5.869582
## 2		2005-2010	60-69	25.395531
## 3		1950-1955	10-14	5.972093
## 4		1950-1955	55-59	24.433007
## 5		2005-2010	30-34	10.603913
## 6		1950-1955	60-69	41.996801
## 7		1950-1955	20-24	7.651103
## 8		1950-1955	40-44	12.633744
## 9		1950-1955	0-4	66.826532
## 10		2005-2010	20-24	3.885368

## Question B5 (cont.)

```
sample_n(asdr(sweden), 10)
```

##		period	age	asdr
## 1		1950-1955	80+	167.81702554
## 2		2005-2010	5-9	0.08138094
## 3		2005-2010	40-44	1.03925622
## 4		2005-2010	25-29	0.49414399
## 5		2005-2010	60-69	9.82877193
## 6		1950-1955	10-14	0.48964064
## 7		1950-1955	25-29	1.11409103
## 8		1950-1955	40-44	2.50955411
## 9		1950-1955	70-79	59.98230926
## 10		1950-1955	5-9	0.43205374

## Question B5 (cont.)

```
sample_n(asdr(world), 10)
```

##		period	age	asdr
## 1		2005-2010	30-34	2.623982
## 2		1950-1955	80+	184.364978
## 3		1950-1955	45-49	13.459846
## 4		1950-1955	30-34	7.132501
## 5		2005-2010	80+	120.679385
## 6		2005-2010	70-79	47.457519
## 7		1950-1955	55-59	24.265320
## 8		2005-2010	20-24	1.832602
## 9		2005-2010	35-39	3.031563
## 10		2005-2010	40-44	3.753402

## Question B6

```
# Collect ASFR and ASDR for each country  
ken <- left_join(asfr(kenya), asdr(kenya))  
swe <- left_join(asfr(sweden), asdr(sweden))  
wor <- left_join(asfr(world), asdr(world))
```

## Question B6

```
# Collect ASFR and ASDR for each country  
ken <- left_join(asfr(kenya), asdr(kenya))  
swe <- left_join(asfr(sweden), asdr(sweden))  
wor <- left_join(asfr(world), asdr(world))
```

```
# Create one data frame with all results  
df <- rbind(ken, swe, wor)  
df$country <- c(rep("Kenya", 30),  
                rep("Sweden", 30), rep("World", 30))
```

## Question B6 (cont.)

```
# Transform age groups to ordered factor
df$age <- factor(df$age,
                 levels = c("0-4", "5-9", "10-14",
                             "15-19", "20-24", "25-29",
                             "30-34", "35-39", "40-44",
                             "45-49", "50-54", "55-59",
                             "60-69", "70-79", "80+"))


# Age groups for reproductive age range
age_groups <- c("15-19", "20-24", "25-29",
                "30-34", "35-39", "40-44", "45-49")
```



## Question B6 (cont.)

```
# Visualise ASFR
g1 <- ggplot(subset(df, age %in% age_groups), # Age range
             aes(age, 1000 * asfr, # Modify rate
                 fill = country)) + # Colour code
  geom_col() + # Show as columns
  labs( # Axis labels
    x = "Age",
    y = "Age-specific fertility rate per 1000 women") +
  scale_fill_viridis_d() + # Choose a nice colour palette
  facet_grid(country ~ period) + # Stratify
  theme_bw() + # Remove redundant lines
  theme( # Avoid cluttering
    legend.position = "none",
    axis.text.x = element_text(angle = 90))
```

Age-specific fertility rate per 1000 women



Age

## Question B6 (cont.)

```
# Visualise ASDR
g2 <- ggplot(df, aes(age, asdr,
                      fill = country)) + # Colour code
  geom_col() + # Show as columns
  scale_y_continuous(breaks = seq(0, 200, 50)) + # Y-axis
  labs( # Axis labels
    x = "Age",
    y = "Age-specific death rate per 1000 population") +
  scale_fill_viridis_d(option = "plasma") + # Colour
  facet_grid(country ~ period) + # Stratify
  theme_bw() + # Remove redundant lines
  theme( # Avoid cluttering
    legend.position = "none",
    axis.text.x = element_text(angle = 90))
```

