

# Quantitative Methods

## Human Sciences, 2020–21

Elias Nosrati

Lecture 6: 19 November 2020

Today

# Today

- ▶ Statistical models and the problem of inference.

# Today

- ▶ Statistical models and the problem of inference.
- ▶ Introduction to statistical inference.

# Today

- ▶ Statistical models and the problem of inference.
- ▶ Introduction to statistical inference.
- ▶ Introduction to linear regression models.

# Statistical models

# Statistical models

- ▶ A model is an abstraction of reality.

# Statistical models

- ▶ A model is an abstraction of reality.
- ▶ Judged by its usefulness, in the context of a particular purpose.



# Statistical models

- ▶ A model is an abstraction of reality.
- ▶ Judged by its usefulness, in the context of a particular purpose.
- ▶ Model might not be “true” or entirely realistic, but does it provide us with useful information about some quantity of interest?

## Data generating process

You are given a data set called `lifexp`:

```
head(lifexp)
```

```
## [1] 69 74 78 73 70 75
```

```
tail(lifexp)
```

```
## [1] 73 65 68 69 80 72
```

```
summary(lifexp)
```

##	Min.	1st Qu.	Median	Mean	3rd Qu.	Max.
##	59.00	69.00	72.00	71.93	74.00	84.00

# Data generating process

You are given a data set called `lifexp`:

```
head(lifexp)
```

```
## [1] 69 74 78 73 70 75
```

```
tail(lifexp)
```

```
## [1] 73 65 68 69 80 72
```

```
summary(lifexp)
```

```
##      Min. 1st Qu.  Median    Mean 3rd Qu.    Max.
##    59.00   69.00   72.00   71.93   74.00   84.00
```

What is the underlying data generating process? What are the key features of the underlying distribution?

# Constructing a model

## Constructing a model

- ▶ Any model has to make some assumptions.

# Constructing a model

- ▶ Any model has to make some assumptions.
- ▶ In the lifexp example, assume Normal distribution:

$$Y \sim \mathcal{N}(\mu, \sigma^2).$$

# Constructing a model

- ▶ Any model has to make some assumptions.
- ▶ In the lifexp example, assume Normal distribution:

$$Y \sim \mathcal{N}(\mu, \sigma^2).$$

- ▶ Quantity of interest:

## Constructing a model

- ▶ Any model has to make some assumptions.
- ▶ In the lifexp example, assume Normal distribution:

$$Y \sim \mathcal{N}(\mu, \sigma^2).$$

- ▶ Quantity of interest:  $\mu = \mathbb{E}(Y)$ .



# Constructing a model

- ▶ Any model has to make some assumptions.
- ▶ In the lifexp example, assume Normal distribution:

$$Y \sim \mathcal{N}(\mu, \sigma^2).$$

- ▶ Quantity of interest:  $\mu = \mathbb{E}(Y)$ .
- ▶ Decompose the model into two parts:  $M = \{A, \mu\}$ .

# Constructing a model

- ▶ Any model has to make some assumptions.
- ▶ In the lifexp example, assume Normal distribution:

$$Y \sim \mathcal{N}(\mu, \sigma^2).$$

- ▶ Quantity of interest:  $\mu = \mathbb{E}(Y)$ .
- ▶ Decompose the model into two parts:  $M = \{A, \mu\}$ .
- ▶  $A$  is the *assumed* part (the data generating process is Normal) and  $\mu$  is to be *estimated* using the data we have.

## Constructing a model

- ▶ Any model has to make some assumptions.
- ▶ In the lifexp example, assume Normal distribution:

$$Y \sim \mathcal{N}(\mu, \sigma^2).$$

- ▶ Quantity of interest:  $\mu = \mathbb{E}(Y)$ .
- ▶ Decompose the model into two parts:  $M = \{A, \mu\}$ .
- ▶  $A$  is the *assumed* part (the data generating process is Normal) and  $\mu$  is to be *estimated* using the data we have.
- ▶ But how do we estimate  $\mu$ ?

# Inverse probability and the problem of inference

# Inverse probability and the problem of inference

- ▶ *Inference* is the process of using facts we have (the data) to learn about facts we do not have (the underlying data generating process): involves uncertainty.

# Inverse probability and the problem of inference

- ▶ *Inference* is the process of using facts we have (the data) to learn about facts we do not have (the underlying data generating process): involves uncertainty.
- ▶ Probability as a model of uncertainty.

# Inverse probability and the problem of inference

- ▶ *Inference* is the process of using facts we have (the data) to learn about facts we do not have (the underlying data generating process): involves uncertainty.
- ▶ Probability as a model of uncertainty.
- ▶ Probabilities are traditionally expressed as  $\mathbb{P}(\text{data} \mid \text{model})$ .

# Inverse probability and the problem of inference

- ▶ *Inference* is the process of using facts we have (the data) to learn about facts we do not have (the underlying data generating process): involves uncertainty.
- ▶ Probability as a model of uncertainty.
- ▶ Probabilities are traditionally expressed as  $\mathbb{P}(\text{data} \mid \text{model})$ .
- ▶ For the purposes of inference, we are interested in the *inverse probability*  $\mathbb{P}(\text{model} \mid \text{data})$ .



# Inverse probability and the problem of inference

- ▶ *Inference* is the process of using facts we have (the data) to learn about facts we do not have (the underlying data generating process): involves uncertainty.
- ▶ Probability as a model of uncertainty.
- ▶ Probabilities are traditionally expressed as  $\mathbb{P}(\text{data} \mid \text{model})$ .
- ▶ For the purposes of inference, we are interested in the *inverse probability*  $\mathbb{P}(\text{model} \mid \text{data})$ .
- ▶ Uncertainty lies with the model (the process that generates the data) rather than the (already observed) data.

# Inverse probability and the problem of inference

- ▶ *Inference* is the process of using facts we have (the data) to learn about facts we do not have (the underlying data generating process): involves uncertainty.
- ▶ Probability as a model of uncertainty.
- ▶ Probabilities are traditionally expressed as  $\mathbb{P}(\text{data} \mid \text{model})$ .
- ▶ For the purposes of inference, we are interested in the *inverse probability*  $\mathbb{P}(\text{model} \mid \text{data})$ .
- ▶ Uncertainty lies with the model (the process that generates the data) rather than the (already observed) data.
- ▶ How to go from  $\mathbb{P}(\text{data} \mid \text{model})$  to  $\mathbb{P}(\text{model} \mid \text{data})$ ?

# Inverse probability and the problem of inference

- ▶ *Inference* is the process of using facts we have (the data) to learn about facts we do not have (the underlying data generating process): involves uncertainty.
- ▶ Probability as a model of uncertainty.
- ▶ Probabilities are traditionally expressed as  $\mathbb{P}(\text{data} \mid \text{model})$ .
- ▶ For the purposes of inference, we are interested in the *inverse probability*  $\mathbb{P}(\text{model} \mid \text{data})$ .
- ▶ Uncertainty lies with the model (the process that generates the data) rather than the (already observed) data.
- ▶ How to go from  $\mathbb{P}(\text{data} \mid \text{model})$  to  $\mathbb{P}(\text{model} \mid \text{data})$ ?
- ▶ How to go from  $\mathbb{P}(\text{known} \mid \text{unknown})$  to  $\mathbb{P}(\text{unknown} \mid \text{known})$ ?

# Inverse probability and the problem of inference

- ▶ *Inference* is the process of using facts we have (the data) to learn about facts we do not have (the underlying data generating process): involves uncertainty.
- ▶ Probability as a model of uncertainty.
- ▶ Probabilities are traditionally expressed as  $\mathbb{P}(\text{data} \mid \text{model})$ .
- ▶ For the purposes of inference, we are interested in the *inverse probability*  $\mathbb{P}(\text{model} \mid \text{data})$ .
- ▶ Uncertainty lies with the model (the process that generates the data) rather than the (already observed) data.
- ▶ How to go from  $\mathbb{P}(\text{data} \mid \text{model})$  to  $\mathbb{P}(\text{model} \mid \text{data})$ ?
- ▶ How to go from  $\mathbb{P}(\text{known} \mid \text{unknown})$  to  $\mathbb{P}(\text{unknown} \mid \text{known})$ ?
- ▶ How to go from  $\mathbb{P}(\text{lifexp} \mid \mu)$  to  $\mathbb{P}(\mu \mid \text{lifexp})$ ?

# Building a theory of inference

## Building a theory of inference

- ▶ Let  $y$  be the realisation of  $Y$ .

## Building a theory of inference

- ▶ Let  $y$  be the realisation of  $Y$ .
- ▶ Let  $\theta$  be any parameter of interest.

## Building a theory of inference

- ▶ Let  $y$  be the realisation of  $Y$ .
- ▶ Let  $\theta$  be any parameter of interest.
- ▶ Inverse probability — use Bayes' Rule:



## Building a theory of inference

- ▶ Let  $y$  be the realisation of  $Y$ .
- ▶ Let  $\theta$  be any parameter of interest.
- ▶ Inverse probability — use Bayes' Rule:

$$\mathbb{P}(\theta \mid y) = \frac{\mathbb{P}(\theta, y)}{\mathbb{P}(y)}$$

## Building a theory of inference

- ▶ Let  $y$  be the realisation of  $Y$ .
- ▶ Let  $\theta$  be any parameter of interest.
- ▶ Inverse probability — use Bayes' Rule:

$$\mathbb{P}(\theta \mid y) = \frac{\mathbb{P}(\theta, y)}{\mathbb{P}(y)} = \frac{\mathbb{P}(\theta)\mathbb{P}(y \mid \theta)}{\mathbb{P}(y)}.$$

## Building a theory of inference

- ▶ Let  $y$  be the realisation of  $Y$ .
- ▶ Let  $\theta$  be any parameter of interest.
- ▶ Inverse probability — use Bayes' Rule:

$$\mathbb{P}(\theta \mid y) = \frac{\mathbb{P}(\theta, y)}{\mathbb{P}(y)} = \frac{\mathbb{P}(\theta)\mathbb{P}(y \mid \theta)}{\mathbb{P}(y)}.$$

- ▶ Let  $\mathbb{P}(\theta)/\mathbb{P}(y)$  be an unknown constant function  $k(y)$ .

## Building a theory of inference

- ▶ Let  $y$  be the realisation of  $Y$ .
- ▶ Let  $\theta$  be any parameter of interest.
- ▶ Inverse probability — use Bayes' Rule:

$$\mathbb{P}(\theta \mid y) = \frac{\mathbb{P}(\theta, y)}{\mathbb{P}(y)} = \frac{\mathbb{P}(\theta)\mathbb{P}(y \mid \theta)}{\mathbb{P}(y)}.$$

- ▶ Let  $\mathbb{P}(\theta)/\mathbb{P}(y)$  be an unknown constant function  $k(y)$ .
- ▶ Define the *likelihood* function:

$$L(\theta \mid y) = k(y)\mathbb{P}(y \mid \theta).$$

## Building a theory of inference

- ▶ Let  $y$  be the realisation of  $Y$ .
- ▶ Let  $\theta$  be any parameter of interest.
- ▶ Inverse probability — use Bayes' Rule:

$$\mathbb{P}(\theta \mid y) = \frac{\mathbb{P}(\theta, y)}{\mathbb{P}(y)} = \frac{\mathbb{P}(\theta)\mathbb{P}(y \mid \theta)}{\mathbb{P}(y)}.$$

- ▶ Let  $\mathbb{P}(\theta)/\mathbb{P}(y)$  be an unknown constant function  $k(y)$ .
- ▶ Define the *likelihood* function:

$$L(\theta \mid y) = k(y)\mathbb{P}(y \mid \theta).$$

- ▶ The numerical value of  $L$  is interpretable as the *likelihood* of a model having generated the observed data.

## Building a theory of inference

- ▶ Let  $y$  be the realisation of  $Y$ .
- ▶ Let  $\theta$  be any parameter of interest.
- ▶ Inverse probability — use Bayes' Rule:

$$\mathbb{P}(\theta | y) = \frac{\mathbb{P}(\theta, y)}{\mathbb{P}(y)} = \frac{\mathbb{P}(\theta)\mathbb{P}(y | \theta)}{\mathbb{P}(y)}.$$

- ▶ Let  $\mathbb{P}(\theta)/\mathbb{P}(y)$  be an unknown constant function  $k(y)$ .
- ▶ Define the *likelihood* function:

$$L(\theta | y) = k(y)\mathbb{P}(y | \theta).$$

- ▶ The numerical value of  $L$  is interpretable as the *likelihood* of a model having generated the observed data.
- ▶ This numerical value is not equal to, but is *proportional* to a probability density:

$$L(\theta | y) \propto \mathbb{P}(y | \theta).$$

## Building a theory of inference

- ▶ Let  $y$  be the realisation of  $Y$ .
- ▶ Let  $\theta$  be any parameter of interest.
- ▶ Inverse probability — use Bayes' Rule:

$$\mathbb{P}(\theta | y) = \frac{\mathbb{P}(\theta, y)}{\mathbb{P}(y)} = \frac{\mathbb{P}(\theta)\mathbb{P}(y | \theta)}{\mathbb{P}(y)}.$$

- ▶ Let  $\mathbb{P}(\theta)/\mathbb{P}(y)$  be an unknown constant function  $k(y)$ .
- ▶ Define the *likelihood* function:

$$L(\theta | y) = k(y)\mathbb{P}(y | \theta).$$

- ▶ The numerical value of  $L$  is interpretable as the *likelihood* of a model having generated the observed data.
- ▶ This numerical value is not equal to, but is *proportional* to a probability density:

$$L(\theta | y) \propto \mathbb{P}(y | \theta).$$

- ▶ A *relative* (not absolute) measure of uncertainty.

# Likelihood theory of inference



## Likelihood theory of inference



- ▶ A normal probability density  $f(y \mid \theta)$  holds  $\theta$  fixed and is a function of the data, defined on the *data scale*.

# Likelihood theory of inference

- ▶ A normal probability density  $f(y \mid \theta)$  holds  $\theta$  fixed and is a function of the data, defined on the *data scale*.
- ▶ The likelihood function holds the data fixed and is a function of  $\theta$ , defined on the *parameter scale*.

# Likelihood theory of inference

- ▶ A normal probability density  $f(y \mid \theta)$  holds  $\theta$  fixed and is a function of the data, defined on the *data scale*.
- ▶ The likelihood function holds the data fixed and is a function of  $\theta$ , defined on the *parameter scale*.



## Likelihood theory of inference (cont.)

## Likelihood theory of inference (cont.)

- ▶ For mathematical and computational reason, it is common to use the *log*-likelihood function.

## Likelihood theory of inference (cont.)

- ▶ For mathematical and computational reason, it is common to use the *log*-likelihood function.
- ▶ Key question: given what we observe, which model is most likely to have generated our data?

## Likelihood theory of inference (cont.)

- ▶ For mathematical and computational reason, it is common to use the *log*-likelihood function.
- ▶ Key question: given what we observe, which model is most likely to have generated our data?
- ▶ The most likely value of  $\theta$  is where  $L$  is at its maximum.

## Likelihood theory of inference (cont.)

- ▶ For mathematical and computational reason, it is common to use the *log*-likelihood function.
- ▶ Key question: given what we observe, which model is most likely to have generated our data?
- ▶ The most likely value of  $\theta$  is where  $L$  is at its maximum.
- ▶ Maximum likelihood estimation: *maximise* the likelihood.




## Likelihood theory of inference (cont.)

- ▶ For mathematical and computational reason, it is common to use the *log*-likelihood function.
- ▶ Key question: given what we observe, which model is most likely to have generated our data?
- ▶ The most likely value of  $\theta$  is where  $L$  is at its maximum.
- ▶ Maximum likelihood estimation: *maximise* the likelihood.
- ▶ The result: a maximum likelihood estimate (MLE)  $\hat{\theta}$ .

## Likelihood theory of inference (cont.)

- ▶ For mathematical and computational reason, it is common to use the *log*-likelihood function.
- ▶ Key question: given what we observe, which model is most likely to have generated our data?
- ▶ The most likely value of  $\theta$  is where  $L$  is at its maximum.
- ▶ Maximum likelihood estimation: *maximise* the likelihood.
- ▶ The result: a maximum likelihood estimate (MLE)  $\hat{\theta}$ .



# Bayesian theory of inference

# Bayesian theory of inference

- Back to the inverse probability:

$$\mathbb{P}(\theta \mid y) \propto \mathbb{P}(\theta) \mathbb{P}(y \mid \theta).$$

# Bayesian theory of inference

- ▶ Back to the inverse probability:

$$\mathbb{P}(\theta \mid y) \propto \mathbb{P}(\theta) \mathbb{P}(y \mid \theta).$$

- ▶  $\mathbb{P}(\theta \mid y)$ : posterior density.

# Bayesian theory of inference

- ▶ Back to the inverse probability:

$$\mathbb{P}(\theta \mid y) \propto \mathbb{P}(\theta) \mathbb{P}(y \mid \theta).$$

- ▶  $\mathbb{P}(\theta \mid y)$ : posterior density.
- ▶  $\mathbb{P}(y \mid \theta)$ : traditional probability ( $\propto$  likelihood).

# Bayesian theory of inference

- ▶ Back to the inverse probability:

$$\mathbb{P}(\theta \mid y) \propto \mathbb{P}(\theta) \mathbb{P}(y \mid \theta).$$

- ▶  $\mathbb{P}(\theta \mid y)$ : posterior density.
- ▶  $\mathbb{P}(y \mid \theta)$ : traditional probability ( $\propto$  likelihood).
- ▶  $\mathbb{P}(\theta)$ : prior density.

# Bayesian theory of inference

- ▶ Back to the inverse probability:

$$\mathbb{P}(\theta \mid y) \propto \mathbb{P}(\theta) \mathbb{P}(y \mid \theta).$$

- ▶  $\mathbb{P}(\theta \mid y)$ : posterior density.
- ▶  $\mathbb{P}(y \mid \theta)$ : traditional probability ( $\propto$  likelihood).
- ▶  $\mathbb{P}(\theta)$ : prior density.
- ▶ Bayesian inference differs from likelihood inference in its use of *prior* information about  $\theta$ :

$$\text{posterior} \propto \text{prior} \times \text{likelihood}.$$



# Bayesian theory of inference

- ▶ Back to the inverse probability:

$$\mathbb{P}(\theta \mid y) \propto \mathbb{P}(\theta) \mathbb{P}(y \mid \theta).$$

- ▶  $\mathbb{P}(\theta \mid y)$ : posterior density.
- ▶  $\mathbb{P}(y \mid \theta)$ : traditional probability ( $\propto$  likelihood).
- ▶  $\mathbb{P}(\theta)$ : prior density.
- ▶ Bayesian inference differs from likelihood inference in its use of *prior* information about  $\theta$ :

$$\text{posterior} \propto \text{prior} \times \text{likelihood}.$$

- ▶ A way of integrating evidence from outside the data into the model.

# Bayesian theory of inference

- ▶ Back to the inverse probability:

$$\mathbb{P}(\theta \mid y) \propto \mathbb{P}(\theta) \mathbb{P}(y \mid \theta).$$

- ▶  $\mathbb{P}(\theta \mid y)$ : posterior density.
- ▶  $\mathbb{P}(y \mid \theta)$ : traditional probability ( $\propto$  likelihood).
- ▶  $\mathbb{P}(\theta)$ : prior density.
- ▶ Bayesian inference differs from likelihood inference in its use of *prior* information about  $\theta$ :

$$\text{posterior} \propto \text{prior} \times \text{likelihood}.$$

- ▶ A way of integrating evidence from outside the data into the model.
- ▶ Practical difference: minor unless prior is actually important.

## Election example

## Election example

- ▶ Candidate  $A$  and candidate  $B$ .

## Election example

- ▶ Candidate  $A$  and candidate  $B$ .
- ▶ Outcome of experiment is Bernoulli random variable  $Y$  with parameter  $p = \mathbb{P}(A \text{ wins})$ .

## Election example

- ▶ Candidate  $A$  and candidate  $B$ .
- ▶ Outcome of experiment is Bernoulli random variable  $Y$  with parameter  $p = \mathbb{P}(A \text{ wins})$ .
- ▶ We use polling data calculate posterior density  $\mathbb{P}(p \mid y)$ .

## Election example

- ▶ Candidate  $A$  and candidate  $B$ .
- ▶ Outcome of experiment is Bernoulli random variable  $Y$  with parameter  $p = \mathbb{P}(A \text{ wins})$ .
- ▶ We use polling data calculate posterior density  $\mathbb{P}(p \mid y)$ .
- ▶ Likelihood:  $L(p \mid y) \propto \mathbb{P}(y \mid p)$ .

## Election example

- ▶ Candidate  $A$  and candidate  $B$ .
- ▶ Outcome of experiment is Bernoulli random variable  $Y$  with parameter  $p = \mathbb{P}(A \text{ wins})$ .
- ▶ We use polling data calculate posterior density  $\mathbb{P}(p \mid y)$ .
- ▶ Likelihood:  $L(p \mid y) \propto \mathbb{P}(y \mid p)$ .
- ▶ Bayesian: use prior information about  $p$  (e.g., previous poll data or evidence of rigged election).




## Election example

- ▶ Candidate  $A$  and candidate  $B$ .
- ▶ Outcome of experiment is Bernoulli random variable  $Y$  with parameter  $p = \mathbb{P}(A \text{ wins})$ .
- ▶ We use polling data calculate posterior density  $\mathbb{P}(p \mid y)$ .
- ▶ Likelihood:  $L(p \mid y) \propto \mathbb{P}(y \mid p)$ .
- ▶ Bayesian: use prior information about  $p$  (e.g., previous poll data or evidence of rigged election).
- ▶ Challenge: what is the exact value of the prior?


## Election example

- ▶ Candidate  $A$  and candidate  $B$ .
- ▶ Outcome of experiment is Bernoulli random variable  $Y$  with parameter  $p = \mathbb{P}(A \text{ wins})$ .
- ▶ We use polling data calculate posterior density  $\mathbb{P}(p \mid y)$ .
- ▶ Likelihood:  $L(p \mid y) \propto \mathbb{P}(y \mid p)$ .
- ▶ Bayesian: use prior information about  $p$  (e.g., previous poll data or evidence of rigged election).
- ▶ Challenge: what is the exact value of the prior?
- ▶ Typical solution: try a range of possible values.

## A non-theory of inference: ordinary least squares




## A non-theory of inference: ordinary least squares




- ▶ A common approach: minimise (“*least*”) the squared deviations (“*squares*”) of the observed data from the estimated parameter value.

## A non-theory of inference: ordinary least squares



- ▶ A common approach: minimise (“*least*”) the squared deviations (“*squares*”) of the observed data from the estimated parameter value.
- ▶ Classical example: how to fit a line to a cloud of points.

## A non-theory of inference: ordinary least squares



- ▶ A common approach: minimise (“*least*”) the squared deviations (“*squares*”) of the observed data from the estimated parameter value.
- ▶ Classical example: how to fit a line to a cloud of points.
- ▶ The corresponding parameter estimate (slope of line) is known as the *ordinary least squares* (OLS) estimate.

A non-theory of inference: ordinary least squares (cont.)

## A non-theory of inference: ordinary least squares (cont.)

- ▶ Not a theory of inference ...



## A non-theory of inference: ordinary least squares (cont.)

- ▶ Not a theory of inference ...
- ▶ ...but, under certain assumptions, coincides with maximum likelihood.

## A non-theory of inference: ordinary least squares (cont.)

- ▶ Not a theory of inference ...
- ▶ ...but, under certain assumptions, coincides with maximum likelihood.
- ▶ Bonus: easy to implement.

## Regression model notation

## Regression model notation

- ▶ Notation:  $Y \sim \mathcal{N}(y \mid \mu, \sigma^2)$ .

# Regression model notation

- ▶ Notation:  $Y \sim \mathcal{N}(y \mid \mu, \sigma^2)$ .
- ▶ Equivalently:  $y = \mu + \epsilon$ .

# Regression model notation

- ▶ Notation:  $Y \sim \mathcal{N}(y \mid \mu, \sigma^2)$ .
- ▶ Equivalently:  $y = \mu + \epsilon$ .
- ▶  $y$ : outcome (dependent) variable.

# Regression model notation

- ▶ Notation:  $Y \sim \mathcal{N}(y \mid \mu, \sigma^2)$ .
- ▶ Equivalently:  $y = \mu + \epsilon$ .
- ▶  $y$ : outcome (dependent) variable.
- ▶  $\mu$ : intercept (mean).

# Regression model notation

- ▶ Notation:  $Y \sim \mathcal{N}(y \mid \mu, \sigma^2)$ .
- ▶ Equivalently:  $y = \mu + \epsilon$ .
- ▶  $y$ : outcome (dependent) variable.
- ▶  $\mu$ : intercept (mean).
- ▶  $\epsilon$ : error (disturbance) term, with  $\epsilon \sim \mathcal{N}(0, \sigma^2)$ .



## OLS regression in R

```
lm(lifexp ~ 1)
```

# OLS regression in R

```
lm(lifexp ~ 1)
```

```
##  
## Call:  
## lm(formula = lifexp ~ 1)  
##  
## Coefficients:  
## (Intercept)  
##      71.93
```

# OLS regression in R

```
lm(lifexp ~ 1)
```

```
##  
## Call:  
## lm(formula = lifexp ~ 1)  
##  
## Coefficients:  
## (Intercept)  
##      71.93
```

```
mean(lifexp)
```

```
## [1] 71.932
```